Pathology Day 3 Table of Contents

Pediatric Pathology I 2
Pediatric Pathology II 15
Pulmonary Pathology 33
Head and Neck Pathology 83
Bone Pathology...107
Soft Tissue Pathology 123
Liver Pathology 149
Inflammation and Repair 167

Pediatric Pathology – Part 1

Archana Shenoy, MD
Staff Pathologist and Assistant Professor
Nationwide Children's Hospital/ The Ohio State University
Columbus, OH

Outline

- Part 1: Pediatric surgical pathology
  - Neoplasms: Small round blue cell tumors
  - Non-neoplastic entities
- Part 2: Perinatal autopsy and Placental pathology
  - Congenital anomalies
  - Molar pregnancy

General scheme of discussion

- Gross and microscopic images
  (Recognize the diagnosis)
- Salient features and pathology
- Molecular attributes
- Prognostic factors

Small round blue cell tumor # 1



Slides courtesy: Anthony Yachnis, MD

Diagnosis: Retinoblastoma



Flexner–Wintersteiner rosettes

- Most common primary intraocular malignancy of childhood
- Loss or inactivation of both alleles of *RB1* (RB) gene
  - Located at 13q14
  - Tumor suppressor gene
- Knudson "2-hit" hypothesis
  - RB results from 2 independent mutations
  - 1st mutation may be either somatic (sporadic) or germinal (inherited)
  - 2nd mutation is sporadic
- Bilaterality 20% to 35%
- Most often presents with leukocoria, cat's eye reflex

Slides courtesy: Anthony Yachnis, MD

Small round blue cell tumor # 2



Flecks of calcification







Diagnosis: Neuroblastoma

- Immunohistochemical stains:
  - PHOX2B
  - PGP 9.5
  - Tyrosine hydroxylase
  - Synaptophysin
  - Chromogranin
  - Neuron specific enolase
- Urine catecholamines in > 90%
- Imaging: Metaiodobenzyl guanidine (MIBG) tracer scan



Peripheral neuroblastic tumors

Epidemiology

- Common sites: Adrenal gland and retroperitoneum, Paraspinal along sympathetic chain
- International neuroblastoma Pathology Committee Classification, INPC (Shimada classification)
- < 50% Schwannian Stroma: Neuroblastoma
- > 50% Schwannian Stroma:
  - Ganglioneuroblastoma, nodular:
    - Gross nodules present
  - Ganglioneuroblastoma, intermixed
  - Ganglioneuroma

Associations

- "Blueberry muffin baby"
  - Blue-red cutaneous metastases
- Myoclonus-opsoclonus syndrome
  - Rapid, alternating eye movements and myoclonic movements of extremities
  - Resolves with tumor eradication



Mitotic karyorrhectic index

Calculated per 5000 neuroblasts

Classified as:

- Low = <100 (<2%)
- Intermediate = 100-200 (2-4%)
- High = >200 (>4%)

INPC Classification: Neuroblastoma

No ganglionic differentiation or neuropil	<5% Ganglionic differentiation	>5% Ganglionic differentiation
Undifferentiated	Poorly differentiated	Differentiating



Undifferentiated

Age	MKI
Any	Any
Unfavorable histopathology	

Poorly differentiated

Age	MKI
<1.5 yrs	<4%
Favorable histopathology	

Differentiating

Age	MKI
<1.5 yrs	<4%
1.5-5 yrs	< 2%
Favorable histopathology	

All other ages and MKIs: Unfavorable histopathology

Prognostic factors

Favorable

- Age at diagnosis: < 1.5 years
- Favorable histology
- Stages 1, 2
- Stage 4S (Stage 1 or 2 with mets to skin, liver, &/or Bone marrow)
- High expression of TrKA (high-affinity nerve growth factor receptor)
- Normal serum ferritin, NSE, and LDH
- Urinary VMA:HVA ratio > 1.5

Unfavorable

- MYCN amplification
- Near diploidy or tetraploidy
- Loss of heterozygosity of 1p and 11q

Ahmed AA et. al. Pediatr Hematol Oncol. 2017 Apr;34(3):165-185. PMID: 28662353

GANGLIONEUROMA



Small round blue cell tumor # 3









Diagnosis: Nephroblastoma (Wilm's tumor)

- Most common renal tumors in children
- Immunohistochemical stains:
  - WT1 (nuclear)
  - CD56
- Syndromic associations:
  - Beckwith-Wiedemann syndrome
  - Perlman familial nephroblastomatosis syndrome
  - Denys-Drash syndrome
  - WAGR syndrome

    (Wilms tumor, aniridia, genitourinary anomalies, and mental retardation)
  - Trisomy 18
  - Neurofibromatosis
  - Bloom syndrome

Deletion or mutation of WT1: 11p13

Beckwith-Wiedemann syndrome

- Loss of heterozygosity or imprinting at 11p15 (locus called WT2)
- Congenital anomalies
  - Macroglossia
  - Macrosomia
  - Midline abdominal wall defects
    - Omphalocele
- Ear abnormalities
- Neonatal hypoglycemia
- Tumors:
  - Wilms tumor
  - Hepatoblastoma

Key aspects of specimen handling

- Weight: No chemotherapy in younger patients with stage I and total kidney weight <550gm
- Perihilar and perirenal lymph nodes needs to be sampled
  - Lack of lymph node evaluation excludes patients from "no-chemotherapy-required" approach
- Documentation of exact site is necessary
  - Critical for evaluating for focal versus diffuse anaplasia





Anaplasia (unfavorable histology):
Nuclear size 3x background
and atypical multipolar/polypoid mitosis

Prognostic factors

- High stage at presentation
- Diffuse anaplasia (unfavorable histology)
- Nephrogenic rests in resected kidney indicates greater probability of bilateral Wilms tumor
- Tumor-specific loss of heterozygosity in 1p and 16q: high risk of relapse and death



- Nephrogenic rests can be perilobar (Subcapsular) or intralobar
- Presence of tumor capsule is important in the distinction of nephroblastoma from a hyperplastic nephrogenic rest

Dome JS et. al. Am Soc Clin Oncol Educ Book. 2014:215-23 PMID: 24857079

Other renal tumors in children

1 day old boy kidney tumor





Dense cellularity, mitoses, and sarcomatous appearance



Intersecting bundles of spindle cells with minimal atypia and infrequent mitoses

Diagnosis: Congenital mesoblastic nephroma

- Demographics
  - Most common renal tumor of infancy
  - Excellent outcome with complete resection
  - Increased risk of local recurrence with incomplete excision
- Microscopic features:
  - Classic variant
  - Cellular variant : t(12;15)
  - Mixed: Shows combination of both histologic patterns



Fascicles of plump spindle cells

Cellular mesoblastic nephroma

ETV6 translocation +

Frequent mitoses

Soft tissue tumor with ETV6 translocation t(12;15) : Infantile fibrosarcoma

- Age: First year of life, sometimes congenital
- Site: Distal extremity soft tissue (two thirds)
  - Head and neck
  - Visceral (rare)
- Solitary rapidly growing mass, ulcerates overlying skin





Spindle cells with a “herring bone” pattern

Diagnosis:

Congenital / Infantile Fibrosarcoma

Mitotically active with little nuclear pleomorphism

Knezevich SR Nat Genet. 1998 Feb;18(2):184-7. PMID: 9462753.

Renal tumor .. cont’d..



Malignant rhabdoid tumor

- Predominantly presents in infancy
- Aggressive tumor
- Biallelic inactivation of gene, located at 22q11.2
  - ~1/3 germline
    - Rhabdoid tumor predisposition syndrome
- Extrarenal sites possible
  - Soft tissue
  - Liver

Clear cell sarcoma of kidney


Photo courtesy: Danielle Harrell, DO

- Presents around 2- 3 years of age
- Tumor necrosis: poor survival

- Nests and cords of cells separated by arborizing fibrovascular septa
- Septa may be thin and delicate or cellular

Next blastoma............







Diagnosis: Hepatoblastoma

Classification:

- Epithelial subtype
  - Fetal pattern
  - Embryonal pattern
  - Macrotrabecular pattern
  - Small cell undifferentiated type (subset have INI-1 loss)
  - Cholangioblastic pattern
  - Mixed
- Mixed epithelial-mesenchymal subtype
  - Stromal derivatives
    - Osteioid
  - Teratoid
    - Melanin, squamous or glandular structures

Syndromic associations:

- Familial adenomatous polyposis of colon
- Beckwith-Wiedemann syndrome
- Li-Fraumeni syndrome
- Trisomy 18

Molecular alteration:

-β-catenin mutation-associated Wnt pathway activation

Immunohistochemistry:

Beta-catenin, AFP, Glypican-3

Prognostic factors:
Stage is most important(completely resectable- good); Pure fetal histology (good); Small cell undifferentiated (Poor)



Hepatoblastoma



Slide courtesy of Nicoleta Arva, MD/PhD

Mesenchymal hamartoma



- First year of life
- Multi-loculated cystic mass, well demarcated grossly.



Pancreatoblastoma

- Most frequent pancreatic tumor of childhood
- Rarely occurs in adults
- Associated with Beckwith-Wiedeman
- Epithelial neoplasm with multiple lines of differentiation
- Large masses, usually encapsulated
- Abnormalities of APC/β-catenin pathway demonstrated
- Good response to chemotherapy, better than in adults
- Immunohistochemistry:
  - Trypsin, chymotrypsin
  - Alpha-fetoprotein



Small round blue cell tumor #4
34 month old female with vaginal mass







“Cambium”: Subepithelial epithelial hypercellularity

Arrows: strap cells



Diagnosis:
Embryonal rhabdomyosarcoma, botryoid variant

Desmin



Myogenin





17 year old maxillary mass



Giant cells



Final diagnosis: Alveolar rhabdomyosarcoma



Myogenin: Diffusely positive

Rhabdomyosarcoma - Demographics

Embryonal

- Commonest pediatric ST sarcoma
  - Most <10years, 50% <5years, 5% <1year
- 80% of all RMS
- Involve wide range of sites
  - 50% head and neck
  - 30% GU and paratesticular
  - Rarely arise in extremities
  - Very rarely involve other sites

Alveolar

- Approx 20% of all RMS
- Rapidly growing mass lesion
- Involves any site
  - 40% extremities
- May present with disseminated disease

Rhabdomyosarcoma - histology

Embryonal

- Sheets of rhabdomyoblasts
  - Immature round to ovoid to spindle cells
  - Differentiating strap or spider cells
- Variably cellular/loose stroma
- Histologic variants include:

Spindle cell variant

Botryoid variant

Anaplastic/pleomorphic variant

Alveolar

- Round cells with variably fibrous stroma
- Giant cells prototypical

Classical alveolar pattern

Solid variant

Rhabdomyosarcoma:Ancillary studies

Embryonal

- Immunohistochemical stains:
  - Desmin
  - Myogenin: >90% specific, usually stains <30% of cells
  - MyoD1
- Cytogenetics
  - 11p15 loss well-reported
  - Not diagnostic

Alveolar

- Immunohistochemical stains:
  - Desmin
  - Myogenin: usually stains >90% of cells
  - MyoD1
- Cytogenetics:
  - FHKR (FOXO1) translocation: diagnostic
    - t(2;13)(q35;q14) PAX3-FHKR (~60%)
      - More aggressive
    - t(1;13)(p36;q14) PAX7-FKHR (~20%)
- Detectable by FISH or RT-PCR, Fusion Panel



- Spindle cell rhabdomyosarcoma
- NCOA2 Fusion detected
- Favorable in pediatric age group



Adult spindle cell/ sclerosing rhabdomyosarcoma
- Recurrent MYOD1 mutations in a subset
- Relatively worse prognosis

Ewing Sarcoma (formerly Peripheral neuroectodermal tumor in soft tissue)

- Bone: Painful, enlarging mass in diaphysis of long bones
- Balanced translocation t(11;22)(q24;q12) involving the EWS gene on chromosome 22 and the FLI gene on chromosome 11
  - 85% have t(11;22)(q24;q12) the EWS/FL11
  - 5-10% have t(21;22)(q22;q12) the EWS/ERG
- Homer-Wright (Pseudo) Rosettes
- Immunohistochemical stains: CD99


CD99: Strong diffuse membranous pattern

Immunohistochemical stain: NKX2-2

- Nuclear stain
- Helps distinguish Ewing Sarcoma from mimickers



Hung YP et. al. Mod Pathol. 2016 Apr;29(4):370-80. PMID: 26847175.

NON- NEOPLASTIC

Diagnosis?



Diagnosis: Hirschsprung Disease (Aganglionosis)

- Characterized by an absence of intramural parasympathetic ganglion cells in the distal gastrointestinal tract resulting in persistent contraction of the affected segment and subsequent colonic obstruction.
- 70%, isolated ; 30%, associated with:
  - Abnormal karyotype 12%; i.e. Down's syndrome
  - Congenital anomalies 18%
- Histologic features
  - Aganglionosis
  - Submucosal nerve hypertrophy (> 40 microns)

Kapur RP. Hirschsprung disease. PathologyOutlines.com website. http://www.pathologyoutlines.com/topic/colonhirschsprung.html. Accessed September 22nd, 2020.

Calretinin

Acetyl choline esterase (frozen section)



Meckel's diverticulum

- Most common diverticulum and most common congenital anomaly of GI tract
- The vitelline duct is usually obliterated by the 10th week of embryonic life; in 2% of the population, it may persist in various forms

- Protrusion of intestine on antimesenteric border of terminal ileum



- 50% contain heterotopic gastric mucosa



Necrotizing enterocolitis

- Most common surgical emergency in the newborn period
- Incidence
  - From 1-2.4 per 1,000 live births in USA
  - Infants < 1,000 g have highest incidence (9.5/1,000 live births)
  - Incidence inversely proportional to gestational age
- Complications: Bowel perforation, sepsis and death
- Signs:
  - Abdominal distension
  - Absent bowel sounds
  - Vomiting, and diarrhea
  - Radiographic findings: Ileus and pneumatosis intestinalis (air in bowel wall)





Abetalipoproteinemia









Acanthocytosis

- Autosomal recessive
- Defect in gene coding microsomal triglyceride transport protein; Chr 4q22
- Irregular vacuoles
- Non-membrane bound
- Absence of apo-B lipoproteins

Slide courtesy: Pierre Russo MD

Severe diarrhea of early infancy

	Microvillous inclusion disease	Tufting enteropathy	Enteroendocrine cell dysgenesis	Autoimmune enteropathy
Presentation	First 2 weeks	First 2 weeks	First 2 weeks	After 1 month
Gene defect	MYO5b (18q21)	EpCAM (2p21)	NEUROG 3 (10q21.3)	FOXP3 (Xp11.23) in IPEX syndrome
Extraintestinal disease	Low GGT cholestasis post bowel transplantation	Dysmorphism; keratitis arthritis	Insulin-dependent diabetes	Polyendocrinopathy
Villous atrophy	Yes	variable	variable	variable
Surface epithelium	Absent brush border	Tufting and desquamation	normal	Normal or atrophic
Salient features	CD10, PAS abnormal	EpCAM IHC Negative	Negative chromogranin	Anti-enterocyte antibodies







Extrahepatic biliary atresia



Atresia: Luminal obliteration



- Portal expansion fibrosis
- ^ Bile duct proliferation
- * Bile stasis in bile ducts
- Changes may be subtle < 1 month

Biliary Atresia – the explant



Trichrome stain



CIRRHOSIS WITH BILIARY PATTERN

Questions?

14 year old presents with a testicular mass. Serum AFP is normal. The patient undergoes a radical orchiectomy. What is the diagnosis?

- A. Seminoma
- B. Yolk sac tumor
- C. Embryonal rhabdomyosarcoma
- D. Leydig cell tumor



What is the likely immunohistochemical profile?

- A. SALL 4 positive, Cytokeratin positive
- B. Oct 4 Positive, SALL 4 positive
- C. Desmin Positive, Myogenin Positive
- D. Inhibin Positive, Cytokeratin negative



A 3 year old presents with a posterior knee mass, that is resected

A. Myxoid liposarcoma
B. Lipoblastoma
C. Lipoma
D. Hibernoma



What is the most likely molecular alteration?

A. t (12:16), FUS DDIT3 fusion
B. Rearrangement of 8q11-13 (PLAG1)
C. MDM2 amplification
D. Rearrangement of 11q13



Acknowledgements

- Previous lecturers:
  - Jamie Shutter, MD (Dr. Kumar and Elsevier for permission to use their images and diagrams)
  - Nicoleta Arva MD/PhD
  - Raja Rabah MD

This page was intentionally left blank.

Pediatric Pathology – Part 2

Archana Shenoy, MD
Staff Pathologist and Assistant Professor
Nationwide Children's Hospital/ The Ohio State University
Columbus, OH

Outline

- Part 1: Pediatric surgical pathology
  - Neoplasms: Small round blue cell tumors
  - Non-neoplastic entities
- Part 2: Perinatal and Placental pathology
  - Congenital anomalies
    - Trisomies
    - Heart
    - Lung
    - Kidney
  - Placental pathology
  - Molar pregnancy

TRISOMIES

Trisomy 21



Slide courtesy of Nicoleta Arva, MD/PhD

Trisomy 21 (Down syndrome)

- Hypoplastic maxilla
- Slanting palpebral fissures
- Abundant nuchal skin
- Hypotonia
- Absent Moro reflex



Slide courtesy of Nicoleta Arva, MD/PhD

Trisomy 21

- Primary trisomy (95%); translocation (4%); mosaicism (1%)
- Hematologic abnormalities
  - Increased risk of acute leukemia (ALL/AML)
- Endocrine disorders
- Alzheimer's-like dementia > 40 years
- Other associated tumors
  - Retinoblastoma, testicular tumors, brain tumors
- Other associated disorders
  - Duodenal obstruction/atresia; imperforate anus; congenital megacolon

Slide courtesy of Nicoleta Arva, MD/PhD

Trisomy 21


ATRIOVENTRICULAR CANAL DEFECT
(Endocardial Cushion Defect, Atrioventricular Septal Defect)
NORMAL
Atrial Septum
Right Atrium
Tricuspid Valve
Right Ventricle
Left Atrium
Mitral Valve
Left Ventricle
Ventricular Septum
DEFECT
A.V. Canal Defect
Common Valve

Slide courtesy of Nicoleta Arva, MD/PhD

Trisomy 13 (Patau syndrome)

- Features:
  - Cleft lip/palate
  - Hypertelorism
    - Increased distance between the eyes
  - Proboscis
    - Elongated nose
  - **Polydactyly**
  - Midline scalp defects
  - Microphthalmia
    - Small eyes
  - Sloping forehead
  - Flat low-set ears
- Associated tumors
  - AML, Neuroblastoma, Adrenocortical carcinoma, retinoblastoma [del(13q14)]



Slide courtesy of Nicoleta Arva, MD/PhD


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
X
Y

Slide courtesy of Nicoleta Arva, MD/PhD

Trisomy 18 (Edward syndrome)


1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
X
Y

Slide courtesy of Nicoleta Arva, MD/PhD

Trisomy 18

- Short sternum
- Narrow pelvis
- Clenched fists
  - Overlapping 2nd & 5th fingers
- Rocker bottom feet
- Short dorsiflexed first toe
- Partial syndactyly of toes
- Upturned nares



Slide courtesy of Nicoleta Arva, MD/PhD

Trisomy 18

- Prominent occiput
- Low set ears with pointed upper helix
- Hypertelorism
- Small palpebral fissure
- Small mouth
- Micrognathia



Slide courtesy of Nicoleta Arva, MD/PhD

Klinefelter syndrome
(Incidence 1:850 live male births)

47, XXY males (80%)
- 46, XY/47 XXY mosaicism (10%)
- 48, XXYY and others (10%)
- Testicular abnormalities
  - Approximately 10% of male sterility due to KS
- Renal abnormalities
- Gynecomastia

Slide courtesy of Nicoleta Arva, MD/PhD

Klinefelter syndrome


Tall stature
Poor beard growth
Minor breast development
Testicular atrophy
Female pubic hair pattern
1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
X
X
Y
Sex Chromosomes

(a) Klinefelter Syndrome (47,XXY)

Slide courtesy of Nicoleta Arva, MD/PhD

Turner Syndrome(45, X)


Low posterior hairline
Webbing of neck
Broad chest and widely spaced nipples
Pigmented nevi
Short stature
Coarctation of aorta
Cubitus valgus
Streak ovaries, infertility, amenorrhea
Peripheral lymphedema at birth

TURNER SYNDROME

Incidence: 1 in 3000 female births
Karyotypes:
  Classic: 45,X
  Defective second X chromosome: 46,X,i(Xq)
    46,X,del(Xq)
    46,X,del(Xp)
    46,X, r(X)
  Mosaic type: 45,X/46,XX
    45,X/46,XY
    45,X/47,XXX
    45,X/46,X,i(X)(q10)

Kumar et al: Robbins & Cotran Pathologic Basis of Disease, 8th Edition.
Copyright © 2009 by Saunders, an imprint of Elsevier, Inc. All rights reserved.

Downloaded from: StudentConsult (on 23September 2013 09:35 PM) © 2005 Elsevier

Turner syndrome (45,X)

Preductal coarctation of aorta



Single umbilical artery



Slide courtesy of Nicoleta Arva, MD/PhD

Turner syndrome (45,X)
Cystic hygroma

Prenatal fluid accumulation in lymphatics of neck





Slide courtesy of Nicoleta Arva, MD/PhD

Other Common Anomalies





Slide courtesy of Nicoleta Arva, MD/PhD

Omphalocele vs Gastroschisis

Omphalocele

- Large umbilical cord defect
  - Intestines protrude
- Intestines covered by a membranous sac
  - Amnion and parietal peritoneum
- Often associated with other anomalies
- 15% have chromosomal anomalies



Gastroschisis

- Paraumbilical abdominal wall defect
  - Extruded free loops of bowel not covered by sac
- Failure of normal umbilical cord development



Slide courtesy of Nicoleta Arva, MD/PhD

Congenital diaphragmatic hernia

- Congenital defect within the diaphragm
- Location:
  - –Left posterolateral (> 80%)
  - –Less common: anterior/ right
- Major complication
  - –Pulmonary hypoplasia
  - –Incarceration and necrosis of herniated organs



CONGENITAL LUNG DISEASE: DIFFUSE LUNG DISEASE

#1



Alveolar capillary dysplasia with venous malalignment

- Clinical background
  - –Term or near term infant
  - –Respiratory failure within days of birth
  - –Persistent pulmonary hypertension
  - –No sustained response to therapy
  - –Death within 1 month of life
  - –Additional anomalies common (~1/2 of cases)
    - Heart, GI, GU
- Microdeletions in the FOX gene cluster on 16q24.1 and mutations in the FOXF1 gene, a transcription factor that is involved in organogenesis
- Histology
  - –Variably deficient alveolar development
  - –Deficient capillary development
  - –Malpositioned pulmonary veins/venules
  - –Thickened pulmonary arteries/arterioles
  - –Muscularized intra-acinar pulmonary arterioles
  - –Pulmonary lymphangiectasis (1/3 cases)



The Osler Institute www.osler.org







Misalignment of pulmonary veins

#2



Surfactant Disorders

- Pulmonary surfactant:
  - Mixture of lipids and proteins
  - Reduces alveolar surface tension
  - Prevents end-expiratory atelectasis
  - Produced and stored by alveolar type II cells
- Surfactant deficiencies (multiple mutations):
  - Surfactant protein B
  - Surfactant protein C
  - ABCA3

Surfactant Dysfunction Disorders

- Pulmonary alveolar proteinosis:
  - SpB genetic mutations
  - ABCA3 genetic mutations
- Chronic pneumonitis of infancy:
  - SpC genetic mutations





The Osler Institute www.osler.org

CONGENITAL LUNG DISEASE: LOCALIZED CYSTIC LESIONS

Describe or diagnose



Cystic pulmonary airway malformation (CPAM)

- Most common surgically resected pulmonary malformation in children
- Types: CPAM 1 – 3 based on cyst size and morphology
- CPAM type 1: Associated with mucinous adenocarcinoma
- Complications:
  - Pressure effects
  - Repeated infections



- Cysts >2 cm
- Surrounding smaller cysts

- Cysts <2 cm
- Intervening alveolar parenchyma

Solid/adenomatoid type

Adapted from Stocker JT et al. Congenital cystic adenomatoid malformation of the lung. Classification and morphologic spectrum. Hum Pathol 1977;8:155-171.

Cystic pulmonary airway malformation 1





CPAM type2





Striated muscle fibers



Describe or diagnose

Intralobar sequestration

- Definition: portion of parenchyma within the lung itself with a systemic arterial supply and lack of connection with the remainder of the lung (bronchial atresia).
- Location: lower lobe
- Arterial supply:
  - Most commonly aorta
  - Occasionally celiac, splenic, intercostal, subclavian
- Venous drainage:
  - Most commonly pulmonary
  - Occasionally systemic



Extralobar sequestration (ELS)

- Isolated mass of lung tissue with its own pleural investment, separate from the normal lung
- Systemic arterial supply
  - –most commonly descending thoracic aorta
- Variable venous drainage:
  - –Systemic or pulmonary
- Most common location:
  - –Between lower lobe and diaphragm (~90% left)
- Other locations:
  - –Anterior or posterior mediastinum, within diaphragm, intra-abdominal, retroperitoneal
  - –May be associated with ipsilateral diaphragmatic defect



Bronchogenic cyst

- Solitary, unilocular
- Etiology: accessory foregut bud
- Usually attached
  - –trachea/major bronchi
- Most commonly within mediastinum, above carina
  - –any paramidline site from suprasternal to infradiaphragm
- May cause secondary obstruction



CONGENITAL HEART ANOMALIES

Tetralogy of Fallot

- Most common cyanotic congenital heart disease
- Right to left shunt
  - –Overriding aorta
  - –Pulmonary valve stenosis
  - –Ventricular septal defect
  - –Right ventricular hypertrophy
- Majority of the blood bypasses lungs
  - –Hypoxia and cyanosis
- Tet spells
  - –Squatting to change shunt direction


Normal heart
Tetralogy of Fallot
Overriding aorta
Pulmonic stenosis
ventricular septal defect
Right ventricular hypertrophy

Slide courtesy of Nicoleta Arva, MD/PhD

Transposition of the great vessels-
Complete, dependent on VSD


Normal Heart
Transposition of the Great
SVC
Ao
MPA
LA
RA
LV
RV
IVC
Ao
SVC
Ao
MPA
LA
RA
LV
RV
IVC
Ao
KEY
SVC – Superior Vena Cava
IVC – Inferior Vena Cava
RA – Right Atrium
RV- Right Ventricle
MPA – Main Pulmonary Artery

Slide courtesy of Nicoleta Arva, MD/PhD

Transposition of the Great Vessels


Ao
LA
RA
RV
LV
A Classic Tetralogy of Fallot
Ao
PT
LA
RA
LV
RV
Ao
PT
LA
RA
LV
RV
With VSD
Without VSD
B Complete Transposition

Kumar et al: Robbins & Cotran Pathologic Basis of Disease, 8th Edition.
Copyright © 2009 by Saunders, an imprint of Elsevier, Inc. All rights reserved.



Kumar et al: Robbins & Cotran Pathologic Basis of Disease, 8th Edition.
Copyright © 2009 by Saunders, an imprint of Elsevier, Inc. All rights reserved.

Slide courtesy: Jamie Shutter, MD

Coarctation of the Aorta

- Pre-ductal
  - –Usually detected in infancy
  - –Heart failure, weak pulses & cyanosis of lower extremities
  - –Associated with Turner syndrome
- Post-ductal
  - –Usually present in older children or adults
  - –Hypertension
    - •Upper extremities
  - –Hypotension
    - •Lower extremities

Coarctation of the Aorta


Ao
PT
LA
RA
RV
LV
Ao
PT
LA
RA
RV
LV
With PDA
Without PDA

Coarctation of Aorta

Kumar et al: Robbins & Cotran Pathologic Basis of Disease, 8th Edition.
Copyright © 2009 by Saunders, an imprint of Elsevier, Inc. All rights reserved.

Downloaded from: StudentConsult (on 23 September 2013 09:49 PM)

Slide courtesy: Jamie Shutter, MD

© 2005 Elsevier

RENAL ANOMALIES

AUTOSOMAL RECESSIVE POLYCYSTIC KIDNEY DISEASE

Mutations in PKHD1 gene at 6p12
Early mortality common from pulmonary hypoplasia


Gross
Microscopy
Medullary cysts
“Medullary rays”

Congenital Hepatic Fibrosis: Ductal Plate Malformation

- Constant feature of ARPKD
- Rarely isolated
- Progressive fibrosis of portal tracts





AUTOSOMAL DOMINANT POLYCYSTIC KIDNEY DISEASE



- PKD 1, 16p13.3(85%), polycystin 1
- PKD 2, 4q13-q23, polycystin 2
- Complications: Intracranial and other vascular aneurysms


Glomerular and tubular cysts

Multicystic renal dysplasia


Mesodermal collarettes
Cartilage
200 µm

Oligohydramnios sequence (Potter sequence)


Flat face, beaked nose, low-set ears
Limb deformation
Pulmonary hypoplasia

PLACENTAL PATHOLOGY

GROSS OBSERVATIONS


MARGINAL
CORD INSERTION


VELAMENTOUS INSERTION


VELAMENTOUS INSERTION,
HYPERCOILED CORD
Normal: 2- 3 twists/ 10 cm
"FETUS PAPYRACEUS"


TRUE
CORD
KNOT

MICROSCOPIC FINDINGS

Amniotic Fluid Infection

Maternal response: Membranes



Fetal response: Umbilical cord/chorionic vessel



Grade 1: Mild – moderate
Grade 2: Severe

Stage 1—acute subchorionitis or chorionitis
Stage 2—acute chorioamnionitis: polymorphonuclear leukocytes extend into fibrous chorion and/or amnion
Stage 3—necrotizing chorioamnionitis: karyorrhexis of polymorphonuclear leukocytes, amniocyte necrosis, and/or amnion basement membrane hypereosinophilia

Stage 1—chorionic vasculitis or umbilical phlebitis
Stage 2—involvement of the umbilical vein and one or more umbilical arteries
Stage 3—necrotizing funisitis

Usual organisms: Group B streptococcus, Klebsiella, E.Coli

Candida “peripheral funisitis” ; hyphae are seen (GMS stain)

Khong TY, Arch Pathol Lab Med. 2016 Jul;140(7):698-713. Epub 2016 May 25. PMID: 27223167.

Microscopic pathology: Maternal inflammatory response



Stage 1: Acute subchorionitis /chorionitis

Stage 2: Acute chorioamnionitis / chorionitis involving fibrous chorion

Stage 3: Necrotizing chorioamnionitis

Candida Funisitis



Picture courtesy: Selene Koo, MD PhD





Chronic villitis (villitis of unknown etiology)





Khong TY, Arch Pathol Lab Med. 2016 Jul;140(7):698-713. Epub 2016 May 25. PMID: 27223167.

Classification of Villitis



Chronic villitis
- Low-grade (<10 villi)
  - Focal – one slide
  - Multifocal - multiple
- High-grade (>10 villi)
  - Patchy – multiple foci
  - Diffuse – >30% of distal villi

Khong TY, Arch Pathol Lab Med. 2016 Jul;140(7):698-713. Epub 2016 May 25. PMID: 27223167.

Lymphoplasmacytic villitis: Cytomegalovirus


Fetus:
Periventricular calcifications



Maternal hypertension: Atherosis (decidual vasculopathy)


Basal plate
Fetal membranes

Placental villous infarction


Ghost villi,
back to back

Fetal vascular malperfusion (previously fetal thrombotic vasculopathy)


Normal villus
Cluster of
avascular villi
50 µm

Abruption

Organizing retro-placental thrombus with underlying villous infarction



Chorangioma


Vascular lesion
Normal villi
100 µm

Multiple – chorangiomatosis
Differentiate from chorangiosis – Increased vascularity within terminal villi

Chorangiosis/Chorangiomatosis



Chorangiosis

"Involves terminal villi"



Localized chorangiomatosis

Twin membranes


Dichorionic diamnionic
Monochorionic diamnionic
Chorion "sandwich"

Twin-twin transfusion syndrome

Complication of monochorionic gestation with vascular anastomosis

"Recipient" – Polyuric polyhydramnios


Twin 1
Twin 2
100 µm

"Donor" -- Oliguric oligohydramnios the "stuck twin"

Possible recipient: Appropriately mature; Possible donor: Atrophic (more mature/shriveled)

TRAP (Twin reversed arterial perfusion) sequence→ Acardiac twin



Molar Gestation


COMPLETE MOLE
Two sperm (XX or XY)
+
No Maternal DNA
Egg
or
X sperm
+
No Maternal DNA
Egg
Duplication of sperm DNA
46XX or 46XY
Paternal chromosomes only (Androgenetic mole)
PARTIAL MOLE
23X
23Y
One or two sperm
46XY
+
23X
Egg
23X
23X
23Y
69XXY or 69XXX
Maternal and paternal chromosomes (Triploid)
© Elsevier 2005

Slide courtesy: Jamie Shutter, MD

The Osler Institute www.osler.org

Molar Gestation



Molar Gestation



Molar gestation



p57

- p57KIP2 gene
  - –Cell-cycle inhibitor
  - –Maternally transcribed
  - –Paternally imprinted
  - –Expressed
    - Maternal decidual tissue
    - Cytotrophoblast
    - Stromal cells of villi

Hydropic Abortus & Partial Mole(p57)



Kumar et al: Robbins & Cotran Pathologic Basis of Disease, 8th Edition.
Copyright © 2009 by Saunders, an imprint of Elsevier, Inc. All rights reserved.

Slide courtesy: Jamie Shutter, MD

Complete Mole: p57 negative



- Both X chromosomes Paternally derived
- No expression
  - Cytotrophoblast
  - Stromal cells of villi

p57 positive in maternal decidua (internal control)

Mole mimic: Placental mesenchymal dysplasia (large > 1 kg)



Abnormal large villi, No trophoblast hyperplasia

Normal villi

Ernst, L.M. Placental Mesenchymal Dysplasia. *J. Fetal Med.* **2,** 127–133 (2015).

The Osler Institute www.osler.org

Placental mesenchymal dysplasia



Question 1



Which of the following syndromes is the most likely diagnosis:

A. Turner syndrome
B. Down syndrome
C. Klinefelter syndrome
D. Patau syndrome



a. Juvenile Xanthogranuloma
b. Langerhans cell histiocytosis
c. Glycogen storage disease
d. Wilson disease

Question 2: What is the diagnosis?



Langerhan's cell histiocytosis

- Clonal proliferation of Langerhans cells (modified dendritic cells)
- Skin, bone and bone marrow, and lung
  - Bone: Eosinophilic granuloma
  - Hand-Schüller-Christian disease: Multi-organ involvement and diabetes insipidus
  - Letterer-Siwe disease aggressive with multisystem involvement
- May spontaneously resolve or progress
- Immunohistochemistry:
  - CD1a, S100, Langerin/CD207



SLIDE QUIZ

Juvenile Xanthogranuloma


CD1a negative

Touton giant cells

Infantile hemangioma



Inclusion body fibromatosis (also infantile digital fibromatosis)


Trichrome (paranuclear eosinophilic globular inclusions)

Calcifying aponeurotic fibroma


Calcific deposits

Fibrous areas

Mononuclear cells and giant cells

Fibrous hamartoma of infancy


Fat

Immature mesenchyme

Fibrous areas

Yolk Sac tumor (elevated AFP)


Hyaline globules



Diagnosis?



Gonadoblastoma

- Occur in:
  - 46XY pure gonadal dysgenesis or
  - mixed gonadal dysgenesis associated with 45X (mosaic turner) or 46XY karyotype
- Immature Sertoli and granulosa cells are arranged around pink hyaline bodies, germ cells, or rim periphery of cellular nest
- Calcifications often present

Acknowledgements

- Previous lecturers:
  - Jamie Shutter, MD (Dr. Kumar and Elsevier for permission to use their images and diagrams)
  - Nicoleta Arva MD/PhD
  - Dr. Raja Rabah MD

This page was intentionally left blank.

Pulmonary Pathology

Vijayalakshmi Ananthanarayanan, MD
Associate Professor
Residency Program Director
Loyola University Medical Center
vananth@lumc.edu
@vijuananthan

- This lecture was modified from a previous lecture given by Aaron Auerbach, MD, MPH

Overview

- Anatomy and Histology
- Infections
- ILD
- Vasculitis and pulmonary hemorrhage syndromes
- Vascular disorders
- LAM, LCH
- Granulomatous lesions
- Tumors
- Additional reading material
- General format: Pictures followed by diagnosis and description

Normal Lung

- Air enters the nasal cavities, moves past the larynx, down the tracheobronchial tree, and enters the alveoli where gas exchange occurs
- Arteries follow airways
  - dual arterial supply from bronchial artery and pulmonary arteries
  - So, infarctions rarely occur in the absence of significant cardiac output problems
- Right bronchus is more vertically oriented
  - frequent site of aspiration

Respiratory epithelium

- Normal epithelium of respiratory tract is pseudostratified ciliated columnar epithelium
- From sinuses to terminal bronchioles



This Photo by Unknown Author is licensed under CC BY-SA-NC

TERTIARY BRONCHUS AND BRONCHIOLE
Bronchiole lacks submucosal glands and cartilage



Bronchovascular bundle


Alveolar duct
Bronchiole
Pulmonary artery
Alveoli


Intralobular venule
espiratory bronchiole
erminal bronchiole
Alveoli and their blood capillaries
Lymphatics related to the pulmonary artery
Interlobular connective tissue
Interlobular lymphatic and lymphatics relate to the pulmonary ve
Lymphatics related to the bronchus
Interlobular lymphatics
branch ary vein
ronchus
Pulmonary artery

Lobular anatomy

- Lung is divided into lobular anatomy
- On the outside you have the pleural surface
- Interlobular septae are fairly inconspicuous,
  - Septae contain pulmonary vein and lymphatics
- Bronchovascular bundle includes bronchus, pulmonary artery and lymphatics
- Disease with lymphangitic distribution is in pleura, Bronchovascular bundle and in Septae

Cell types in alveoli


ALVEOLAR SPACE
Type II pneumocyte
Type I pneumocyte
Endothelium
Interstitial cell
CAPILLARY LUMEN
Type I pneumocyte
ALVEOLAR SPACE
Endothelium



EM of type II pneumocyte with surfactant (pathologyoutlines.com)

- Type I cells or squamous pneumocytes - gas exchange
- Type II cells or granular pneumocytes -secrete surfactant

Infections

Viral Infections



Image Courtesy Dr. Aliya N.Husain



Image Courtesy Dr. Aliya N.Husain



Intranuclear and intracytoplasmic inclusions

Image Courtesy Dr. Aliya N.Husain



Cytomegalovirus

- Most common viral organism causing PNA recognized by the pathologist
- Herpesvirus, DS DNA, icosahedral capsid, envelope 120-200 nm
- Large cells with intranuclear and intracytoplasmic inclusions
- Opportunistic infections: congenital, perinatal, hepatitis, post-transfusion, transplants, immunosuppressed
- To know that the infection is in the lung, you need lung bxs or BAL



Image Courtesy Dr. Aliya N.Husain



Tracheobronchitis: Large bronchi with necrosed epithelium



Herpes Simplex

- Two serotypes:
  - HSV-1 respiratory tract infection
  - HSV-2 genitourinary tract infection
- Herpesvirus, similar features to CMV
- Nuclear inclusions, multinucleated, marginated chromatin, molding
- Usually airway epithelium
- HSV PNA seen in neonatal period, as well as immunocompromised including burns patients
- Diffuse HSV appears to be acquired hematogenously from other sites





Measles

- Highly contagious, causes rubeola
- 1-2 wk. incubation, then fever, Kolpiks' spot
- Pulmonary complications include: secondary (bacterial) PNA, giant cell pneumonia
- Giant cell pneumonia
  - Warthin-Finkeldey cells, multinucleated giant cells
  - Nuclear and cytoplasmic inclusions (measles)
  - RSV is the other cause of giant cell pneumonia (only cytoplasmic inclusions)
  - Different from Giant cell interstitial pneumonia (see later)





Adenovirus

- Non-enveloped DS DNA, 70-90 nm
- Person-to-person spread
  - Epidemics in dense populations (military barracks, college dorms)
- Nuclear inclusions and smudge cells
  - Like herpes
- Bronchiolitis in children
- Can cause severe necrotizing bronchiolitis and pneumonitis in neonates and immunocompromised

Bacterial Pneumonias



Bacteria in middle surrounded by eosinophilic debris and neutrophils



Sulfur granule, amorphous, purple surrounded by acute bronchopneumonia



Filamentous bacteria in sulfur granules

Actinomyces

- Anaerobic gram-positive bacilli; found in mouth and nose
- Susceptible to lysozyme
- NOT partially acid fast
  - Unlike Nocardia (partially acid fast)
- Infection through aspiration or direct extension from abscess
- Granulomatous lesion with sulfur granules, abscesses and sinus tracts


Macrophages and neutrophils fill alveoli (non-specific)


No sulfur granules, filamentous organisms

Nocardia

- Aerobic gram-positive bacilli with filamentous beaded morphology
- Partial acid-fastness / Fite stain
  - Unlike Actinomyces
- Infection through inhalation of saprophytic organisms, often immunodeficient patients
- 50% disseminate to other organs
- Dx with culture (1-2 weeks)
  - Direct examination, biopsy or BAL, is quicker


PAS stain
Michaelis-Gutmann bodies
von Kossa stain

TARGET

This Photo by Unknown Author is licensed under CC BY-SA

Rhodococcus equi

- Aerobic Gram-positive/variable coccobacillus
- Infection through inhalation sometimes from animals (horses)
- Dx material from sputum, BAL, or infected tissue
- Most common cause of pulmonary malakoplakia
  - Chronic inflammatory process with a prominent histiocytic component and calcifications (Michaelis-Gutmann bodies)
    - Demonstrable with von Kossa stain


Bronchopneumonia with Michaelis-Gutmann bodies

The Osler Institute www.osler.org





Auramine-Rhodamine Immunofluorescence



© 2005 Elsevier Inc. Leslie & Wick: Practical Pulmonary Pathology

Fluorescent stains: light up and are easier to see

Mycobacterium tuberculosis

- Acid-fast bacillus – aerobic, non-motile, contain mycolic acid in their cell walls
- Chronic recurrent infection of the lung
- Inhaled small droplet nuclei that are effectively dispersed by coughing
- Primary and secondary tuberculosis

Primary TB

- Alveolar bacilli are ingested by neutrophils and macrophages - macrophages may go to regional lymph nodes
- Bacilli often lodge in posterior half of upper lobes due to the high oxygen tension
- When 1º infection resolves and forms a parenchymal nodule (Ghon focus)
- When Ghon focus is associated with calcified hilar nodes called a Ranke's complex
- > 90% of 1º TB heal without progression
  - More likely to progress in patients with impaired cellular immunity
  - Progressive infections result in necrotizing granulomas

Secondary TB

- Occurs when:
  - Dormant but viable mycobacteria becomes active
  - Previously infected individual is reinfected
- Usually cavitary lesion in one or both upper lobes
- Miliary TB results from hematogenous spread
  - Many nodules in lung, liver, etc..
- Most common form of clinical TB in adults

The Osler Institute www.osler.org





Non-TB Mycobacteria


Lady Windermere's Fan (1892).
A Play About a Good Woman is a four-act comedy.
Oscar Wilde

- M. avium-intracellulare, M. kansasii, M. fortuitum, and M. chelonei
  - Ubiquitous organisms; infection usually from environment
- Majority of patients have underlying lung disease
  - Often immunodeficient patients
  - Unsure of clinical significance of infection in immunocompetent patients; colonization vs infection
- Assoc with middle lobe syndrome (Lady Windermere syndrome)
  - Middle lobe has a steep angle so patients have trouble clearing secretions from that lobe
  - They get atypical MAI infections
- Assoc with hot tub lung
  - Especially if hot tub is indoors or if water is not changed

Fungal Infections















Aspergillus

- Ubiquitous molds found worldwide
- Septate and 45º branching hyphae
- Spectrum of clinical disease:
  - Fungus ball (aspergilloma)
    - Colonization of previous cavity
    - This is the most common form of aspergillus
  - Acute angioinvasive aspergillosis
  - Hypersensitivity reactions:
    - Allergic bronchopulmonary aspergillosis (ABPA)
    - Hypersensitivity pneumonitis
  - Chronic necrotizing aspergillosis (fungal ball that is starting to invade into tissue)

Allergic bronchopulmonary aspergillosis

- Hypersensitivity reaction to an Aspergillus antigen
  - usually in asthmatics
- Airways filled with allergic mucin
  - Tide mark lines
  - Collections of eosinophils
- Mucous
- Charcot-Leyden crystals (hexagonal)



Tide marks, septate hyphae and 45 angle septation

Peudoallescheria boydii

- Immunocompromised patients
- Associated with allergic bronchopulmonary disease
- Morphologically very similar to Aspergillus Sp
- Tide marks, septate hyphae and 45 angle septation







Irregular (different widths) ribbon-like broad non-septate hyphae

Zygomycetes

- Uncommon but frequently fatal infection
- Mostly Rhizopus, Mucor, Rhizomucor
- Transmitted by inhalation, ingestion, or percutaneously
- Usually immunosuppressed patients
- Histology:
  - Irregular ribbon-like broad (empty appearing) non-septate hyphae
  - 90 degree branching
- vascular invasion



GMS, 2-4 µm, budding yeasts

All same size, except the ones that are budding

Histoplasma

- Dimorphic fungi
  - Exists in soil in mycelial form
  - Mississippi and Ohio river valleys
- Infection through dust inhalation of dust
  - in bird droppings
  - infectivity determined by ability to convert to yeast form
- Pathogenesis similar to TB
  - Primary
  - Progressive disseminated
  - Chronic cavitary histoplasmosis

54-year old male, smoker, lung mass





Suppurative Granulomatous Inflammation









Blastomyces dermatitidis

3Bs

- Dimorphic – Inhaled into lungs
- Occurs primarily in US, around Great Lakes
- Disseminates preferably to skin and bone
- Acute deterioration can result in ARDS
- Mixed acute and granulomatous inflammation
- Large yeast with “broad-based” budding
- Characteristic internal nuclear structure



Coccidioides – Book image from : Non-neoplastic Pulmonary Pathology: Sanjay Mukhopadhyay



Coccidioides – Book image from : Non-neoplastic Pulmonary Pathology: Sanjay Mukhopadhyay

Coccidioides

- Dimorphic fungus
  - US southwest, grows in hot climate
  - AKA San Joaquin Valley Fever
- Often asymptomatic
  - 40% have flu-like illness for 1-3 wks
  - 20% have erythema multiforme or erythema nodosum
- Rare dissemination, usually immunocompromised
- In contrast to Histo and Blasto, serologic studies are very useful







Cryptococcus

- Found worldwide
  - soil with pigeon droppings
- Two major infections
  - Pulmonary
  - Cerebromeningeal (hematogenous spread from lung)
- Hosts can be immunocompetent, especially if large inoculum
- Patients with primary infections usually do not have symptoms

46-year old with CVID







GMS





Pneumocystis

- Formerly P. carinii now P. jirovecii
- Mostly immunosuppressed
- Probably causes subclinical pneumonia in childhood
  - 60-100% of kids have abs by 4 y/o
- Intra-alveolar frothy material with minimal inflammation
  - GMS stains show 7-8 micron non-budding cysts
  - Dented ping-pong balls
  - Some with central black dot
  - PAS negative
- Routine lab studies not helpful
- Dx by BAL, bronchial washings, sputum specimens and bx

	Blastomyces	Coccidioides	Cryptococcus	Histoplasma	Pneumocystis
Average Size	8-15 um	30-60 um	4-7um	3 um	4-6 um
Morphology on H&E	Round, thick Walled with basophilic nuclei	Round, large, thick-walled with basophilic endospores or empty	Pale, round, thin, cell wall with halo (mucinous capsule)	Not visible in granulomatous lesions	Not visible in granulomatous lesion Dot-like structures in frothy exudate
Morphology on GMS	Round, uniform yeasts	Large spherules with endospores	Round yeasts, varying sizes, fragmentation	Oval shaped, small yeasts	Indented, helmet shaped
Budding	Yes, broad based	NO	Yes, narrow based	Yes	NO

The Osler Institute www.osler.org

Parasitic Infections





Toxoplasma gondii

- Obligate intracellular protozoa
- Three forms:
  - Tachyzoite (invasion, proliferation)
  - Tissue cyst (reservoir for tachyzoites)
  - Oocyst (infectious after sporulation)
- Cat is reservoir
- 80-90% of healthy hosts have no sxs
- In cell-mediated immunity deficiency have fever, encephalitis, brain abscesses, pneumonitis
- Serologies with titers to prove infection



Strongyloides

- Skin is penetrated via contact with soil contaminated with larvae-containing feces
- Pass to lung, coughed up, swallowed, GI tract infection
- Can complete life-cycle entirely within human host – can persist for decades
- Can be disseminated in immunocompromised

The Osler Institute www.osler.org



Echinococcus

- Species: E. granulosus, E. multilocularis, E. vogeli
- Usually rural, associated with sheep, cows, dogs
- Liver most common site in adults; lungs in children



Longitudinal section; image courtesy, Dr. Madhavi Naik, Bangalore, India

Ferrari, P.A., Grisolia, A., Reale, S. et al. A rare case of human pulmonary dirofilariasis with nodules mimicking malignancy: approach to diagnosis and treatment. J Cardiothorac Surg **13,** 65 (2018). https://doi.org/10.1186/s13019-018-0750-5

Dirofilariasis


DPDx
Pulmonary Dirofilariasis
CDC
8 Migrate to head and mosquito's proboscis
1 Mosquito takes a blood meal (L3 larvae enter bite wound)
7 L3 larvae
9 Lesions in pulmonary vessels
6 L1 larvae
Dirofilaria immitis
5 Microfilariae pentrate mosquito's midgut and migrate to Malpighian tubules.
4 Mosquito takes a blood meal (ingests microfilariae)
2 Adults in pulmonary arteries.
3 Adults produce microfilariae that typically are found in peripheral blood.
Infective stage
Diagnostic stage



Question time

Morphological findings shown in the image are most compatible with which of the following clinical scenarios

a) 70-year old male with LVH, dyspnea on exertion, pink, frothy sputum
b) 55-year old male with HIV presenting with cough and dyspnea on exertion
c) 70-year old with lambda restricted multiple myeloma with an isolated mass on imaging
d) 65-year male with dyspnea on exertion and microscopic honeycombing on imaging



Art by
Dr. EM
Wojcik

Interstitial Lung Diseases



Serena Williams of the USA grimaces in pain after being injured during her match against Bojana Jovanovski during day four of the 2012 Brisbane International at Pat Rafter Arena on January 4, 2012 in Brisbane, Australia.

The 6 Patterns of Interstitial Lung Disease

1. Acute Injury Prototype: DAD	4. Alveolar Filling Prototype: OP
2. Fibrosis Prototype: UIP	5. Nodules Prototype: Wegener's
3. Cellular Interstitial Prototype: HSP	6. Minimal Changes Prototype: SAD







Diffuse alveolar damage

- Form of acute lung injury to endothelium and epithelium, progresses through sequential stages
- Exudative stage
  - damaged alveoli, surfactant loss, alveolar collapse
  - Forming hyaline membranes
    - composed of dead cells (nucleoplasm, cytoplasm, fibrin)
    - Eosinophilic material mashed up against the alveoli
- Organizing phase
  - Organizing pneumonia (fibroblasts and myofibroblasts) with interstitial fibrosis
- ~60% mortality

Etiologies

- Infection (viral inclusions, adenovirus)
  - Eosinophilic pneumonia
- Drug toxicity
- Connective tissue disease
- Diffuse alveolar hemorrhage
- Irradiation injury
- Idiopathic (AIP)

Acute Interstitial Pneumonia

- AIP - idiopathic form of DAD - called Hamman-Rich syndrome
- Very rare, must rule out infection, drug toxicity, CVD and pancreatitis
- Chest X-ray shows diffuse ground glass infiltrates to complete “white out”
- Histological exam demonstrates hyaline membranes, interstitial and alveolar edema in acute phase Organizational phase displays organizing pneumonia.
- If resolves – can return to normal. More often, fibrosis develops with eventual honeycomb change



Organizing pneumonia

- New fibrosis
  - follow distal airways
  - spindled myofibroblasts and fibroblasts within loose collageous tissue
- Not old dense collagen fibrosis



Organizing Pneumonia

- Polypoid plug of fresh fibrous tissue
- Nonspecific reaction to numerous injuries
- Not to be confused with bronchiolitis obliterans









Cryptogenic Organizing Pneumonia

- Formerly “broncholitis obliterans organizing pneumonia”
    - BOOP
- Classified as interstitial lung disease but is really an airway disease and causes obstructive lung disease
- COP is idiopathic, but other entities can cause COP like changes in the lung
    - Characteristically develops 4-6 weeks after viral URI
- COP is a clinical diagnosis in which imaging shows ground glass opacities

COP

- Histologic features:
    - Patchy airway centered fibrous polyps of organizing pneumonia
    - Normal lung architecture – no significant fibrosis
    - No granulomas or acute or chronic inflammation
    - Must rule out infection (special stains)
- Treatment
    - Most resolve without treatment
    - Or steroids, very good prognosis


ABNORMAL LUNG
NORMAL LUNG




UIP
Patchy fibrosis, subpleural, paraseptal- Dr. Churg's atlas of pulmonary pathology
00--
-35803


Patchy fibrosis, subpleural, paraseptal- Dr. Churg's atlas of pulmonary pathology


Microscopic Honeycombing


Old fibrosis
New fibrosis (fibroblast foci)



Usual interstitial pneumonia

- Clinical term is idiopathic pulmonary fibrosis
- Diffuse, bilateral lung disease
- Poor prognosis
  - survival ≤ 5years
  - risk of lung cancer
- Unknown etiology
  - ? cigarette smoking
- Treatment
  - Drugs don't work
  - Lung transplant in young patients
- Characteristic clinical, radiologic, and histologic features
- Pattern can be seen in connective tissue disease, asbestosis and drug reactions

Histologic features

- Geographic and temporal heterogeneity
- Geographic heterogeneity
  - Patchy lung involvement
  - Periphery of lung and lower lobes i.e. subpleura and paraseptal
  - ADJACENT NORMAL LUNG TISSUE
- Temporal heterogeneity
  - Dense fibrosis with remodeling and usually honeycombing
  - Fibroblastic foci at edge of fibrosis
- Inflammation, smooth muscle proliferation, fatty metaplasia

COP vs fibroblast foci

Granulation tissue	Granulation tissue
In airways, bronchi	Not in airways
Branching	No branching
No fibrosis	Sits on fibrosis





Nonspecific Interstitial Pneumonia

- Uniform septal thickening
- Usually no honey combing/architectural distortion
- Fibrotic and cellular patterns
- Causes include CVD, drugs, infection, pneumoconiosis
- Mimickers include hypersensitivity pneumonitis and lymphoma

NSIP

- Cellular Pattern
  - Interstitial infiltrates of lymphocytes and plasma cells
  - Initially no architectural remodeling

  Fibrotic pattern
  - Fibrosis in each alveolar septae
  - Fibrosis blends gradually with the normal lung
  - Fibroblastic foci and granulomas are rare to absent
- Responds better to steroids than UIP
- Cellular-NSIP better prognosis than fibrotic-NSIP
  - Fibrotic-NSIP with honeycomb changes is similar to UIP



All alveoli are filled with infiltrate (like cellular NSIP)



Interstitial inflammation, nodular to diffuse lymphoplasmacytic infiltrate

Lymphocytic Interstitial Pneumonia

- Bilateral diffuse lymphoplasmacytic infiltrates
  - Overlap with cellular NSIP
  - More dense than NSIP
- Assoc with HIV, CVD (Sjögren's syndrome) , EBV, Pneumocystis
- Small % develop extranodal marginal zone lymphoma
- Interstitial inflam, nodular to diffuse lymphoplasmacytic infiltrate
  - Some use the term follicular bronchitis if the infiltrate is nodular with germinal centers and LIP if it is diffuse
- Fibrosis & honeycombing can develop later

Pleuritis



Pleura with lymphocytes and plasma cells



RA - Follicular lymphoid hyperplasia

http://granuloma.homestead.com/rheumatoid_nod_s89-2617-5.jpg


Central
necrotic
nodule
Palisaded
Histiocytes







Hypersensitivity Pneumonitis

- Reaction to inhaled antigen that results in chronic bronchocentric inflammation
- Acute, subacute (intermittent) or chronic
  - Acute forms are usually not biopsied
- Acute reactions due to type III hypersensitivity; chronic reaction due to type IV hypersensitivity
- Dyspnea, fever, malaise, weight loss, clubbing
- > 300 antigens have been described.
  - bird peptides (especially parakeet)
  - fungi (in moldy hay, dust, hot tubs, humidifiers)

HP

- Histologic features:
  - Bronchiolocentric interstitial lymphoplasmacytic infiltrates
  - Poorly formed interstitial granulomas
  - Organizing pneumonia
  - Chronic bronchiolar inflammation with peribronchiolar metaplasia (Lambertosis)
  - Chronic UIP or NSIP patterns of fibrosis
    - These can act aggressively
- May be subtle histologically
  - Does not correlate with disease severity
- Treatment
  - Remove offending agent
  - Get rid of your bird

Smoking-Related Lung Diseases

- RB-ILD : Respiratory Bronchiolitis (RB)- ILD
- DIP : DIP – Desquamative IP –misnomer – responds to smoking cessation and steroids
- SRIF
- LCH


RB – very common incidental finding in smokers

Image Courtesy : Dr. Churg's presentation


DIP- Misnomer

Image Courtesy : Dr. Churg's presentation


DIP

Image Courtesy : Dr. Churg's presentation

Desquamative Interstitial Pneumonia

- Uncommon smoking related lung disorder
- Primarily an alveolar filling process with secondary interstitial fibrosis
- Associated with other smoking related lung diseases: respiratory bronchitis/bronchiolitis, emphysema, Langerhans cell histiocytosis
- Dense aggregate of pigmented macrophages (smokers macrophages) and temporally uniform alveolar wall fibrosis
- Treat with steroids and cessation of smoking







Poll-time: This lesion is likely to be positive for

A. CD1a, S-100
B. Keratin, Calretinin
C. Keratin, TTF-1
D. EMA, PR



© 2005 Elsevier Inc. Leslie & Wick: Practical Pulmonary Pathology

Langerhans cells, coffee bean shaped, grooves

Pulmonary Langerhans cell histiocytosis

- A.K.A. eosinophilic granuloma, histiocytosis X
- Reactive smoking-related disease
  - Often seen with DIP
- Stellate centrilobular scar with eosinophils and Langerhans cells
  - CD1a and S-100+
- EM - Birbeck granules (tennis racket)



https://commons.wikimedia.org/

Giant Cell Interstitial Pneumonia

- Rare ILD
- Multinucleated histiocytic giant cells in the alveolar spaces
- Emperiopolesis/cannabalism
- Hard-metal disease (Cobalt, tungsten)
- Clinically different from giant cell pneumonia associated with measles and no nuclear/cytoplasmic inclusions

Art by
Dr. EM
Wojcik



https://emedicine.medscape.com/article/2078412-overview#a5



Eosinophils and macrophages in alveolar space

Eosinophilic pneumonia

- May be simple, acute or chronic
- Peripheral infiltrates on chest x-ray
- Loeffler's syndrome (EosPNA+ allergy)
- Eosinophilic granulomatosis with polyangiitis : Churg Strauss/ (EosPNA + asthma+ eosinophilia)
- Acute idiopathic EP causes fevers and acute lung injury
  - PB eosinophils may be normal or high
  - Looks like DAD with eosinophils
- Must rule out secondary causes:
  - Drugs
  - Parasites (Tropical EosPNA due to Wucheria bancrofti)
- Histology demonstrates numerous eosinophils, macrophages
  - Acute EosPNA can have hyaline membranes
  - Chronic EosPNA can have interstitial fibrosis
- Patients often respond to steroids





Irregular granuloma with basophilic necrosis



Courtesy: Dr. S. Mehrotra



American College of Rheumatology: Image Library
Image 99-03-0022
http://images.rheumatology.org/bp/#/search?q=anca&filters=%257B%257D#2862324

https://twitter.com/edgarvlermamd





Wegener granulomatosis
Granulomatosis with Polyangiitis (GPA)

- Vasculitis, granulomas and geographic necrosis involving the upper and lower respiratory tract and kidneys
- Usually C-ANCA+
- Present with bloody nasal discharge, ulcers, polyarthralgias, sinus pain
- Untreated is fatal
  - 85-90% respond to steroids







Goodpasture's syndrome

- Usually young men, present with hemoptysis
- Anti-alveolar basement membrane antibodies
- Immunofluorescence
  - linear staining along alveolar septal basement membranes with IgG and complement
- Nonspecific histology
  - Intra-alveolar hemorrhage
  - ± acute capillaritis

Art by
Dr. EM
Wojcik



Vascular Lung Disease

Pulmonary thromboembolism
with infarction



Thromboembolism

- Saddle embolus
  - involves both main branches of the pulmonary artery
  - causes sudden death.
- Smaller emboli cause wedge shaped infarction with hemorrhage.
- Paradoxical embolus travels across a defect in the heart (ASD or VSD) resulting in thromboemboli in the systemic circulation – danger of stroke.







Pulmonary hypertension

- Primary or secondary (vast majority are secondary)
- Secondary causes include:
  - Obstructive or interstitial lung disease
  - Heart disease
  - Recurrent thromboemboli
- PPH seen in young to middle age women, due to dysfunction of endothelial cell
- Elevated pulmonary artery pressures
  - Echocardiogram
  - Cardiac catheratization
- Features in PPH and not secondary PH
  - Plexiform lesion
  - Angiomatoid lesion
  - Necrotizing arteritis



Art by
Dr. EM
Wojcik





Lymphangioleiomyomatosis

- Cystic lung disease with numerous cysts lined by plump smooth muscle cells
  - Young to middle aged women
- Present with hemoptysis, chylothorax and pneumothorax
- Associated with tuberous sclerosis
  - Mutation on TSC2 gene
- PEComa family of tumors
  - Melanocytic and muscle markers, positive
  - estrogen
- Assoc with surgar tumor (clear cell carcinoma)
- Sometimes treated with lung transplant





© 2005 Elsevier Inc. Leslie & Wick: Practical Pulmonary Pathology



Book image from : Non-neoplastic Pulmonary Pathology: Sanjay Mukhopadhyay



Cartoon Depiction of PAP

Pulmonary Alveolar Proteinosis (PAP)

- Rare disorder of unknown etiology with alveolar filling of granular surfactant material
- Patchy involvement of lung
- Most common: Autoimmune PAP:
  - Neutralizing Ab to GM-CSF –macrophages non-functional –no clearance
- Secondary forms
  - Patients with infections or hematologic malignancies, MDS, AML, inhaled dusts (Acute Silicosis)
- Congenital form / Hereditary PAP
  - Defect in surfactant production and metabolism (SFTPB and SFTPC mutations)
  - CSF2RA mutations may cause a rare hereditary form of PAP

PAP

- Radiology: Crazy pavement
- PAS stain
- Elevated serum anti-GM-CSF (primary cases)
- Rule out PCP
- RX:
  - therapeutic BAL
  - GM-CSF therapy
  - Rituxan
  - Transplantation



Art by
Dr. EM
Wojcik



Lymph node replaced by granulomas



Epithelioid histiocytes (arrowheads): Folded, bland, "bathroom slipper/ flip flop/ insole" shaped nuclei



Bronchocentric distribution of granulomas



Kumar et al: Robbins & Cotran Pathologic Basis of Disease, 8th Edition.
Copyright © 2009 by Saunders, an imprint of Elsevier, Inc. All rights reserved.

Interstitial distribution of granulomas





Multinucleated giant cells are characteristically present in the disease, often accompanied by a variety of distinctive (but not specific) cytoplasmic inclusions: (A) upper left, asteroid body; (B) upper right, Schaumann (conchoidal) body; (C) lower left, Schaumann bodies; (D) lower right, Schaumann body in polarized light.

Sarcoidosis

- Multisystem disease of unknown etiology
- Characterized by non-caseating granulomas
- Diagnosis of exclusion
  - For example, must rule out infections (TB, fungi) when see granulomas

Sarcoidosis

- Adults < 40
- African Americans
- Danish and Swedish
- Typically disease of young adults
- Tends to be upper lobe predominant
- 2/3 rd patients are asymptomatic
- CXR abnormalities in 90% patients
  - Bilateral hilar lymphadenopathy
  - Lung parenchymal abnormalities



Images courtesy CXR - Dr. A. Chandrasekhar, https://www.flickr.com/photos/pulmonary_pathology/7609643866

Sarcoidosis
Laboratory/Diagnostic Abnormalitiesc

- Elevated serum angiotensin converting enzyme (ACE) level
- Hypercalcemia
- Markers to systemic immune abnormalities
  - Anergy to common skin test antigens
    - Candida, PPD
  - Polyclonal hypergammaglobulinemia

Sarcoidosis

- Noncaseating granulomas
  - Rarely shows caseating necrosis
  - Beryliosis is diff dx of non caseating lung granulomas
- Granulomas are present around lymphatics – accessible to transbronchial biopsy
- Rule out infection with special stains
- Can lead to transplant

Sarcoidosis and Vitamin D Hypercalcemia


25-OH Vitamin D
1,25 - OH Vitamin D
Calcium

Diagnosis of Sarcoidosis

- Diagnosis:
  - Made on a transbronchial biopsy in 90% if 4 pieces were submitted
  - Routine AFB and GMS
  - Diagnosis of exclusion









Silica crystals
Under polarized
light

Web Path

Silicosis

- Foundries, sandblasting, mines
- Most prevalent chronic world occupational disease
- Inhalation of crystalline silica
  - Within macrophages causes activation, release of mediators
    - Fibrogenic

Silicosis

- Nodules
  - Concentric hyalinized collagen
  - Whorled appearance
- Coalescence of nodules
- Progressive massive fibrosis
- Fibrotic lesions in hilar lymph nodes, pleura

“Eggshell” calcification of lymph nodes



Silicosis

- Incidental finding on CXR
- PMF – dyspnea, pulm HTN, cor pulmonale
- Increase susceptibility to tuberculosis
  - Silica depresses cell-mediated immunity
  - Pulmonary macrophage phagocytosis impaired
- Carcinogenic – lung cancer
  - Still under debate



Art by
Dr. EM
Wojcik







Fat and loose connective tissue



Hamartoma

- Benign neoplasm with various mesenchymal tissues
- Contains entrapped, “clefted” respiratory epithelium



Round cells with eosinophilic cytoplasm and bland nuclei

Sclerosing hemangioma

- Frequently on boards
- Benign, well circumscribed tumor
- AKA pneumocytoma and sclerosing epithelioma
- Tumor of primitive respiratory epithelium, not vascular tumor (not hemangioma)
  - Type 2 pneumocytes (hobnailing) line the surface of papillary structures with underlying round cells
- 2 types of cells
  - Surface epithelial cells (hobnailed cells)
    - Keratin +, EMA+ TTF-1+
  - Round cells (bland cells)
    - Keratin -, TTF-1 +, EMA+



Keratin formation

The Osler Institute www.osler.org




Central mass
Necrotic and cavitating center

http://library.med.utah.edu/


Intercellular bridges
Keratinization


Bright Orange: Keratinized cells

Cytology : PAP stain http://pathology.jhu.edu/

Squamous cell CA

- Closely linked to smoking
- Usually central
  - arise from dysplastic squamous metaplasia
- Usually p63+, CK5/6+, CK7+, CK20- TTF1-
- Basaloid variant has peripheral palisading
- Not associated with EGFR mutations
- Histology
  - Keratin formation
  - Intracellular bridges
- Local nodes involved in 70-90%
- Spreads outside the thorax, later than other histologic types
- 5-7.5% : 5 year survival

Squamous cell carcinoma -Molecular Genetics

- Highest frequency of p53 mutations
- Others include CDKN2A, PTEN, PIK3CA, NOTCH1, RB1
- Very rare EGFR and KRAS mutations

The Osler Institute www.osler.org





Irregular spiculated mass with entrapped anthracosis

Spiculated lung mass on CT



http://library.med.utah.edu/

Precursor lesions



Atypical Adenomatous Hyperplasia (AAH)

Adenocarcinoma in situ (AIS)

Minimally invasive adenocarcinoma

Precursor lesions -contd

AAH

- Well demarcated focus
- Less than 5 mm
- Lined by cuboidal to low columnar epithelium
- Mild atypia

AIS

- Formerly known as Bronchioloalveolar carcinoma (BAC)
- 3 cm or less
- Grow along pre-existing structures
- Preservation of alveolar architecture
- NO INVASION

MIA

- ✓ 3 cm or less AND
- ✓ Invasive focus less than 5 mm
- ✓ No invasion into pleura, lymphatics or blood vessels



Well-demarcated nodule

AAH

Alveoli lined by cuboidal epithelium with mild atypia



Normal alveoli

Cellular proliferation along septae without invasion

Moderate to marked atypia

AIS (aka) BRONCHIOLOALVEOLAR CARCINOMA



Invasive adenocarcinoma

Kumar et al: Robbins & Cotran Pathologic Basis of Disease, 8th Edition.
Copyright © 2009 by Saunders, an imprint of Elsevier, Inc. All rights reserved.





Bronchoalveolar carcinoma, hobnail neoplastic cells spreading out along alveoli

Subtypes of Adenocarcinoma

a. Acinar adenocarcinoma

b. Papillary adenocarcinoma

c. Solid adenocarcinom

d. Mucinous adenocarcinoma



Adenocarcinoma

- Relatively more frequent in women (most common type in women)
- More peripheral, but can be central also
- Less strongly associated with smoking (most common type in non-smokers)
- Progresses through AAH – AIS
- Spiculated mass, central scarring
- Adenocarcinoma: gland forming or mucin containing or TTF-1 positive
- Slower growing, smaller mass but can metastasize early
- Local nodes involved in over 50%
- 10-12% 5 year survival

Molecular targets for lung cancer (drug examples)

- EGFR, (cetuximab, gefitinib, erlotinib), non-smokers, women, Asian
- EML4-ALK (Crizotinib), young non-smokers
- VEGF, (Avastin, etc.)
- HER2, (Herceptin)
- KRAS, smokers
- BRAF

- Test for ALK and EGFR mutations in anyone with a component of adenocarcinoma
  - Don't test for squamous cell carcinoma or small cell carcinoma

Summary and Current Recommendations for Mol. Testing

- Recommendations to test all advanced stage IIIB and IV adenocarcinoma patients regardless of clinical characteristics
- Squamous cell carcinoma – at least PDL1
- Early stage NSCLC is left to institutional policy!!
- Reflex and Bespoke testing
- EGFR, ALK and ROS1 and PDL1 have to be absolutely tested
- Suggested additional testing: BRAF*, NTREK*, MET, RET, HER2 and KRAS
- *: NCCN guidelines

- 54-year old non-smoker Asian woman presents with cough and right sided abdominal pain. Imaging reveals a 2.1 cm lung mass and multiple PET-avid lesions in the liver. Liver core biopsy shows malignant, infiltrative tumor with acinar pattern. Immunostatins demonstrate lesional cells to be positive for TTF-1 and Napsin-A. Which of the following molecular alterations is likely to be detected in this tumor?
- BRAF V600E mutation
- BAP1 mutation
- EGFR L858R mutation
- Homozygous deletion of CDKN2A



















Neuroendocrine Tumors of the Lung

An Update

Natasha Rekhtman, MD, PhD

Table 1. Summary of Diagnostic Criteria and Grading of Lung Neuroendocrine Tumors Based on the 2004 World Health Organization Classification

	Typical Carcinoid	Atypical Carcinoid	Large Cell Neuroendocrine Carcinoma	Small Cell Lung Carcinoma
Grade	Low	Intermediate	High	High
Morphology	Well-differentiated NET	Well-differentiated NET	Poorly differentiated NET	Poorly differentiated NET
Mitoses per 10 HPFs[a]	<2	2–10	>10 (median, 70)	>10 (median, 80)
Necrosis	None	Present (focal punctate)	Present (extensive)	Present (extensive)

Abbreviations: HPFs, high-power fields; NET, neuroendocrine tumor.

[a] Field diameter with a ×40 objective (0.2 mm^2).

NEUROENDOCRINE TUMORS LUNG



Neuroendocrine tumor	Mitosis/2 mm^2 (%)	Necrosis	Grade
Typical carcinoid	<2	No necrosis	1
Atypical carcinoid	2–10	Foci of necrosis	2
LCNEC/SCLC	>10	Necrosis	3

LCNEC: Large cell neuroendocrine cancer; SCLC: Small cell lung cancer.

https://www.researchgate.net/publication/280288714_Neuroendocrine_tumors_of_the_lung_a_comprehensive_overview/figures?lo=1

Typical carcinoid

- Typical carcinoid: organoid pattern, nested, trabecular, rosettes, uniform nuclei, granular (salt and pepper) chromatin.
- Minimal mitotic activity (1-2/10 HPFs)
- NO necrosis



Trabecular pattern (cords of cells)

Atypical carcinoid

- Increased mitotic activity (3-10/10HPFs)
- Or minimal necrosis






[redacted]


Ki67
High proliferative rate




Synaptophysin
Often patchy neuroendocrine markers

Small cell carcinoma

- Sheets of cells with scant cytoplasm
  - can see nuclear molding
  - Usually lack nucleoli
  - Numerous mitoses
  - Punctate and zonal necrosis
- 75% express neuroendocrine markers
  - NSE, chromogranin, synaptophysin
- >50% Ki67 staining



Epithelioid hemangioendothelioma

- Usually present as multiple nodules in young women
- Circumscribed pale nodules on H&E
- Periphery is more cellular, central area is hyalinized or necrotic
- IVBAT

Pleural Disease

- Often results in a pleural effusion, or fluid in the pleural space. The type of fluid is very important:
  - Transudate: a low specific gravity (or thin) fluid without inflammatory cells.
    - Secondary to heart failure.
  - Exudate: a more viscous fluid with inflammatory cells.
    - Secondary to infection or cancer
  - Lymph fluid: milky lymphatic fluid rich in triglycerides.
    - Secondary to lymphatic duct obstruction. Called chylothorax.
  - Blood: called hemothorax.
    - Secondary to trauma or malignancy



Pleurectomy – Decortication: Measure Volume and Weight



Extrapleural Pneumonectomy

Histologic Patterns of Pleural MM

- Three major subtypes
  - Epithelioid – Most common pattern (65-70%)*
  - Sarcomatoid (< 10%)
    - Desmoplastic (< 2 %)
  - Biphasic – 10-15% (at least 10% of epithelioid and sarcomatoid components)
- Most epithelioid tumors are readily recognized or suspected on H&E
  - Bland vesicular chromatin, abundant eosinophilic cytoplasm
  - Infrequent mitoses
- Wide range of histologic patterns within the same tumor and between different cases
  - Important for diagnostic and prognostic considerations

Wide variety of histologic patterns in diffuse malignant mesothelioma

- Epithelioid
  - Tubulopapillary
  - Adenomatoid
  - Solid well-differentiated
  - Acinar
  - Clear cell
  - Deciduoid
  - Adenoid cystic
  - Signet ring
  - Small cell
- Sarcomatous (these have a worse prognosis)
  - Lymphohistiocytoid
  - Desmoplastic (bland, hard to diagnosis)
- Biphasic

Histologic Patterns of Pleural MM



Epithelioid



Biphasic



Sarcomatoid



Immunohistochemical stains for diagnosis of EMM and distinguishing it from lung tumors with pleural involvement



WT1	Calretinin	Keratin 5/6	Podoplanin (D2-40)
Nuclear	Nuclear and cytoplasmic	Cytoplasmic	Membrane
70-95% of EMM	Nearly 100% of EMM	75-100% of EMM	80-100% of EMM
Negative in lung ADC	Positive in 5-10% of lung ADC (focal)	Focal positivity in 2-20% of lung ADC	Focal positivity in less than 15% of lung ADC
Negative in lung SQC	Approximately 40% of lung SQC (focal)	Positive in 100% of lung SQC	50% of lung SQC











Mesothelioma

- Mesothelioma is a rare, but highly lethal, malignancy arising from the mesothelial cells lining the pleural surface
- Asbestos exposure is the number one risk factor
- Bland epithelioid cells invading chest wall
- Positive markers: calretinin (cyto and nuclear), CK 5/6, thrombomodulin, WT-1 (nuclear)
- Sarcomatoid mesothelioma loss of these antigens
  - may only be calretinin +



Asbestos and MM

- Hydrated magnesium silicate
- Asbestos – Greek for "unquenchable"
- Extensively used in construction industry
- Two types of fibers:
  - Curly, serpentine, pliable, chrysotile fibers (most commonly used)
  - Rod or needle-like amphibole fibers – higher order of carcinogenicity
- Complex relationship b/w asbestos and MM
  - long latency period
  - 70-80% of MM in men assoc with asbestos while only 20% of women have this assocn
  - Only 5% of asbestos workers develop MM
- Unusual carcinogen: asbestos is neither metabolized nor incorporated into DNA

The End

Additional reading material



Acute bronchopneumonia, neutrophils filling alveolar spaces

Streptococcus pneumoniae

- Gram-positive, catalase-negative cocci
- Grows in short chains or pairs
- Causes 50% of bacterial community acquired pneumonia (CAP)
- Capsule is virulence factor
- Winter-spring; often follows influenza
- 60% of general population colonized

Pseudomonas aeruginosa

- Gram-negative, motile, strictly aerobic, oxidase positive, respiratory metabolism
- Resistant to many disinfectants (common nosocomial pathogen)
- Common hospital acquired pathogen
- Complication of COPD with steroids, cystic fibrosis, contaminated respiratory equipment

Legionella pneumophila

- Gram-negative bacilli, non-spore forming
- Live in aquatic environments, infects through aerosolized droplets (A/Cs, humidifiers, respiratory therapy equipment)
- Not person-to-person
- 8% of CAP
- Pontiac fever is a milder clinical syndrome caused by L. pneumophila and other Legionella species
- Histology, acute bronchopneumonia, neutrophils in the airways

Sarcoidosis – Systemic Involvement

- Heart
  - Restrictive cardiomyopathy
  - Conduction abnormalities
- Bell's Palsy
  - CN VIII
- Eye (20-50% pts)
  - Uveitis
- Skin (30-50% pts)
  - Erythema nodosum : acute, nodular erythematous eruptions on the extensor aspect of LL.
  - Lupus Pernio: Chronic indurated plaques on the mid-face

Erythema Nodosum

Lupus Pernio

http://www.aafp.org






Obliterated bronchus, still has smooth muscle but no epithelium

Constrictive bronchiolitis

- Histologic changes seen with the clinical syndrome of bronchiolitis obliterans
- Associated with many conditions
  - Pulmonary Graft vs Host disease
  - Healed infection
  - Fume or toxin exposure
  - Connective tissue disease, especially RA
  - Lung transplant rejection
  - Drug reaction
  - Inflammatory bowel disease-associated
  - DIPNECH
  - Asthma
  - Hypersensitivity pneumonitis
  - Idiopathic

Histologic features

- THE MOST DISTINCTIVE FEATURE MAY BE THE SUBTLETY OF CHANGES.
  - Patient symptoms worse than histology
- submucosal scarring, concentric luminal narrowing,
- adventitial scarring, chronic inflammation,
- bronchiolar ectasia
- mucus stasis, smooth muscle hypertrophy



Rejection

- Acute rejection
  - Changes in alveoli
  - Intraparenchymal perivascular mononuclear infiltrates
  - More PMNs, eos and macrophages as disease develops
  - Endothelialitis
- Lymphocytic bronchitis/bronchiolitis
  - Combined large and small airway inflammation
- Chronic airway rejection
  - Bronchiolitis obliterans
    - Prevalence 40%, mortality 25%
    - i.e fibrosis, scarred lumen of bronchiole
- Chronic vascular rejection
  - Fibrointimal thickening of arteries and veins





Solitary Fibrous tumor

- Uncommon spindle cell pleural based tumor
- Hemangiopericytoma-vascular pattern
  - Staghorn blood vessels
- “patternless-pattern”
- Ropey collagen
- Phenotype: CD34, Bcl-2, CD99, STAT-6
- Malignant features are: greater cellularity, infiltrative growth pattern, celllular atypia, > 4 mits/ 10 HPFs

This page was intentionally left blank.

Head & Neck Pathology

Pathology Board Review
Osler Institute
April 2021

Nicole A. Cipriani, MD
Nicole.Cipriani@uchospitals.edu

AT THE FOREFRONT
UChicago Medicine

Outline

- Squamous Lesions & Carcinomas
- Mandible & Maxilla
- Nose & Paranasal Sinuses & Nasopharynx
- Larynx
- Salivary Glands
- Ear & Temporal Bone

Squamous Lesions

- Non-neoplastic / Benign
  - Leukoplakia
  - Candidiasis
  - Lichen Planus
  - Vesiculobullous Diseaes
  - Epulis / Granular Cell Tumor
  - Pyogenic Granuloma
- Dysplasia
- Keratinizing Squamous Cell Carcinoma
- Nonkeratinizing Squamous Cell Carcinoma
  - HPV-related Oropharynx
  - EBV-related Nasopharynx (Nasopharyngeal CA)

Leukoplakia / Erythroplakia

- Are clinical appearances of oral mucosa
  - White / Red / Both
  - Histologically, may represent hyperkeratosis, hyperplasia, dysplasia, etc





Oral Hairy Leukoplakia

- Occurs mostly in HIV patients
- Associated with Epstein-Barr virus (EBV)
- Lateral border of tongue
- Cannot be "wiped off" tongue
  - Unlike thrush
- Histology:
  - Hyperkeratosis, Hyperplasia
  - Balloon cells in spinous layer (below granular layer)
    - Cytoplasmic clearing / ballooning
    - Chromatin margination
  - Little inflammation
  - Decreased Langerhans cells (may promote EBV persistence)



Oral Candidiasis

- Most common oral fungal infection
- Usually C. albicans
- Pseudomembranous (thrush): white plaques that "wipe off" with scraping
- Erythematous: red atrophic mucosa with burning
- Hyperplastic: white plaques that cannot be wiped off
  - Can be difficult to separate from oral leukoplakia secondarily superimposed with Candida
  - Thick epithelium, broad rete / Superficial neutrophils / Yeast piercing through surface





Lichen Planus

- Autoimmune mucositis
- F>M
- Young adult – middle age at presentation
- Usually asymptomatic
- Clinically: Wickham's striae
  - Reticular pattern of raised, white (hyperkeratotic) streaks on red background
- Histologically:
  - Sawtooth rete ridges
  - Band-like subepithelial chronic inflammatory infiltrate
- IF: granular fibrinogen at basement membrane


http://www.ijdr.in/articles/2011/22/6/images/IndianJDentRes_2011_22_6_827_94679_f5.jpg

Vesiculobullous Diseases

Pemphigus Vulgaris

- Supra-basal / Intra-epithelial blister
- IF: intercellular "fishnet" IgG, IgM, C3
- Anti-desmoglein1/3

Bullous Pemphigoid

- Sub-basal, Sub-epithelial blister
- IF: basement membrane (subepithelial) linear IgG, C3
- Anti-BP-antigen (BPAG1/2)


Vulgar Fishnets



Oral Hairy Leukoplakia	Oral Candidiasis	Lichen Planus	Pemphigus Vulgaris	Bullous Pemphigoid
EBV Immunocomp.	C. albicans	Autoimmune	Autoimmune Anti-desmoglein INTERcellular	Autoimmune Anti-BP-Antigen Basemt. memb
Lateral border tongue	White plaques	Wickham's striae white on red	Blister	Blister
Balloon cells	Hyperplasia Neutrophils Yeast forms	Sawtooth rete Band-like stromal CI	INTRAepithelial blister	SUBepithelial blister
		Granular fibrinogen at BM	IgG, C3 vulgar fishnets	IgG, C3 at BM



Subepithelial Squamous Lesions

Epulis		Granular Cell Tumor	
Newborns Alveolar ridge Exophytic mass		Adults Dorsal tongue Deep / sessile mass	 Schwann cell?
Granular cells Distinct cell borders S-100 negative		Granular cells Indistinct cell borders S-100/SOX-10 positive	
No pseudoepitheliomatous hyperplasia		Pseudoepitheliomatous hyperplasia present	



Lobular Capillary Hemangioma
(Pyogenic Granuloma)

- Benign tumor / Reactive vascular proliferation
- Lobules
  - Central larger vessel
  - Peripheral smaller capillaries




Lobular Capillary Hemangioma
(Pyogenic Granuloma)

- Often on lip
- Slit-like capillary spaces
- Lined by flat endothelial cells
- Mitoses are ok




Squamous Dysplasia

- The usual suspects: Tobacco, Alcohol, Betel nuts.
- Must eliminate superficial hyper/parakeratotic layer when grading dysplasia.
- Grade of dysplasia does not always correlate to presence of invasive carcinoma.
  - Can have no / low grade dysplasia and adjacent carcinoma.
- Prognosis in oral cavity relates to tumor size and nodal status.



Keratinizing dysplasia of UADT usually leads to Keratinizing SCC:

Dysplasia	Risk of SCC	
Hyperplasia	0-3%	
Mild	Up to 30%	Some low grade lesions progress
Moderate	Up to 45%	
Severe	Up to 60%	Not all high grade lesions progress
CIS	Up to 80%	

DOI - Depth of Invasion

- DOI (important for stage) is measured from adjacent normal basement membrane to deepest tumor island
- This is NOT tumor thickness


- - - - Normal basal line
.... Maximum depth of invasion
Exophytic
Ulcerated



Nonkeratinizing SCC

Oropharynx	Nasopharynx
HPV (High Risk 16, 18, 31, 33)	EBV, nitrosamines > HPV
Middle aged M Cystic neck nodes	Middle aged M Cystic neck nodes, Epistaxis Hong Kong, China, Southeast Asia, North Africa, Arctic
No surface dysplasia Blue on blue	No surface dysplasia Often syncytial
p16+ / HPV E6-E6 mRNA ISH / DNA PCR GOOD PROGNOSIS	EBER RNA ISH Somewhat more radiosensitive if EBV+



Squamous Cell Carcinoma Non-Keratinizing

- Oropharynx (base of tongue, tonsil)
  - Related to HPV-infection (High Risk 16, 18, 31, 33)
  - Men, age 50s-60s, cystic lymph node metastases
  - Good prognosis
  - p16 positive



HPV Non-keratinizing SCC



The Oropharynx is a "Special Site"

- HPV-related SCC of the oropharynx is thought to originate in the deep tonsillar crypts, where the basement membrane is discontinuous and lymphatics exist in close association with the epithelium
- ALL such HPV-related carcinomas should be considered invasive, discarding the concept of in-situ carcinoma and "crypt dysplasia"
  - Usually the invasion appears broad-based without single cell invasion

p16

- p16 is a surrogate marker for HPV-induced oncogenesis
  - Cell cycle protein upregulated by HPV viral proteins (E6/E7)
  - Works well in the OROPHARYNX (BOT, TONSIL)
  - It should show diffuse, strong, nuclear and cytoplasmic staining in most cells (some say ~70%)
  - Low PPV for HPV-involvement in oral cavity & other sites where p16 can be upregulated for reasons OTHER than virus
- p16 may be combined with a confirmatory test
  - HPV E6-E7 RNA ISH or HPV DNA PCR



p16-ICH



HPV-ISH



HPV-PCR

HPV & Neck Cysts

- Malignant appearing squamous epithelium in a neck lymph node
  - Is NOT carcinoma arising in a branchial cleft cyst
  - Is highly likely to be metastatic HPV-related SCC
    - p16+ suggests primary Oropharynx



Benign epithelium





Malignant epithelium

Images courtesy Jamie Steinmetz, MD

Human Papillomavirus Testing in Head and Neck Carcinomas

Guideline From the College of American Pathologists

James S. Lewis Jr, MD; Beth Beadle, MD, PhD; Justin A. Bishop, MD; Rebecca D. Chernock, MD; Carol Colasacco, MLIS, SCT(ASCP); Christina Lacchetti, MHSc; Joel Todd Moncur, MD, PhD; James W. Rocco, MD, PhD; Mary R. Schwartz, MD; Raja R. Seethala, MD; Nicole E. Thomas, MPH, CT(ASCP)^{CM}; William H. Westra, MD; William C. Faquin, MD, PhD

- Bottom line:
  - Perform HR-HPV testing by p16 on all newly diagnosed OPSCC.
  - Do NOT perform HR-HPV testing on nonOP primary SCC.
  - Do NOT provide a tumor grade or differentiation status for HPV-positive/p16-positive OPSCCs.
  - Perform HR-HPV testing on metastatic CUPs in Neck Levels II-III

Lewis JS Jr, Beadle B, Bishop JA, et al. Human Papillomavirus Testing in Head and Neck Carcinomas: Guideline From the College of American Pathologists. Arch Pathol Lab Med. 2018;142(5):559-597. doi:10.5858/arpa.2017-0286-CP

Table 2. Summary of Guideline Statements

Guideline Statement	Strength of Recommendation
1. Pathologists should perform HR-HPV testing on all patients with newly diagnosed OPSCC, including all histologic subtypes. This testing may be performed on the primary tumor or on a regional lymph node metastasis when the clinical findings are consistent with an oropharyngeal primary.	Strong recommendation
2. For oropharyngeal tissue specimens (ie, noncytology), pathologists should perform HR-HPV testing by surrogate marker p16 IHC. Additional HPV-specific testing may be done at the discretion of the pathologist and/or treating clinician, or in the context of a clinical trial.	Recommendation
3. Pathologists should not routinely perform HR-HPV testing on patients with nonsquamous carcinomas of the oropharynx.	Expert consensus opinion
4. Pathologists should not routinely perform HR-HPV testing on patients with nonoropharyngeal primary tumors of the head and neck.	Recommendation
5. Pathologists should routinely perform HR-HPV testing on patients with metastatic SCC of unknown primary in a cervical upper or mid jugular chain lymph node. An explanatory note on the significance of a positive HPV result is recommended.	Recommendation
6. For tissue specimens (ie, noncytology) from patients presenting with metastatic SCC of unknown primary in a cervical upper- or mid-jugular chain lymph node, pathologists should perform p16 IHC. *Note:* Additional HR-HPV testing on p16-positive cases should be performed for tumors located outside of level II or III (nonroutine testing) in the neck and/or for tumors with keratinizing morphology.	Expert consensus opinion
7. Pathologists should perform HR-HPV testing on head and neck FNA SCC samples from all patients with known OPSCC not previously tested for HR-HPV, with suspected OPSCC, or with metastatic SCC of unknown primary. *Note:* No recommendation is made for or against any specific testing methodology for HR-HPV testing in FNA samples. If the result of HR-HPV testing on the FNA sample is negative, testing should be performed on tissue if it becomes available. If pathologists use cytology samples for p16 IHC testing, they should validate the criteria (ie, cutoff) for a positive result.	Expert consensus opinion
8. Pathologists should report p16 IHC positivity as a surrogate for HR-HPV in tissue specimens (ie, noncytology) when there is at least 70% nuclear and cytoplasmic expression with at least moderate to strong intensity.	Expert consensus opinion
9. Pathologists should not routinely perform low-risk HPV testing on patients with head and neck carcinomas.	Expert consensus opinion
10. Pathologists should *not* repeat HPV testing on patients with locally recurrent, regionally recurrent, or persistent tumor if primary tumor HR-HPV status has already been established. If initial HR-HPV status was never assessed or results are unknown, testing is recommended. HPV testing may be performed on a case-by-case basis for diagnostic purposes if there is uncertainty regarding whether the tumor in question is a recurrence or a new primary SCC.	Expert consensus opinion
11. Pathologists should *not* routinely perform HR-HPV testing on patients with distant metastases if primary tumor HR-HPV status has been established. HPV testing may be performed on a case-by-case basis for diagnostic purposes if there is uncertainty regarding whether the tumor in question is a metastasis or a new primary SCC.	Expert consensus opinion
12. Pathologists should report primary OPSCCs that test positive for HR-HPV or its surrogate marker p16 as HPV positive and/or p16 positive.	Expert consensus opinion
13. Pathologists should *not* provide a tumor grade or differentiation status for HPV-positive/p16-positive OPSCCs.	Expert consensus opinion
14. Pathologists should *not* alter HR-HPV testing strategy based on patient smoking history.	Expert consensus opinion

Abbreviations: FNA, fine-needle aspiration; HPV, human papillomavirus; HR-HPV, high-risk HPV; IHC, immunohistochemistry; OPSCC, oropharyngeal SCC; SCC, squamous cell carcinoma.

JOURNAL OF CLINICAL ONCOLOGY ASCO SPECIAL ARTICLE

Human Papillomavirus Testing in Head and Neck Carcinomas: ASCO Clinical Practice Guideline Endorsement of the College of American Pathologists Guideline

Carole Fakhry, Christina Lacchetti, Lisa M. Rooper, Richard C. Jordan, Danny Rischin, Erich M. Sturgis, Diana Bell, Mark W. Lingen, Seema Harichand-Herdt, John Thibo, Jose Zevallos, and Bayardo Perez-Ordonez

- Similar Bottom line:
  - Perform HR-HPV testing by p16 on all newly diagnosed OPSCC.
  - Do NOT perform HR-HPV testing on nonOP primary SCC.

Recommendations
It is recommended that HPV tumor status should be determined for newly diagnosed oropharyngeal squamous cell carcinomas. HPV tumor status testing may be performed by surrogate marker p16 immunohistochemistry either on the primary tumor or from cervical nodal metastases only if an oropharyngeal primary tumor is present. The threshold for positivity is at least 70% nuclear and cytoplasmic expression with at least moderate to strong intensity. Additional confirmatory testing may be done at the discretion of the pathologist and/or treating clinician. Pathologists should not routinely determine HPV tumor status in nonsquamous carcinomas of the oropharynx or non–oropharyngeal squamous cell carcinomas of the head and neck. When there is uncertainty of histologic type or whether a poorly differentiated oropharyngeal tumor is nonsquamous, HPV tumor status testing may be warranted and at the discretion of the pathologist and/or treating clinician. Additional information is available at: www.asco.org/head-neck-cancer-guidelines.

J Clin Oncol 36. © 2018 by American Society of Clinical Oncology

Fakhry C, Lacchetti C, Rooper LM, et al. Human Papillomavirus Testing in Head and Neck Carcinomas: ASCO Clinical Practice Guideline Endorsement of the College of American Pathologists Guideline [published correction appears in J Clin Oncol. 2019 Jan 20;37(3):261]. J Clin Oncol. 2018;36(31):3152-3161. doi:10.1200/JCO.18.00684

Nasopharyngeal (Lymphoepithelial) Carcinoma

- High prevalence in:
  - Hong Kong, China, Southeast Asia, North Africa, Arctic
  - EBV exposure, nitrosamine ingestion
- M > F, Age > 30
- Presentation: Epistaxis, Otitis media, Neck metastases
- Can be thought of as a variant of squamous cell carcinoma
  - p63/p40+, HMWK+
  - Keratinizing: Not associated with EBV
  - NON-keratinizing w/ lymphoid stroma: USUALLY EBV+
- Somewhat more radiosensitive if EBV+

Nasopharyngeal Carcinoma Non-Keratinizing

- Related to EBV-infection
- Syncytial growth pattern
- Inflammatory infiltrate
- EBER in-situ hybridization = detects EBV RNA





Why do we care about virus?

- HPV-associated oropharyngeal SCCs have much better prognosis
  - Clinical trials
  - De-intensified treatment



N Engl J Med 2010;363:24-35.

Why do we care about virus?

- EBV-associated nasopharyngeal SCC have somewhat better prognosis
  - Serum EBV titers are monitored in these patients



International Journal of Radiation Oncology, Biology, Physics. 2014. 88(3): 580–588.

Mandible & Maxilla

- Medication (Bisphosphonate) Related Osteonecrosis of Jaw and Osteoradionecrosis
- Odontogenic Lesions
  - DC vs OKC
  - Ameloblastoma
- Mesenchymal Neoplasms
  - Osteoma
  - Osteosarcoma
  - Fibro-osseous lesions
- Misc
  - Giant Cell Lesion, Central vs Peripheral
  - Cherubism


Oral flora colonization: Actinomyces, Cocci


Acute osteomyelitis

MRONJ vs ORN

Medication (Bisphosphonate) Related Osteonecrosis of Jaw vs Osteoradionecrosis



- Tenderness and exposed bone in posterior mandible.
- Triad of: 1) Osteonecrosis, 2) Acute inflammation, 3) Oral flora colonization.
- Secondary to bisphosphonate use (for tumor, osteoporosis) or Radiation

Alendronate (Fosamax®), Risedronate (Actonel®), Ibandronate (Boniva®), Pamidronate (Aredia®), Zolendronate (Zometa®) (Reclast®), Denosumab (Xgeva®) (Prolia®)

Odontogenic Cysts

- Around the crown of an un-erupted tooth, DDx includes:
  - Dentigerous Cyst
  - Odontogenic Keratocyst (OKC)
  - Cystic Ameloblastoma


http://images.radiopaedia.org/images/529991/8abce424a14c555d776b2809c41d70.jpg





Diagnosis?
Syndrome?


http://images.radiopaedia.org/images/529991/8abce424a14c555d776b2809c41d70.jpg



Odontogenic Keratocyst
Gorlin – AKA – Nevoid BCC Syndrome

- AD Chr9
  - Odontogenic Keratocysts / "Gorlin cysts"
  - Multiple BCCs (even on non-sun exposed skin)
  - Medulloblastomas
  - Palmar / plantar pits
  - Skeletal anomalies
- Features of OKC squamous epithelium
  - 5-7 cells thin
  - Palisaded basal layer (fence)
  - Corrugated surface (cardboard)
  - May have daughter cysts in wall – leads to recurrence
- Treatment is marsupialization / caustic treatment of cavity

Diagnosis?

- Dentigerous cyst
  - Proliferative squamous lining
  - Inflammation
  - OKC can look like this but with only focal diagnostic lining – look carefully!



http://images.radiopaedia.org/images/529991/8abce424a14c555d776b2809c41d70.jpg



Diagnosis?

- Ameloblastoma, unicystic







http://images.radiopaedia.org/images/529991/8abce424a14c555d776b2809c41d70.jpg

Ameloblastoma

- Can be cystic or solid
- Solid: Large epithelial nests
  - Peripheral reverse polarity (nuclei away from basement membrane)
  - Central stellate reticulum
- Locally destructive, requires hemi-mandibulectomy
- Many are BRAF V600E mutated





Recapitulates Tooth Development


Odontoblasts (dentin)
Stellate reticulum
Ameloblasts (enamel)

Ameloblastoma


ENUCLEATION
Unicystic, Luminal
Unicystic, Intraluminal
Unicystic, Mural
HEMI-MANDIBULECTOMY
Multicystic
Solid

Histologic variants (Plexiform, Follicular, Acanthomatous, etc) are NOT as important as growth pattern

OKC Odontogenic Keratocyst	Dentigerous Cyst	Ameloblastoma
• Thin 5-7 cells • Palisaded basal layer (picket fence) • Corrugated surface (like cardboard)	• Proliferative squamous lining • Inflammation • OKC can look like this but with only focal diagnostic lining!	• Can be cystic or solid • Solid: Large epithelial nests • Peripheral reverse polarity (nuclei away from basement membrane) • Central stellate reticulum
Marsupialization, Caustic Rx	Curettage is curative	Locally destructive, requires hemi-mandibulectomy
Daughter cysts in wall → recurrence		
Gorlin Syndrome		



http://images.radiopaedia.org/images/529991/8abce424a14c555d776b2809c41d70.jpg

Ameloblastic Fibroma

- Children > Adults
- Swelling of posterior mandible
- +/- unerupted tooth, radiolucent
- Hypocellular bland myxoid stroma
- Compressed nests and anastomosing chords of ameloblastic epithelium
- May recur
- Risk for development of ameloblastic fibrosarcoma



Diagnosis? Syndrome?

http://www.hindawi.com/journals/ijd/2010/834761.fig.003.jpg



Osteoma
Gardner – AKA – variant of FAP

- APC gene Chr 5q21
  - Colonic polyposis / carcinoma
  - Osteomas of the skull
  - Desmoid tumors (fibromatosis)
  - Thyroid cancer
  - Epidermoid cysts
  - Sebaceous cysts
- Osteomas in general
  - Are comprised of compact cortical bone with little stroma
  - Most frequent in frontal / ethmoid sinus
  - 70-90% of patients with MULTIPLE osteomas have Gardner syndrome

Fibro-Osseous Lesions

(Cemento-) Ossifying Fibroma	Fibrous Dysplasia
F>M, 20s-30s CDC73 (HRPT2) mutation in HPT-JT syndrome Juvenile trabecular or psammomatoid variants	GNAS mutation Monostotic > Polyostotic Craniofacial bones + femur
Sharply demarcated Expansile	Poorly demarcated Ground glass
• Variably cellular stroma • Variably sized/shaped woven bone / cementum • Osteoblast rimming • Can be peripheral (in gingiva) or central (in bone)	• Monotonous stroma • Monotonous curvilinear strands of woven bone • No osteoblast rimming • Sharpey's fibers



Gnathic Osteosarcoma

- M>F, Mean age 30s (vs long bone OS which occurs in teens / elderly)
- Mandible > Maxilla
- More frequently chondroblastic
  - Osteoblastic: Makes osteoid
  - Chondroblastic: Also makes cartilage
  - Fibroblastic: Also makes spindle cells
- Mortality due to local recurrence
- Chemotherapy is less effective than in long bones; primary resection is key



http://www.webpathology.com/slides-13/slides/Bone_Osteosarcoma7_Chondroblastic_BaseofSkull.jpg



Mardinger O et al. Oral Surgery, Oral Medicine, Oral Pathology, Oral Radiology, and Endodontics, 91(4), 445–451.



Giant Cell Lesions

- Giant Cell Lesion / Granuloma / Reparative Granuloma
  - Central: in bone (anterior mandible)
  - Peripheral: in gingiva
- Histology:
  - Lobulated growth of mononuclear spindle-to-polygonal cells + osteoclast like giant cells
  - Richly vascular, hemosiderin
  - DDx: Brown tumor, ABC
- Cherubism (AD, SH3BP2 gene)
  - Symmetric jaw enlargement due to giant-cell rich lesion

Nose & Paranasal Sinuses & Nasopharynx

- Sinusitis
  - Fungal: Mycetoma, Allergic Sinusitis, Invasive Sinusitis, Rhinosporidiosis (vs. Rhinoscleroma)
- Wegener's Granulomatosis
- Tumors & Tumor-like lesions
  - Inflammatory Polyp
  - Respiratory Epithelial Adenomatoid Hamartoma
  - Schneiderian Papilloma
  - Nasopharyngeal Angiofibroma
  - Sinonasal Adenocarcinomas
  - Sinonasal Sarcomas
  - Nasal Glial Heterotopia
  - High Grade Tumor Differential (SNUC, etc)



Mycetoma (Fungus Ball)

- Ball of fungus in sinus
  - Usually Aspergilloma
- NO tissue invasion

Aspergillus
Septate hyphae
Acute angle branching



Allergic Fungal Sinusitis

- * Most common fungal sinusitis
- Hypersensitivity to inhaled fungus in atopic people
  - Usually Dematiaceous fungus
- Respiratory mucosa with prominent eosinophilis
- NO fungus in tissue



Allergic Fungal Sinusitis

- Extracellular mucin is full of
  - Eosinophilis
  - Charcot-Leyden crystals
  - Fungus (GMS stain)



Inflammatory Polyp

- NON-neoplastic
- Chronic sinusitis
  - Cystic fibrosis in kids
- Lateral nasal wall masses
- Thick basement membrane
- Myxoid stroma with Eos
  - Atypical stromal fibroblasts
- Cystically dilated glands



Respiratory Epithelial Adenomatoid Hamartoma (REAH)

- Benign NON-neoplastic hamartoma
  - May also be related to chronic sinusitis
- Adult Males
- Olfactory clefts
- Proliferation of respiratory glands w/ goblet cells
  - Surface invaginations
  - Stromal hyalinization / thick basement membrane
  - +/- atrophic acinar proliferation



Invasive Fungal Sinusitis

- Immunocompromised (HIV, hematologic malignancy, DM)
- Fungal hyphae invading tissue with associated
  - Angioinvasion (into vessel wall / lumen with thrombosis)
  - Tissue necrosis
- Mucor in diabetics; also Aspergillus, Candida, etc

Hard to see on frozen?
Consider Diff Quick on air-dried slide



Similar Names

Rhinoscleroma
- Klebsiella rhinoscleromatis (bacteria)
- Bacteria in foamy macros
- Warthin-Starry, Giemsa

Rhinosporidiosis
- Rhinosporidium seeberi (fungus)
- Sporangia in tissue
- DDx Oncocytic Schneid. Pap

Microcysts in Epithelium AND Stroma

http://www.brown.edu/Courses/Digital_Path/systemic_path/hn/rhinoscleroma2.jpg
http://www.dermpedia.org/files/imagecache/book_node/images/Image3_12.jpg



Myospherulosis

- Due to packing of sinus with petrolatum-coated gauze
- RBCs interact with lipid-rich ointment and cause structures that mimic fungus
- GMS stains negative

https://app.expertpath.com/document/myospherulosis/95d2750f-6e19-4e64-884f-2eee48299453?searchTerm=myospherulosis

RBCs encased in membrane



Rhinoscleroma	Rhinosporidiosis	Myospherulosis
Klebsiella rhinoscleromatis (bacteria)	Rhinosporidium seeberi (fungus)	RBCs NOT an organism
Bacteria in foamy macros Warthin-Starry, Giemsa	Sporangia in tissue	RBCs surrounded by petroleum products from sinus-packing gauze

Granulomatosis With Polyangiitis (Wegener's Granulomatosis)

- Systemic necrotizing vasculitis
- Respiratory tract +/- glomerulonephritis
- c-ANCA+ in 50-90%
- Treatment with immunosuppression

- DDx:
  - Sarcoid
  - Fungal sinusitis
  - Cocaine related necrosis
  - Neoplasms



Wegener's Granulomatosis

Granulomatosis With Polyangiitis

c-ANCA associated destructive vasculitis

Granulomas
(collections of epithelioid histiocytes)

Vasculitis
(inflammatory cells in wall of vessel)

* Neutrophilic microabscesses
(small collections of neutrophils)



Sinonasal Tumors

- Most common site is:
  - Maxillary sinus 60%
- Followed by:
  - Nasal cavity 20-30%
  - Ethmoid sinus 10-15%
  - Sphenoid sinus 1%

http://www.formulamedical.com/Assets/Pictures/Head&Neck/sinus1.jpg



Sinonasal (Schneiderian) Papilloma

Classic Inverted Type	Oncocytic Type (“cylindrical cell”)	Exophytic Type (“fungiform”)
M > F, age 40-70	M = F, age > 50	M > F, age < 50
Lateral Nasal Wall	Lateral Nasal Wall	Nasal Septum
Predominantly endophytic growth Neutrophilic microabscesses in epithelium	Endophytic and exophytic growth Neutrophilic microabscesses in epithelium	Predominantly exophytic growth
Carcinoma in up to 20%	Carcinoma in up to 20%	Carcinoma rare
EGFR mut ~90% HPV-LR>HR	KRAS mut ~100%	HPV-LR

https://c1.staticflickr.com/9/8123/8613998440_5eca0bb9d7_n.jpg



Sinonasal Papilloma, Inverted



Sinonasal Papilloma, Oncocytic

Microcysts in Epithelium only



Sinonasal Adenocarcinoma

Non-intestinal Type (Non-ITAC)	Intestinal Type (ITAC)
Slight M > F, low or high grade	M > F, high grade & locally aggressive
No predisposing condition	Occupational: wood and leather dust exposure
Low grade: ethmoid sinus High grade: maxillary sinus	Most common type is colonic (int grade) Most common location is ethmoid sinus
CK20-, CDX2-, CK7+	CK20+, CDX2+, CK7 var
Small back to back glands, Round to oval nuclei	Large dilated glands, Dark elongated nuclei



NON-ITAC



ITAC

Sinonasal Spindle Cell Tumors



Nasopharyngeal Angiofibroma

- Adolescent Males
  - FAP patients
- Slit-like blood vessels
  - Clinically bloody
- Surrounded by dense eosinophilic fibrous stroma with bland spindle cells
- Stromal cells:
  - AR+
  - Beta-catenin + (mutation)
- Surgery +/- Embolization

http://www.webpathology.com/image.asp?case=152&n=1



Sinonasal Hemangiopericytoma vs Solitary Fibrous Tumor

- Histologically similar
  - Sheets of spindle cells with thin-walled dilated blood vessels
  - Both occur in adults; treated surgically
- In most sites (brain, soft tissue), share identical genetics (NAB2-STAT6 fusion)
- In sinonasal cavity, can be separated into 2 distinct tumors:
  - Hemangiopericytoma (HPC)
  - Solitary fibrous tumor (SFT)

Agaimy et al. Histopathology. 2014: 65; 667-673.



Sinonasal SFT vs HPC

Histologically similar, genetically different

Solitary Fibrous Tumor SFT	Hemangiopericytoma HPC
SMA -	SMA +
CD34 +	CD34 -/weak
STAT6 + (NAB2–STAT6 fusion)	STAT6 -
Beta-catenin +/- (not mutated)	Beta-catenin + (B-cat mutation)
Dilated vessels	Hyalinized vessels

+/- Hypoglycemia *



Biphenotypic Sinonasal Sarcoma

(neural & myogenic differentiation)

- Monotonous, cellular spindle cell neoplasm
- Infiltrative
- A/W proliferative respiratory glands
- S-100 + ★
- SMA +
- t(2;4) PAX3-MAML3
- Locally aggressive but no metastases

Lewis JT, et al. Am J Surg Pathol. 2012: 36(4), 517–525.



Synovial Sarcoma

- Young adults (20-40)
- Highly cellular spindle cell neoplasm
- Fasciculated to storiform
- Dilated vessels (like SFT/HPC)

- Monophasic
  - Spindle only
- Biphasic
  - Spindle & Epithelioid

- t(X;18) SYT-SSX1 or 2
- TLE-1 overexpressed



Sinonasal Spindle Cell Tumors

Tumor	CD34	STAT6	SMA	Beta-catenin	S100	Genetics	Other
SFT	strong	+ nuclear	-	+/- nuclear pitfall	-	inv12 NAB2-STAT6	Stromal collagen
HPC	weak	-	+	+ nuclear	-	Beta-catenin mutation	Hyalinized vessels
BSS			+ (desmin often neg)	+/- nuclear pitfall	+ (SOX10 neg)	t(2;4) PAX3-MAML3	Proliferative respiratory epithelium
Syn Sarc				+/- nuclear pitfall		t(X;18) SYT-SSX1 or 2 TLE-1	+/- biphasic



High Grade / Blue Cell Tumors



High Grade Blue Cell Sinonasal Tumor	Neuroendocrine markers (Synaptophysin, Chromogranin)	Keratin (CAM5.2, AE1/AE3)	Squamous markers (p63, HMWK)	S-100	Other
Sinonasal undifferentiated CA (SNUC)	--	✚✚	--	--	High grade IDH2 mut
NUT midline CA	--	✚	✚	--	NUT nuclear stain + NUT translocation +
INI-1 deficient CA	--	✚	Focal +/-	--	INI-1 loss
HPV-related Multiphenotypic CA (AdCC-like)	--	✚ luminal stronger	✚ abluminal	+ var	p16, HPV33
Nasopharyngeal CA	--	✚✚	✚	--	Lymphoid stroma +/- EBV
Neuroendocrine CA	✚	✚✚	--	--	High grade
Olfactory neuroblastoma	✚	-/rare	--	+ sust	Neuropil / Rosettes Bland cytology
Ewing sarcoma / PNET	-/+	-/+ in Adamantinoma-like	--	-/+	CD99+, FLI-1+
Melanoma	--	--	--	✚	HMB-45, MelanA, SOX-10, MiTF



NUT Midline Carcinoma

- Head & Neck (35%) and Mediastinum
- High grade carcinoma with squamoid features
  - May be keratin / p63 positive
- Primitive, monotonous cells with
  - Round nuclei, Nucleoli
  - Amphophilic to clear cytoplasm
  - Occasional abrupt squamous differentiation
- NUT;BRD4or3 translocation (IHC/FISH)
- Median survival < 1 year

INI-1 Deficient Sinonasal Carcinoma



- Younger age (30s-40s)
- Aggressive
- Histology:
  - Basaloid nests with peripheral palisading
  - Rhabdoid cells
  - Intraepithelial spread
- IHC:
  - Pankeratin+
  - CK5/p63 patchy+
  - INI-1 lost
- FISH
  - SMARCB1 homozygous deletion



Am J Surg Pathol 2017;41:458–471)

---

a histologic mimic

HPV-Related Multiphenotypic Sinonasal Carcinoma

(HPV-related Carcinoma with Adenoid Cystic Like Features)

- Basaloid nests with solid, cribriform, tubular growth +/- surface involvement
- Myoepithelial, Ductal, Squamous differentiation
- p16+, HPV RNA ISH+ (more frequently HPV 33)
- Local recurrence > Distant mets > Nodal mets
- Low mortality, good prognosis


Am J Surg Pathol • Volume 41, Number 12, December 2017 *HPV-related Multiphenotypic Sinonasal Carcin*

PMID: 28877065

---

a histologic mimic

Adamantinoma-like Ewing Sarcoma

- Basaloid nests with peripheral nuclear palisading and prominent interlobular fibrosis
  - +/- keratin pearls
  - +/- myxoid stroma
- Positive for CD99, pankeratin, p40
  - +/- synaptophysin, S100, p16
- Positive for EWSR1-FLI1 rearrangement





Bishop JA, et al. Am J Surg Pathol. 2015 Sep;39(9):1267-74.

Larynx



Supraglottis

Glottis

Ventricle to 1 cm below TVC

False vocal cord

Ventricle

True vocal cord

Thyroid cartilage

Infraglottis

Larynx

- Reactive process
  - Vocal cord polyp
- Most common benign tumor
  - Papilloma
- Most common malignant tumor
  - SCC
- Most common neuroendocrine tumor
  - Atypical carcinoid
- Most common sarcoma
  - Chondrosarcoma

Vocal Cord Polyp

- "Singer's nodule"
- Related to vocal abuse
  - Singers, Teachers, etc
- Clinically diverse:
  - unilateral, bilateral, edematous
- Histologically similar:
  - Stromal edema, vascular proliferation, hyalinized amyloid-like material
  - Stromal cell atypia = NOT malignant



Squamous Papilloma


Papillomas



- Low Risk HPV 6/11
- Single
  - Adult Males
- Multiple
  - Children
  - "Recurrent respiratory papillomatosis"
  - Acquired in birth canal of mother
  - Multiple throughout larynx, tracheobronchial tree
- Occasionally lead to SCC, more so with radiation/smoking

Epidemiology of SCC

- M > F, Age 50s-60s
- Tobacco, Alcohol
- Most common site:
  - Glottis (in US)
  - Supraglottis (in most of western Europe)

- Prognosis: Stage-dependent
  - Cartilage invasion / bulky tumour volume → poor response to chemoradiotherapy
  - Histologic grade has limited impact

Squamous Cell Carcinoma


False vocal cord
True vocal cord

Invasive, keratinizing SCC

Staging


Epiglottis
Vestibule
Supraglottis
Epiglottis, AE folds, Arytenoids, False Cords
Glottis
True Cords to 1 cm below
Subglottis
To lower margin of cricoid
False vocal cord
Ventricle
True vocal cord
Anterior Commissure 12:00
Cricoid Cart
Trachea

- Is based on epicenter of tumor
  - Supraglottis
  - Glottis
  - Infraglottis
- And extent of invasion
  - Paraglottic space
  - Pre-epiglottic space
  - Strap muscles
  - Post-cricoid space


Pre-epiglottic Space
Paraglottic Space

- Looking at sagittal section through ANTERIOR larynx:

- Pre-epiglottic space
  - bounded by epiglottic cartilage posteriorly
  - filled with adipose and connective tissue.
- Paraglottic space
  - potential space deep to the vocal cords, filled with adipose and connective tissue
  - bounded by thyroid cartilage anteriorly
- If carcinoma invades anterior to paraglottic space, it may invade thyroid cartilage, strap muscles

Neuroendocrine Tumors of Larynx

- Atypical carcinoid is most common
  - M>F, a/w smoking
  - Most common in supraglottis
  - 50% nodal and distant mets
  - Surgery (chemo/radio insensitive)
  - Keratin +, Synap +, CMG +
  - May have: Calcitonin +, mCEA +, TTF-1 RARE

- Small cell carcinoma is second most common
- Rare: paragangliomas, typical carcinoids

Atypical Carcinoid

Carcinoid:
<2 mitoses / 10 hpf
Absent nucleoli,
pleomorphism, necrosis

Atypical Carcinoid:
2-10 mitoses / 10 hpf
Nucleoli present
Mild pleomorphism

Small Cell Carcinoma:
>10 mitoses / 10 hpf
Absent nucleoli
Necrosis

Nested growth
“Salt & Pepper” nuclei
Mitoses





Laryngeal Sarcomas

- Chondrosarcoma is most common
  - Rare but more common than benign chondroma
  - Any cartilaginous tumor > 2 cm with atypia should be regarded as chondrosarcoma
- Most common in cricoid cartilage
- Age 40+
- Usually low-int grade, low metastatic rate, curable by excision with clean margins

Chondrosarcoma





Salivary Glands

- Normal Histology
- Non-Tumors
  - Mucocele
  - Sialometaplasia
  - Sjogren Syndrome
- Benign Tumors
  - Pleomorphic Adenoma
    - Most common tumor
    - Most common benign tumor
  - Warthin tumor
    - Second most common benign tumor
- Malignant Tumors
  - Mucoepidermoid carcinoma
    - Most common malignant tumor
  - Adenoid Cystic
    - Second most common malignant tumor

Normal Salivary Gland

Gland	Low Power	High Power
Parotid SEROUS (zymogen granules)		
Submandibular & Minor MIXED		
Sublingual & Minor MUCINOUS		

Necrotizing Sialometaplasia

- Most common in palate
- Frequent ulceration
- ? Ischemia
  - following surgery, radiation
- Lobular configuration maintained
- Ductal structures
  - With lumina, +/- mucinous cells, intraluminal necrosis


http://static.wikidoc.org/7/72/Salivary_gland_enlargement_11.jpg





Oral Mucocele

- The most common minor salivary gland lesion
  - Mechanical trauma to the excretory duct
  - Soft, fluctuant nodule
  - Most often on lower lip > ventral tongue
  - Called RANULA on floor of mouth
- Pool of mucin surrounded by muciphages / fibrous pseudocyst
  - Adjacent atrophic / inflamed salivary tissue





Sjogren Syndrome

- Labial minor salivary gland

Langerman, A. J., Blair, E. A., Sweiss, N. J., & Taxy, J. B. (2007). Utility of Lip Biopsy in the Diagnosis and Treatment of Sjogren's Syndrome. The Laryngoscope, 117(6), 1004–1008.



4 mm^2 field
(10x obj)
Focus Score = 1
Grade = 3



TABLE I.
Histological Salivary Gland Grading System Developed by Chisholm and Mason[4] Compared With Focus Score Modification by Greenspan et al.[7]

Grade	Lymphocytes per 4 mm^2	Focus Score
0	No infiltrate	0
1	Slight infiltrate	0
2	Moderate infiltrate but less than one focus	0
3	One focus	1
4	More than one focus	2–12*

A "focus" is defined as an aggregate of 50 or more lymphocytes.
*The focus score is reported as raw number of lymphocytic foci per 4 mm^2, with a maximum of 12 (confluent foci).

Sclerosing Polycystic Adenosis / Adenoma (SPA)

- Rare mass-forming lesion
  - Predominantly in parotid gland
  - No age/sex predilection
- Proliferation of dilated ductal and acinar structures and fibrous stroma
  - Cells can be vacuolated, apocrine, clear, oncocytic, granular
- May recur after long follow up
- Recurring mutations in PTEN gene (?neoplastic)





https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3712090/



Tumor Trends

- Salivary carcinomas = 6% of head & neck cancers
- Benign tumors occur more often than malignant tumors
  - Most common location of all tumors: Parotid
  - Site with highest rate of malignant tumors: Sublingual
- The most common:
  - Tumor = Pleomorphic adenoma
  - Benign tumor = Pleomorphic adenoma > Warthin
  - Malignant tumor = Mucoepidermoid (adults & children) > Adenoid Cystic

Most Important For Diagnosis & Reporting

- Preoperative Biopsy (usually FNA, occasionally core)
  - Site: Primary salivary type tumor vs Metastasis
  - Grade: Low vs High
    - Low (benign or LG malignant) → Resection with margins
    - High → Resection with margins and neck dissection
  - Definitive diagnosis on biopsy can be challenging and histologic features overlapping
    - Can say "low grade salivary neoplasm" or "high grade carcinoma"
- Resection
  - Confirm Salivary type tumor vs Metastasis
  - Confirm Low vs High grade
  - Margins
  - T & N Stage

Benign Tumors +/– Malignant Counterparts

# of cell types, overall morphology, genetics

Benign	Cell Types	Malignant Counterpart	Genes
Pleomorphic Adenoma	1. Epithelial (ducts-tubules) 2. Myoepithelial • Stroma: Myxo-chondroid	Carcinoma ex pleomorphic adenoma Metastasizing pleomorphic adenoma	PLAG1 HMGA2
Basal cell adenoma	1. Epithelial (central, +/- squamoid) 2. Myoepithelial (peripheral palisading)	Basal cell adenocarcinoma	CYLD CTNNB1
Myoepithelioma	1. Myoepithelial	Myoepithelial carcinoma	EWSR1 in clear cell variants
Oncocytoma	1. Oncocytes, solid sheets/nests	Oncocytic carcinoma	None
Warthin tumor	1. Oncocytes, 2 rows, cystic-solid • Stroma: Lymphoid	Carcinoma / Lymphoma	None
Canalicular adenoma	1. Epithelial (Single row cuboidal) • Stroma: Prominent loose vascular	None	None



Pleomorphic Adenoma

Epithelial & Myoepithelial Cells:
- Ducts / single cells in extensive stroma
- Cellular sheets without much stroma



Pleomorphic Adenoma

- Things PA can do:
  - Recur > incomplete excision
    - Multiple small nodules in stroma / “buckshot”



Pleomorphic Adenoma

- Things PA can do:
  - Metastasize (“benign” metastasizing PA)
    - Bland cytology, up to 40% death
  - Transform (carcinoma ex PA)
    - Infiltrative, destructive growth (Invasive type)
      - Can resemble any type of salivary carcinoma
      - Most commonly CA-NOS or salivary duct
    - Nuclear atypia without infiltrative growth
      - Confined to tubules (In-situ)
      - Beyond tubules but Confined within capsule (Intracapsular)



Warthin Tumor

- Smokers / can be Bilateral
- Pain with infarction
- Cysts lined by bilayered oncocytic epithelium
  - Mucinous / Squamous met.
- Lymphoid stroma
  - PET-avid
  - Some are NON-clonal
    - ? NON-Neoplastic
  - 1% rate of malignant transformation
    - Carcinoma or Lymphoma



Malignant Tumors

Malignant	Cell types	Grade	High –Grade “Undifferentiated” Transformation	Genes
Mucoepidermoid carcinoma	1. Mucinous 2. Squamoid 3. Intermediate	LOW INT HIGH	YES	CRTC1-MAML2 t(11:19)
Adenoid cystic carcinoma	1. Epithelial 2. Myoepithelial	LOW INT HIGH	YES	MYB-NFIB t(6;9)
Epithelial-Myoepithelial carcinoma	1. Epithelial 2. Myoepithelial	LOW	YES	HRAS mut
Acinic cell carcinoma	1.	LOW	YES	NR4A3 or MSTATD3 fusion
Secretory carcinoma	1.	LOW	YES	ETV6-NTRK3 t(12;15)
Polymorphous adenocarcinoma Cribriform adenocarcinoma	1.	LOW	YES	PRKD1 mut PRKD1 fusion
Clear cell carcinoma	1.	LOW	YES	EWSR-ATF1 t(12;22)
Intraductal carcinoma	1. But intraductal so surrounded by myoeps	LOW	Not known to	RET fusion
Salivary duct carcinoma	1.	HIGH	Already High	AR, HER2



Mucoepidermoid Carcinoma

- 3 cell types

Mucinous
- Pale intracytoplasmic mucin

Intermediate
- Minimal pink-purple cytoplasm
- Small round nuclei

Squamoid
- Abundant eosinophilic cytoplasm
- Prominent cell borders

Mucoepidermoid Carcinoma


SOLID


CYSTIC

MEC – 4 Grading Systems

	Healy (Healy & Batsakis)	AFIP (Auclair & Goode)	Brandwein	Katabi
Intracystic	L: macro + micro cysts I: micro cysts + solid H: solid +/- micro cysts	2 (<20%)	2 (<25%)	L: predominantly cystic (>80%) I: predominantly solid H: any (usually solid)
PNI	H: present	2	3	n/a
Necrosis	n/a	3	3	L: absent I: absent H: present
Mitosis	L: rare I: few H: many	3 (4/10 H	3 (5/10 HP	L: 0-1/10 HPF I: 2-3/10 HPF H: 4+/10 HPF
Nuclear Anaplasia / Pleomorphism	L: absent/minimal I: slight/moderate H: considerable (including nucleoli)	4	2	L: no significant I: no significant H: any
Border / Invasive Front	L: broad /circumscribed I: uncircumscribed H: soft tissue/perineural/vascular invasion	n/a	2 (small nes	L: well circumscribed I: well circumscribed or infiltrative H: any (usually infiltrative)
LVI	H: present	n/a	3	n/a
Bony Invasion	n/a	n/a	3	n/a
Intermediate Cells	L: rare I: more common H: predominant	n/a	n/a	n/a
Stroma	L: extravasated mucin + fibrosis + CI I: fibrosis separating nests + CI H: desmoplasia, minimal CI	n/a	n/a	n/a
Architecture	L: daughter cysts from larger I: large duct less conspicuous H: variable architecture/cell morphology	n/a	n/a	n/a
Low	Qualitative	0-4	0	Qualitative
Intermediate		5-6	2-3	
High		7-14	4+	

Under-grades

Over-grades

High grade ☹

Low/Int Grade ☺

Significance of FUSION & GRADE:
-Tumors that are morphologically MEC have MAML2 fusion
-Can be high or low grade
-Tumors WITHOUT MAML2 fusion are likely NOT MEC

Adenoid Cystic Carcinoma



- 2 cell types
- 3 grades
  - Tubular (grade 1)
  - Cribriform (grade 2)
  - Solid (grade 3)
- Can undergo "dedifferentiation" / "high grade transformation"
- Perineural invasion
- MYB-NFIB fusion

Adenoid Cystic Carcinoma

- 2 cell types seen by IHC staining


Epithelial cells
- c-kit (CD 117)
- Inner cells


Myoepithelial cells
- p63
- Outer edge of cribriform space, touching stroma

Polymorphous (low grade) Adenocarcinoma (PAC)

- Epidemiology:
  - Most common location is palate
  - But most common tumor in palate is NOT PAC
  - Cribriform adenocarcinoma is morphologically similar, minor glands outside of palate, more propensity for nodal metastases
- Histology:
  - Single cell type: Pale oval nuclei (compare to AdCC)
  - Various architectural patterns (polymorphous): Lobular, papillary, cystic, cribriform, trabecular, ductal, single file, whorling
  - Infiltrative + Perineural invasion
  - S-100+, p40-
  - Can be high grade
- Genetics:
  - PRKD1 mutation



Polymorphous Adenocarcinoma


No myoeps evident around tubules



PAC vs AdCC vs PA

PAC
AdCC
PA
Polymorphous low-grade adenocarcinoma
Adenoid cystic carcinoma
Pleomorphic adenoma
A
B
C
D
E
F
G
H
I
Hematoxylin and eosin
p63
p40

Rooper et al. Head and Neck Pathol. 2015 Mar; 9(1): 78-94.



Acinic Cell Carcinoma

- Resemble serous acinar cells, but grows in sheets without ducts or fat
- +/- prominent lymphoid stroma
- Granular cytoplasm
- Considered low grade
  - Aggressive features: >2/10 mitoses, necrosis, LVI, PNI, pleomorphism, EPE
  - Can undergo "dedifferentiation" / "high grade transformation"

PAS (Periodic Acid Schiff) Stain:
- Stains glycogen and granules

Diastase:
- Digests out glycogen sugars
- Remaining granules are ZYMOGEN



Acinic Cell Ca
Adenoid Cystic Ca – Solid (Grade 3)

"High grade transformation"
Looks the same for AdCC, AcCC, etc



(Mammary Analogue)

Secretory Carcinoma

- Resembles secretory carcinoma of breast:
  - Stains with Mammaglobin and S-100
  - Carries same translocation (>90%):
    - ETV6-NTRK3 -> tyrosine kinase fusion protein (absent in Acinic Cell)
  - Can undergo high grade transformation

PAS-D stains secretions NOT zymogen granules

* Other tumors that diffusely stain with S100
- MASC
- PLGA
- LGCC

Classic microcystic pattern
Eosinophilic secretions
Monotonous, round-oval nuclei
Papillae
Intracytoplasmic vacuoles



Salivary Duct Carcinoma

- Epidemiology:
  - De novo or Ca-ex-PA
  - Aggressive
- Histology:
  - Resembles high grade ductal carcinoma of breast: large nests with peripheral cribriforming and central necrosis
  - PNI and LVI are common
- IHC/Genetics:
  - Androgen Receptor +
    - Although ER/PR neg
  - Her-2-neu amplification in some

Use AR/Her-2 for Rx not for Dx
- Other HG Carcinomas can have expression

Apocrine morphology
Cribriforming / Necrosis
AR
Her2
Micropapillary
PNI



Epithelial Myoepithelial Carcinoma

- Epidemiology:
  - 1% of salivary gland tumors, mostly parotid
  - 14% metastasis, < 10% death
- Histology:
  - Bi-layered tubules with clear myoepithelial (outer) cells
  - Solid clear myoepithelial areas

Basal Cell Adenoma / Adenocarcinoma

- Epidemiology:
  - Rare, mostly parotid
- Histology:
  - Epithelial & Myoepithelial cells
  - Solid, trabecular, tubular architecture
  - Peripheral palisading
  - Globules and bands of PINK hyalinized stroma
  - Beta-catenin nuclear +
- Genetics:
  - CTNNB1 mut in adenoma
  - CYLD1 mut in adenomas/carcinomas
    - Also implicated in familial dermal cylindromas



Am J Surg Pathol 2016;40:1143–1150

(Hyalinizing) Clear Cell Carcinoma

- Epidemiology:
  - Intraoral minor salivary glands
  - Intraosseous (Odontogenic CCC)
- Histology:
  - Monomorphous, clear cells with wrinkled nuclei
  - Pink hyaline stroma: thin septae or dense bands
- IHC: Squamous-like:
  - Pos: Keratin, HMWK, p63/p40
  - Neg for other myoepithelial markers: S-100, GFAP, SMA, calponin
- Genetics:
  - EWSR-ATF1
  - Identical to clear cell odontogenic carcinoma



Think about HCCC when you see a tumor with myoepithelial-like morphology but with only squamous (not myoep) markers

Ear & Temporal Bone

- Normal
- Chronic otitis media
- Cholesteatoma
- Tumors
  - Jugulotympanic Paraganglioma
  - Middle Ear Adenoma
  - Ceruminous Adenoma
  - Papillary Endolymphatic Sac Tumor

Normal Ear Histology

Middle Ear Mucosa
- Cuboidal

Tympanic Membrane
- Stratified Squamous



Chronic Otitis Media





- Chronic inflammatory cells
- Epithelial invaginations forming glands
  - Not normally found in middle ear

Cholesteatoma

- Squamous epithelium from external canal migrates onto eardrum
- Destructive invasion into drum / mastoid
- Related to chronic infections
- Morbidity if deep into skull



Jugulotympanic Paraganglioma

"glomus tympanicum"

- * Most common tumor of middle ear
- NE markers +
- Keratin –
- S-100 + in sustentacular cells
- May be locally infiltrative







synaptophysin



S-100

An aside on paragangliomas

- In the head & neck they occur:
  - Carotid body 60%
  - Middle ear 30%
  - Vagus/Larynx
- Spectrum of metastatic potential
  - No good morphologic criteria / No histologic scoring system
  - Not diagnosed as benign or malignant
  - Only "Paraganglioma" or "Metastatic paraganglioma"
- The best predictor of behavior
  - Is germline SDH mutation (succinate dehydrogenase)
  - Highest risk of aggressive behavior is SDHB mutation (30% metastasis)
  - Genes are A, B, C, D, AF2
- Cases should be screened with SDHB stain
  - Catches mutations in ANY SDH gene



Absent (or very weak) cytoplasmic SDHB staining should be worked up for germline mutation.



Adenomas

Middle Ear

- Glands/Trabeculae
- NO myoepithelial layer
- Most express neuroendocrine markers + keratins

Ceruminous

- Unencapsulated
- Inner oncocytic layer
- Outer myoepithelial layer





Papillary Endolymphatic Sac Tumor

- Inner ear (temporal bone)
- Associated with VHL
  - Hemangioblastomas, renal cysts and tumors, pancreatic cysts and tumors, pheochromocytomas, epididymal / ovarian cystadenomas
- Hearing loss
- Local invasion
- No metastasis





Thank you!

Questions? Comments?
Multiple Choice Questions to Follow

University of Chicago Photographic Archive, [apf6-02100], Special Collections Research Center, University of Chicago Library.

1. A 45 year old female is noted to have an erythematous patch on her posterolateral tongue. An excisional biopsy is performed. One of the sections demonstrates the pictured finding. What is the best diagnosis?

A. Mild squamous dysplasia
B. Squamous cell carcinoma in situ
C. Melanoma in situ
D. Paget's disease
E. Normal





1. A 45 year old female is noted to have an erythematous patch on her posterolateral tongue. An excisional biopsy is performed. One of the sections demonstrates the pictured finding. What is the best diagnosis?

A. Mild squamous dysplasia
B. Squamous cell carcinoma in situ
C. Melanoma in situ
D. Paget's disease
E. Normal

This image depicts a normal taste bud. Taste buds are nodular clusters of pale spindled epithelial cells within squamous epithelium. They are positive for CK7, while adjacent squamous epithelium is negative.



2. What is the most common location for this lesion?

A. Parotid gland
B. Palate
C. Base of tongue
D. Sinonasal cavity
E. Middle Ear



2. What is the most common location for this lesion?

A. Parotid gland
B. Palate
C. Base of tongue
D. Sinonasal cavity
E. Middle Ear

These images depict a polymorphous adenocarcinoma (low grade), which most commonly occurs on the palate.

It is strongly S100 positive and p40 negative (confirming absence of two cell-types).



3. The pictured image is from a curettage of an intra-osseous lesion. What is the most likely associated clinical finding?

A. Cutaneous basal cell carcinoma
B. Desmoid tumor
C. Nerve pain
D. Root canal
E. Unerupted molar



3. The pictured image is from a curettage of an intra-osseous lesion. What is the most likely associated clinical finding?

A. Cutaneous basal cell carcinoma
B. Desmoid tumor
C. Nerve pain
D. Root canal
E. Unerupted molar

This lesion is an ameloblastoma, with both cystic and infiltrative "mural" involvement. Ameloblastomas, Dentigerous cysts, and OKCs are often associated with unerupted molars. Other than pure cystic/intracystic ameloblastomas, all are treated by hemimandibulectomy.



4. A 35 year old female presents with an exophytic lesion on the mandibular gingiva / floor of mouth. What is the most likely diagnosis?

A. Peripheral giant cell granuloma
B. Pyogenic granuloma
C. Giant cell tumor of bone
D. Aneurysmal bone cyst
E. Brown tumor

4. A 35 year old female presents with an exophytic lesion on the mandibular gingiva / floor of mouth. What is the most likely diagnosis?

A. Peripheral giant cell granuloma
B. Pyogenic granuloma
C. Giant cell tumor of bone
D. Aneurysmal bone cyst
E. Brown tumor

This lesion is a giant cell reparative granuloma, peripheral type (located in the gingiva rather than within bone). It shows surface ulceration with an underlying lobular stromal proliferation of giant cells, spindle cells, and extravasated red blood cells.





5. A 56 year old male presents to ENT clinic with a neck mass. FNA shows metastatic squamous cell carcinoma. The patient is taken for panendoscopy to identify the primary site. The surgeon sees a fullness in the base of tongue and sends biopsies (pictured) for frozen section. What is the best diagnosis?

A. No evidence of neoplasia
B. Severe dysplasia
C. Carcinoma in situ
D. Squamous cell carcinoma, nonkeratinizing
E. Invasive squamous cell carcinoma, keratinizing



5. A 56 year old male presents to ENT clinic with a neck mass. FNA shows metastatic squamous cell carcinoma. The patient is taken for panendoscopy to identify the primary site. The surgeon sees a fullness in the base of tongue and sends biopsies (pictured) for frozen section. What is the best diagnosis?

A. No evidence of neoplasia
B. Severe dysplasia
C. Carcinoma in situ
D. Squamous cell carcinoma, nonkeratinizing
E. Invasive squamous cell carcinoma, keratinizing



Nonkeratinizing squamous carcinoma of the oropharynx (base of tongue and tonsil) is thought to originate in the deep tonsillar crypts, where the basement membrane is discontinuous and lymphatics exist in close association with the epithelium. HPV-related neoplasia of the tonsillar crypts should be considered invasive, discarding the classifier in situ and eliminating the concept of "crypt dysplasia."

This page was intentionally left blank.

Bone Pathology

Pathology Board Review
Osler Institute
April 2021

Nicole A. Cipriani, MD
Nicole.Cipriani@uchospitals.edu

AT THE FOREFRONT
UChicago Medicine

Outline

- Normal tissues
- Non-tumor lesions of bone
- Lesions of synovium
- Tumors **
  - DDx by histology

Normal

Ossification

Endochondral
- Replacement of hyaline cartilage template with bone
- Forms most bones: long bones of limbs, pelvis, ribs, and vertebral column


Epiphyseal plate

Intramembranous
- Direct conversion of mesenchymal tissue into bone
- Forms flat bones: mandible, maxilla, clavicles, frontal and parietal bones, parts of temporal and occipital bones

Bone


Periosteum
Cortex
Marrow
Immature WOVEN bone
Mature LAMELLAR bone

Cartilage

Types of cartilage:
- Hyaline (most prevalent):
  - Articular surfaces, larynx, trachea, nasal septum
- Fibro:
  - IV discs, ligaments
- Elastic:
  - External ear, epiglottis


Elastic Cartilage


Fibrocartilage


Hyaline Cartilage



Connective



Synovium

Synovium
- Scaffold of fibrous, adipose, vascular tissue
- Condensation of modified histiocytes / fibroblasts at surface (synoviocytes)

Non-Neoplastic Bone & Joints



Osteoarthritis

- Gradual degeneration of articular surface:
  - Cracking/fraying
  - Erosion
  - Eburnation
  - Subchondral cysts/sclerosis

Don't need to "tap" for eburnation – you can see it when you cut across the surface



AVN:Avascular Necrosis

- Fracture, steroids, EtOH, ALL
- Necrosis under articular surface
- Gross:
  - Opaque white wedge-shaped area
  - Uplifting of cartilage
- Histo:
  - Zone of necrosis of bone and marrow (very pink, loss of nuclei)

Legg-Calve-Perthes = idiopathic hip avascular necrosis in children



Rheumatoid Arthritis

- Rheumatoid Factor, HLA DR4
- Hyperplastic synovium with lymphoplasmacytic infiltrate
- Pannus: surface erosion with local invasion of synovium, fibroblasts

Figure 1: **Haematoxylin and eosin stains of joint tissues from patients with rheumatoid arthritis**
A: inflamed synovium with hyperplasia of synovial lining layer outlined by arrows. Deep in lining layer is a dense inflammatory infiltrate (original magnification ×40). B: synovial pannus invading bone and cartilage (original magnification ×100).

904 THE LANCET • Vol 358 • September 15, 2001



Fracture Callus

- Differential diagnosis on small biopsy = neoplasm!
  - Initial hemorrhage and bony fragmentation
  - Granulation tissue
  - Fibrocartilaginous matrix
  - Ossification to woven bone
- Admixture of vascular, fibrous, cellular cartilaginous, bony tissue with osteoblast rimming



Osteomyelitis

Acute

- Staph aureus
- Neutrophils, dead bone
- Sequestrum: dead bone in center
- Involucrum: subperiosteal new bone surrounding sequestrum

"Chronic"

- Lymphoplasmacytic inflammation, marrow fibrosis
- Nonspecific so must clinically correlate
- CRMO: Chronic recurrent multifocal osteomyelitis
  - Culture negative, multiple bones, relapsing inflammation ? autoimmune



Paget Disease

- Bone deforming disease, idiopathic
  - osteoclast resorption followed by compensatory increased osteoblast activity leading to unstructured and unstable bone
- Age >50, M>F
- Pelvis > femur, spine > skull
- Serum alk phos elevated
- Stages:
  - Initial stage: Osteoclastic
    - Osteoclast predominant (mimics brown tumor)
    - Stromal vascularity
  - Mid stage: Mixed pattern
    - Osteoclast + Osteoblast
    - Stromal vascularity
  - End stage: Sclerotic
    - Mosaic-pattern "reversal lines" in bone
    - Stromal fibrosis
- Treatment: bisphosphonates
- Secondary sarcomas rare (osteosarcoma second peak in elderly!)



Gout
Uric Acid

- Pale color +/- brown
- Usually surrounded by histiocytes/giant cells
- Needle shape
- * FIX IN ALCOHOL to retain crystals

Pseudogout
Calcium Pyrophosphate

- Purple or pale color
- Not surrounded by histiocytes
- Rhomboid shape
- No special processing



Joint Replacement

- Failure of joint replacement may be due to infection or mechanical failure
  - "Poly count" on frozen sections

Infection
5+ neutrophils/HPF in 5+ fields
- exclude fibrin/inflammatory exudate
- exclude neutrophils within vessels
- ... 43% sensitive and 97% specific for infection compared to culture

Mechanical Failure
"Metal on metal"
- Gray-black fragments, often within histiocytes
- Foreign body giant cells

Abdul-Karim FW, et al. Mod Pathol. 1998 May;11(5):427-31. PMID: 9619594.

J Clin Pathol 2012;65:409e418. doi:10.1136/jclinpath-2011-200398



Tenosynovial Giant Cell Tumors

Tenosynovial Giant Cell Tumor, Localized Type	Tenosynovial Giant Cell Tumor, Diffuse Type
"Giant cell tumor of tendon sheath"	"Pigmented villonodular synovitis (PVNS)"
Adults, F>M, FINGERS (mostly near joint) Well-circumscribed mass	Younger adults, F>M, KNEE (mostly in joint) Ill-defined intra/peri-articular lesions

1. Mononuclear cells (synovial-like) (CD68, CD163+)
2. Foamy histiocytes
3. Osteoclast-like giant cells (less common in diffuse type) (CD68, CD45+)
4. Hemosiderin

CSF1 rearrangement // May recur (more in diffuse type) // Possible "Malignant" variant

How to think about bone tumors

RADIOLOGY!

- Age
  - <30 (skeletally immature)
  - >30 (skeletally mature)
- Location in body
  - Axial skeleton, where?
  - Appendicular skeleton, where?
- Location in bone
  - Epiphysis (skeletally immature)
  - Metaphysis
  - Diaphysis
- Benign vs Malignant
  - Frozen section – worst thing to do is call malignant benign, because they will curet and contaminate the field
  - Caution is your friend

Age

Blue=Benign
- More benign things in kids

Red=Malignant
- More malignant things in adults



http://www.radiologyassistant.nl/en/494e15cbf0d8d

Location in Bone


Adult
Child
Diaphysis
Metaphysis
Diaphysis
Metaphysis
Epiphysis
Physis
(growth plate)
"Epiphyseal Line"
Endochondral oss.

© The Royal Children's Hospital, Melbourne, Australia

Epiphysis

CHILD (skeletally immature)

Epiphysis (Marrow):
- Chondroblastoma
- Infection
- GCT

ADULT (skeletally mature)

Epiphysis:
- GCT
- Infection
- Clear Cell Chondrosarcoma


Articular cartilage
Epiphysis
Epiphyseal line
Metaphysis
Spongy bone (cancellous)
Compact bone (cortical)
Medullary cavity
Diaphysis
Endosteum
Periosteum
Metaphysis
Epiphysis
Articular cartilage

Metaphysis

CHILD (skeletally immature)

Physis/Metaphysis:
- Osteosarcoma
- GCT

Metaphysis:

Marrow:	Cortex:
Enchondroma	Osteochondroma
UBC	ABC
CMF	NOF
	Osteoid Osteoma

ADULT (skeletally mature)

Metaphysis (Marrow):
- GCT
- Enchondroma/Chondrosarcoma
- Metastasis
- Myeloma/Lymphoma


Articular cartilage
Epiphysis
Epiphyseal line
Metaphysis
Spongy bone (cancellous)
Compact bone (cortical)
Medullary cavity
Diaphysis
Endosteum
Periosteum
Metaphysis
Epiphysis
Articular cartilage

Diaphysis

CHILD (skeletally immature)

Diaphysis:

Marrow:	Cortex:
Ewing	Adamantinoma
EG/LCH	Osteofibrous dysplasia
Fibrous dysplasia	

ADULT (skeletally mature)

Diaphysis (Marrow):
- Metastasis
- Myeloma/Lymphoma


Articular cartilage
Epiphysis
Epiphyseal line
Metaphysis
Spongy bone (cancellous)
Compact bone (cortical)
Medullary cavity
Diaphysis
Endosteum
Periosteum
Metaphysis
Epiphysis
Articular cartilage



Facial/Skull bones:
Fibrous dysplasia
Ossifying fibroma
GC reparative granuloma
Adamantinoma
Mesench. Chondrosarc
Langerhans histiocytosis

Ribs:
Fibrous dysplasia
Mesench. Chondrosarc
ABC

Vertebrae:
Osteoblastoma (posterior)
Chordoma (sacrum/skull base)

This diagram is the MOST COMMON site for each tumor. They may occur in locations other than those listed.

Proximal Humerus:
Periosteal chondroma
UBC

Small bones (hands, feet):
Enchondroma

Pelvis/Proximal Femur:
Osteoid osteoma (femoral neck)
Fibrous dysplasia
UBC
Chondrosarc
Clear cell chondrosarc
Ewing's sarcoma (shaft fem)

Around the Knee:
GCT (epi)
Chondroblastoma (epi)
ABC (met)
CMF (met)
NOF (met)
Exostosis (osteochondroma)
Osteosarc

Tibia:
Adamantinoma
Osteofibrous dysplasia

Location in Body

Cartilaginous, @ Surface

- Osteochondroma
- BPOP
- Periosteal Chondroma
- Periosteal Chondrosarcoma





Measure cartilage cap

What's wrong with this osteochondroma?



Benign cartilaginous neoplasm
In endochondrally-ossified bones
With bone induction and marrow continuity

Osteo-chondroma - Exostosis	• Most common primary bone tumor • Endochondrally ossified bones (thought to arise from growth plate) • Teens/20s, M>F , long bones • Single or Multiple • Multiple hereditary exostosis (AD, EXT1, 2, 3 mutation) ↑risk of chondrosarcoma • Stop growing w/ nl growth plate	• Metaphysis • Epiphysis rare (Trevor disease) • Cartilage is the neoplastic component, induces underlying bone growth • Cortex and marrow continuous w/ cortex & marrow space	Organized arrangement of: 1. Outer fibrous periosteal layer 2. Cartilage cap (always measure greatest thickness; thicker have higher risk of chondrosarc transformation) • Chondrocytes at base of cap in columns (like normal growth plate) 3. Undergoes endochondral ossification below 2ary Chondrosarcoma grows outward Cauliflower-like



Reactive proliferation
Of bone, cartilage, fibrous tissue
Usually on small phalangeal bones

BPOP (bizarre parosteal osteochondromatous proliferation) - Nora lesion - Subungual exostosis	• Not a neoplasm • Reactive proliferation / trauma? • All ages, both genders	• Metaphysis or Diaphysis • 75% small bones hands/feet • No continuity with marrow cavity (intact cortex)	Disorganized arrangement of: 1. Cartilage, hypercellular 2. Bone, endochondral ossification of cartilage 3. Spindle cells, hypercellular, mitoses



Periosteal Chondroma

- Rare
- All ages, M>F
- Tubular bones
- May erode cortex / bulge into the underlying marrow cavity, but there is always a border of intervening intact cortex
- Bland cartilage, no endochondral ossification
- Periosteal (juxtacortical) chondrosarcoma
  - Larger size (> 5 cm)
  - Absence of radiologic evidence of solid periosteal new bone buttressing
  - Higher grade cytology

Cartilaginous, in Marrow

- Metaphysis/Diaphysis
  - Enchondroma
  - Conventional Chondrosarcoma
  - CMF
- Epiphysis
  - Clear Cell Chondrosarcoma
  - Chondroblastoma



Intramedullary hyaline cartilage lobules
With peripheral ossification
WITHOUT entrapment of lamellar bone
+/- IDH1/2 mutation

Enchondroma	• Endochondrally ossified bones • Small bones of hands feet (60%) • Long bones (femur, humerus) • Ollier's disease: multiple enchondromas • Maffucci's syndrome: enchondromas w/ soft tissue spindle cell hemangiomas • Metachondromatosis: enchondromas and osteochondromas	• Metaphysis, Diaphysis • Intramedullary • Stippled opacities (calcifications) • Sclerotic rim b/c slow growing • +/- Inner cortical scalloping	• Hyaline matrix, Hypocellular, Bland cytology • Binucleation is OK • Myxoid matrix – think chondrosarcoma • More cellular/atypical in digits • Ossification at edges of lobules • Calcification within matrix • No invasion of bone (i.e. no encasement of mature lamellar bony trabeculae) • IDH1 or 2 mutation in 50% (also in chondrosarcoma)





Intramedullary hyaline-myxoid cartilage
WITH entrapment of lamellar bone
+/- IDH1/2 mutation

Chondro-sarcoma, conventional	• Middle age / older adults • Limb girdles (pelvis, prox femur, prox humerus) • Mass, pain, fracture	• Metaphysis, Diaphysis • Intramedullary • Lytic, opacities (arcs/spicules) • Endosteal scalloping • +/- Cortical destruction, soft tissue invasion	• Infiltrates around preexisting trabeculae +/- woven bone at edges • Hypercellular, nuclear enlargement, hyperchromasia, mitoses, necrosis, myxoid • (Binucleate cells not a great criterion) • Dediff (10 y older): abrupt transition from low to undiff spindle cell sarcoma, very poor prognosis • IDH1 and IDH2 gene mutations



Chondrosarcoma Grading

Cellularity / Nuclear Pleomorphism / Mitoses

Usually arise from grade 1-2

Locally aggressive
Treated with curettage

Worse prognosis
Treated with wide resection



Epiphysis of Adults
Cartilage islands, Clear cell background

Clear cell chondrosarcoma	• Adult Males • Long bones	• Epiphysis	• Lobulated, infiltrative • Clear cytoplasm, distinct boundaries, PAS-D+ glycogen • +/- Foci of conventional hyaline chondrosarcoma • Metaplastic bone +/- osteoblast rimming • May dedifferentiate



Jaws of Young Adults
Cartilage islands, Cellular spindled background
HEY1-NCOA2

Mesenchymal chondro-sarcoma	• Teens/20s (usu < 40) • Jaw > vertebrae, ribs, pelvis, long bones • Soft tissue	• Lytic with stippled calcification • Soft tissue invasion	1. Nodules of hyaline cartilage, may ossify 2. Spindle, ovoid, round cells with HPC-like vessels HEY1-NCOA2 fusion in 75% POOR PROGNOSIS (25% 10 year survival)

Giant-Cell Rich Tumor DDx

- NOF
- CMF
- Chondroblastoma
- GCT of bone
- ABC
- Telangiectatic osteosarcoma
- Osteitis fibrosa cystica (brown tumor)
- Central giant cell granuloma



Eccentric lesion in children
Storiform spindle cells + giant cells
Minimal-no intralesional bone

Non-ossifying fibroma (NOF) (larger, cortex & marrow) - fibrous cortical defect (limited to cortex) - metaphyseal fibrous defect - benign fibrous histiocytoma	• Most common cause of pathologic fracture in children • Teens, M>F • Around knee • May regress spontaneously • May be multiple in association with NF1	• Metaphysis • Cortex or cortex/medulla • Lobulated with sclerotic rim (soap bubble)	• Hypercellular plump spindle cells in a loose storiform pattern with admixed giant cells • Frequent hemorrhage w/ degenerative changes (hemosiderin, foamy histiocytes) • Necrosis if fractured • Looks like BFH/DF with Giant Cells!



Eccentric lesion in children
Lobules of myxoid cartilage
Peripheral hypercellularity + giant cells

Chondromyxoid fibroma (CMF)	• Teens-20s • M>F • Around knee (tibia > femur)	• Metaphysis • Eccentric marrow (rarely cortex) • Lucent, scalloped w/ sclerotic rim	• Lobulated myxoid cartilage w/ ↑cellularity at edge • Small spindle/stellate cells with oval nuclei and pink cytoplasm • Multinucleated giant cells at edge of lobules



Epiphyseal tumor in children
Giant cells + Mononuclear cells with wrinkled nuclei
Chicken wire calcification
Histone K36M

https://doi.org/10.1111/his.12945

Chondro-blastoma	• Teens/20s (skeletally immature) • M>F • Long bones (femur>tibia>humerus)	• Epiphysis • Lucent with sclerotic rim • +/- Cortical scalloping	1. Mononuclear chondroblasts w/ bean-shaped smudgy nuclei, abundant pink cytoplasm, well defined borders, retraction halos 2. Scattered multinucleated giant cells 3. Pink to blue chondroid matrix with chicken wire / stippled calcification Treated with curettage Histone H3F3A or H3F3B mutation (K36M) / IHC+



Lytic, intramedullary, +/- soft tissue invasion

Red-brown friable, well-circumscribed, may erode cortex
Usually we see these as curettage rather than resection



Oval, vesicular nuclei with small nucleoli

Lytic, around knee in young adults
Evenly-spaced giant cells + mononuclear cells with similar nuclei
Maturation with RANK-L inhibitors
Histone G34W

Am J Surg Pathol 2017;41:1059–1068.

Giant cell tumor of bone (GCT)	• 20-40s (skeletally mature) • F>M • Distal femur > proximal tibia > distal radius > sacrum • Local recurrence > curettage • Rare metastasis to lung (<5%) • Rare progression to sarcoma	• Epiphysis / Metaphysis • Intramedullary • Expansile, lytic • +/- Cortical destruction, soft tissue invasion • +/- Secondary ABC	• Numerous multinucleated giant cells evenly scattered throughout (bystanders) • Mononuclear cells with ill-defined borders and nuclei identical to giant cell nuclei (*neoplastic cells*) +/- mitoses, atypia • +/- Reactive bone formation, spindle cell areas, secondary ABC change • Express RANK ligand -> induce giant cells b/c they express RANK • RANK-L inhibitors (denosumab) induce bone formation/maturation • Histone H3F3A mutation (G34W) / IHC+ Mononuclear cells



Post-treatment
Loss of giant cells
Increase spindle cells/bone
DDx: LGOS, FD

Pre-treatment



Fluid fluid levels on MRI

Expansile lytic lesion
+/- Shell of reactive bone



Fluid-fluid levels, around knee in teens
Variably cellular spindle cells + osteoid + scattered giant cells jumping into cysts
USP6 rearrangement

Aneurysmal bone cyst (ABC)	• Teens, M=F
	• Around knee (distal femur and proximal tibia)
	• 70% are primary, 30% are secondary (arising in other tumors such as GCT, CB)
	• CDH11-USP6 t(16;17)
		• Metaphysis
		• Lytic, expansile
		• May be eccentric, central, cortical, or surface of bone
		• Cortex may be destroyed
		• Fluid-fluid levels common on MRI
			• Multiple cystic spaces separated by septa:
			  • Bland plump spindle cells without atypia or necrosis
			  • Multinucleated giant cells "jumping into the void"
			  • Osteoid and woven bone following contours of septae
			  • Occasional "blue bone"
			• Occasionally solid (hypercellular, mitotically active)
			• USP6 FISH (ABC, Nod Fasc, cellular fibroma of TS)



Telangiectatic osteosarcoma
- Low-power ABC-like
- High power malignant cells (hyperchromatic, angulated, nucleoli, atypical mitoses)
- Little osteoid


Lytic, loculated lesion in children
Thin collagenous wall

Unicameral bone cyst (UBC)	• Teens, young adults • M>F • Proximal humerus & femur • Pathologic Fx in 60%	• Metaphysis • Intramedullary • Lucent, trabeculated, thin cortex • Not usually fluid-fluid levels	• Thin cyst wall with bland spindle cells, collagen, fibrin • +/- osteoclast giant cells, foamy histiocytes, hemosiderin • Mimics ABC in cases of fracture


UBC
ABC
- USP6
Bone Cysts
Ganglion
- a/w OA, myxoid, fibrous
Epidermoid

Other giant cell rich lesions

Brown tumor of Hyperparathyroidism (osteitis fibrosa cystica)	Giant cell reparative granuloma (central / peripheral giant cell lesion)
PTH activates osteoblasts -> activate osteoclasts via RANKL-RANK interaction	Single or Symmetric jaw enlargement Cherubism (AD, SH3BP2 gene 4p16.3)

Lobulated growth of mononuclear cells + clustered osteoclast like giant cells
Richly vascular, hemosiderin, RBCs


Bony resorption nearby

Bone forming

- Osteoma
- Osteoid osteoma
- Osteoblastoma
- Osteofibrous dysplasia (Adamantinoma)
- Fibrous dysplasia
- Low grade osteosarcoma
- High grade (conventional) osteosarcoma



Osteoma	• Adults • Craniofacial (Frontal, Ethmoid) • Multiple in Gardner syndrome	• Cortical • Radiodense	• Ovoid • Resembles dense cortical bone (lamellar and woven with haversian canals)


Intracortical with nidus



Young adult with night pain relieved by NSAIDs
Small <2cm cortical lesion with nidus
Woven bone with rimming

Osteoid osteoma	Teens/20s, M>F Long bones (prox femur), posterior spinal elements Night pain relieved w/ NSAIDs Treated with radiofrequency ablation (RFA)	Diaphysis, metaphysis Most cortical < 2 cm (vs osteoblastoma) Radiolucent with central nidus (targetoid)	Woven bone with osteoblast rimming and loose fibrovascular stroma Peripheral sclerotic bone Biopsy often nondiagnostic (clinical Dx -> RFA)



Young adult with pain, less drug relief
Larger >2cm lesion
Woven bone with rimming +/- plump osteoblasts

Osteoblastoma	Teens/20s, M>F Long bones > posterior spinal elements (lamina, pedicles) > mandible Pain, Less response to NSAIDs	> 2 cm (vs osteoid osteoma)	Woven bone with osteoblast rimming More vascular stroma Atypical / Aggressive / Epithelioid OB: epithelioid osteoblasts



Anterior tibial cortex of young male
Fibro-Osseous Lesion: Variably cellular stroma + woven bone with rimming
Like ossifying fibroma of jaws

Osteofibrous dysplasia - Ossifying fibroma	Children, teens M>F Anterior cortex of tibia or fibula May regress > puberty	Diaphysis Cortex Bowing Multiple lytic lesions/cysts	Variably cellular fibrous stroma Irregular woven bony trabeculae with osteoblast rimming May progress to Adamantinoma


Well diff adamantinoma
(looks like OFD with clusters of epithelial cells)
Classic adamantinoma
(looks like ameloblastoma)

Anterior tibial cortex of young male
Epithelial islands with variable amounts of fibrous stroma
Like ameloblastoma of jaws

Adamant-inoma	Well-diff: <20 ; Classic: >20 M>F Tibia >> fibula Late metastasis to lung	Diaphysis Cortex Lytic, soap bubble +/- marrow/soft tissue extension	Well-diff: osteofibrous dysplasia like with inconspicuous keratin-positive nests Classic: predominant epithelial component that resembles ameloblastoma of jaws



Ground glass lesion with sclerotic rim
Fibro-Osseous Lesion: Monotonously cellular stroma + curvilinear woven bone without rimming
GNAS1 mutation in congenital cases

Fibrous dysplasia - Congenital: acquired mutation in GNAS1	Broad age range (usu before 40) Monostotic > Polyostotic Rib > femur > tibia > craniofacial - McCune-Albright: Polyostotic FD, endocrinopathy, cutaneous pigmentation - Mazabraud: FD, Soft tissue myxoma	Intramedullary Elongate/ovoid Well-circumscribed w/ sclerotic rim (more ill-defined in craniofacial) "Ground glass"	Monotonously cellular stroma, bland fibroblasts Curvilinear woven bone trabeculae without osteoblast rimming Appear to arise from stroma (Sharpey fibers) May be bone-poor, cystic, ABC-like, cartilaginous



Fibro-Osseous Lesion: Cellular stroma + long woven/lamellar bone without rimming
Mild cytologic atypia + peripheral infiltration
MDM2 amplification

Well-differentiated intramedullary osteosarcoma - Low grade central osteosarcoma	20s (Teens-40s), M=F Femur > Tibia	Metaphysis Intramedullary Ill-defined, cloud-like mineralization	Low-Moderately cellular fibrous stroma with mild cytologic atypia Long, parallel trabeculae of woven or lamellar bone Infiltrative into adjacent bone MDM2, CDK4 amplification (also in parosteal osteosarcoma, which is histologically similar)





Conventional Osteosarcoma	- Teens/20s, M>F
	  - Around knee (femur, tibia)
	- Second peak >50
	  - Pelvis, axial skeleton
	- Early chemo to treat micromets
		- Medullary, Metaphysis
		- Mixed lytic-blastic
		- Codman's triangle – periosteal reaction; angle created by cortex and elevated periosteum
			- Osteoid produced by malignant cells
			- Hyperchromasia, atypical mitoses
			- Variants: Osteoblastic, chondroblastic, fibroblastic, telangiectatic, giant cell rich, small cell
			- Most are high grade (Grade 2 or 3)
			- SATB2 can help differentiate true osteoblastic cells from mimics (cells in collagen)



Sometimes osteoid is scarce

But atypia should be obvious

Bone-forming, @ surface

- Parosteal osteosarcoma
- Periosteal osteosarcoma



- Heavily mineralized mass attached by a broad base to the underlying cortex



Low grade fibroblastic surface osteosarcoma
In young people
With MDM2 amplification

PARosteal osteosarcoma	- Most common surface OS - Young adults (peak age 20s), ?F>M - Posterior distal femur > proximal tibia > proximal humerus	- Metaphysis - Heavily mineralized mass attached by a broad base to the underlying cortex	- Broad woven bone trabeculae - Spindle cells with moderate atypia - +/- Malignant cartilaginous areas (nodule or cap) - 15% High grade spindle cell component (dedifferentiated) - MDM2 + by IHC, MDM2 amplification - 80% 10-year survival



- Fusiform surface mass with linear mineralization



Intermediate grade chondroblastic surface osteosarcoma
In young people
WITHOUT MDM2 amplification

PERIosteal osteosarcoma	• Rare • Teens-20s, F>M • Distal femur, proximal tibia > humerus, ulna	• Diaphysis > Metaphysis • Fusiform surface mass with linear mineralization	• May scallop / erode cortex into medullary canal • Predominantly chondroblastic (grade 1-2) osteosarcoma • Coarse lace like neoplastic bone • 85% overall survival



DDx

	Parosteal	Periosteal
Location	Distal femur Metaphysis	Distal femur Diaphysis
Gross	Sessile, broad base Nearly circumferential	Sessile, more fusiform Nearly circumferential
Radiology	Can invade marrow (50%)	Can invade marrow (25%)
Histo	Mostly fibroblastic component Minimal cartilage component MDM2 amplified	Mostly cartilaginous component No fibroblastic component No specific genetics
Grade	Usu low, can de-differentiate	Low-Int, can de-differentiate
Treatment	Local resection with clear margins +/- radiation for positive margins +/- Neoadjuvant CRT (controversial) if int-high grade periosteal	

Other things seen in Marrow



Langerhans Histiocytosis - Eosinophilic Granuloma - Histiocytosis X	• Children>Adults, M>F • Flat bones > long bones • Single system: single site or multi site • Multi system • Hand-Schüller-Christian disease (multiple lesions) • Letterer-Siwe disease (disseminated disease or visceral involvement)	• Langerhans cells with folded, grooved nuclei and pale-to-eosinophilic cytoplasm • Variable eosinophils, histiocytes, neutrophils, lymphocytes, and multinucleated giant cells • BRAF V600E mutation in most • CD1a, Langerin, and S100 positive • EM: Birbeck granules



Ewing Sarcoma	• Age 10-15, M>F • Diaphysis of long bones and flat bones of pelvis • Lytic marrow lesion with cortical destruction • Treated with chemotherapy and surgery • Chemotherapy = 75% 5-year survival	• small round blue cell tumor (alsoin soft tissue lecture) • sheets or irregular islands in dense fibrous tissue +/- organoid growth pattern and rosettes • oval nuclei with finely chromatin and inconspicuous nucleoli, scant eosinophilic or clear cytoplasm • +/- larger cells with irregular nuclear contours and prominent nucleoli (previously designated atypical or large cell variant of EWS=same prog.) • CD99, FLI1+ • t(11;22)(q24;q12) EWSR1-FLI1 85% • t(21;22)(q22;q12) EWSR1-ERG 5-10%



Ewing Like Sarcomas

BCOR-CCNB3 tumors	CIC-DUX4 tumors
• most occur in bone	• 2/3 of EWSR1-negative tumors • most occur in soft tissue
• heterogeneous small round and spindle cells • stroma may be myxoid	• sheets of small round cells +/- spindle cells • stroma may be myxoid • less monotonous than EWS • may have more cytoplasm
• usually patchy CD99 positive • stain for BCOR and CCNB3	• usually patchy CD99 positive • stain for WT1, ETV6, and CIC



Adults

Plasma Cell Neoplasms

- Plasmacytoma vs Multiple Myeloma
- Most common primary malignant bone tumor
- CD138, Kappa:Lambda

Metastasis

- More common in adults than primary bone tumors
- Breast, Lung, Thyroid, Kidney, Prostate

MALIGNANCIES THAT METASTASIZE TO BONE
BACON (BREAST CA)
LETTUCE (LUNG CA)
TOMATO (THYROID CA)
KOSHER PICKLE (KIDNEY CA, PROSTATE CA)
THE SPINE IS THE MOST COMMON SITE OF BONE METASTASES
www.medcomic.com
© 2013 Jorge Muniz

RCC tends to be Lytic
Prostate and Breast tend to be Sclerotic



Vascular

* More in soft tissue lecture

Hemangioma

- Usu asymptomatic, incidental
- Adults, F > M
- Vertebral bodies
- Gorham-Stout disease: massive osteolysis due to hemangiomas
- Capillary/cavernous vessels
- Fibrotic stroma

Epithelioid Hemangioendothelioma

- Peak Teens/20s
- Can be multifocal
- Extremities, pelvis, spine
- May recur, metastasize
- Chords/nests of epithelioid cells in myxoid matrix
- Intracytoplasmic lumina



Thank you!

Questions? Comments?
Multiple Choice Questions to Follow

round cell lesions
fibrous dysplasia
chondromyxoid fibroma
solitary bone cyst
non-ossifying fibroma
osteosarcoma
giant cell tumor
physis
chondroblastoma
enchondroma
< 30 years
> 30 years
FD
EG
Ewing EG
Osteoid osteoma
NOF
SBC
Osteo Chondroma
CMF
Enchond
ABC
Osteo sarcoma
Chondroblastoma
infection
Metastasis Myeloma Lymphoma
Metastasis Myeloma HTP
Enchondroma Chondrosarcoma
Geode
Giant CT
infection
RS



1. Pictured is a biopsy from a 15 year old female. What is the most likely site of origin of this tumor?

A. Mandible.
B. Rib.
C. Femur.
D. Vertebra.
E. Phalanx.



1. Pictured is a biopsy from a 15 year old female. What is the most likely site of origin of this tumor?

A. Mandible.
B. Rib.
C. Femur.
D. Vertebra.
E. Phalanx.

This tumor is a mesenchymal chondrosarcoma, composed of primitive blue cells and sparse myxo-chondroid matrix-rich areas. The most frequent location is craniofacial bones. Prognosis is poor with 25% 10 year survival.

2. A 17 year old male presents with leg pain. A lesion is found in the anterior tibial cortex. What is the most likely diagnosis?



A. Fibrous dysplasia.
B. Osteofibrous dysplasia.
C. Parosteal osteosarcoma.
D. Osteoid osteoma.
E. Osteoblastoma

2. A 17 year old male presents with leg pain. A lesion is found in the anterior tibial cortex. What is the most likely diagnosis?



A. Fibrous dysplasia.
B. Osteofibrous dysplasia.
C. Parosteal osteosarcoma.
D. Osteoid osteoma.
E. Osteoblastoma

Osteofibrous dysplasia is a benign fibro-osseous lesion composed of irregular spicules of woven bone with osteoblast rimming in a variably cellular fibrous stroma. It most frequently occurs in the anterior tibial cortex of teenage males.

3. Regarding the patient in question 2, what is the most likely long-term complication?



A. Progression to polyostotic disease.
B. Transformation to adamantinoma.
C. Metastasis to lung.
D. Development of sarcoma.
E. No adverse long-term complications expected.

3. Regarding the patient in question 2, what is the most likely long-term complication?



A. Progression to polyostotic disease.
B. Transformation to adamantinoma.
C. Metastasis to lung.
D. Development of sarcoma.
E. No adverse long-term complications expected.

In some patients, osteofibrous dysplasia transforms to Adamantinoma, the long bone equivalent of Ameloblastoma.

4. A 16 year old male has an exophytic lesion on the posterior surface of the femur. Immunostain for MDM2 is positive. What is the most likely diagnosis?



A. Well-differentiated intramedullary osteosarcoma
B. Parosteal osteosarcoma
C. Periosteal osteosarcoma
D. Fibrous dysplasia
E. Osteofibrous dysplasia

4. A 16 year old male has an exophytic lesion on the posterior surface of the femur. Immunostain for MDM2 is positive. What is the most likely diagnosis?



A. Well-differentiated intramedullary osteosarcoma
B. Parosteal osteosarcoma
C. Periosteal osteosarcoma
D. Fibrous dysplasia
E. Osteofibrous dysplasia

Parosteal osteosarcoma is present on the surface of bones and shows MDM2 amplification. Well-differentiated osteosarcoma is similar but located within the marrow.



5. A 22 year old female presents with knee pain. A tibial epiphyseal lesion is identified on X-ray. Frozen section is performed to guide therapy. What is the best treatment for this lesion?

A. Curettage.
B. Distal tibial resection.
C. Amputation.
D. Chemotherapy.
E. No further treatment.



5. A 22 year old female presents with knee pain. A tibial epiphyseal lesion is identified on X-ray. Frozen section is performed to guide therapy. What is the best treatment for this lesion?

A. Curettage.
B. Distal tibial resection.
C. Amputation.
D. Chemotherapy.
E. No further treatment.

Chondroblastoma is composed of giant cells and mononuclear cells with wrinkled nuclei. Chicken-wire calcification is often present. Curettage is the treatment of choice.

This page was intentionally left blank.

Soft Tissue Pathology

Pathology Board Review
Osler Institute
April 2021

Nicole A. Cipriani, MD
Nicole.Cipriani@uchospitals.edu

AT THE FOREFRONT
UChicago Medicine

Outline

- Overview of approach to soft tissue tumors
- Adipocytic Tumors
- Myogenic Tumors
- Vascular Tumors
- Nerve Sheath Tumors
- Fibroblastic / Myofibroblastic / Undifferentiated Tumors
  - Pattern-based

Soft Tissue Tumor Epidemiology

- Benign >> Malignant
- Benign soft tissue tumors:
  - Most common = Lipoma
  - Most frequent location = Superficial (99%)
  - Mostly small (< 5 cm)
- Malignant soft tissue tumors:
  - Most common = UPS, liposarcoma, leiomyosarcoma, myxofibrosarcoma, synovial sarcoma, MPNST
  - Most frequent location:
    - Extremity (75%) vs trunk/retroperitoneum
    - Deep (60%) vs superficial
  - 10% have metastases at presentation
  - Surgery is primary treatment +/- XRT, chemo

Soft Tissue Tumor Epidemiology

- Incidence varies with age:
  - Children: Embryonal rhabdo
  - Young adults: Synovial sarcoma
  - Elderly: UPS, liposarc, leiomyosarc, myxofibrosarc
- Radiation-induced sarcoma:
  - At least 50 Gy, median 10 yrs from exposure to diagnosis
  - Usually UPS in soft tissue, Angiosarc in skin
  - Increased risk in Retinoblastoma (RB1), Li Fraumeni (TP53), and NF1 patients

Diagnosis: It's Manageable!

Ways to think about soft tissue tumors:



Lineage of Differentiation
- WHO book
- Enzinger & Weiss



Histologic Pattern
- Hornick book



A combination of both
- your brain

Organization

Lineage of Differentiation	Histologic Pattern
- Adipocytic	- Spindle cell
- Fibroblastic/myofibroblastic	- Epithelioid
- Fibrohistiocytic	- Pleomorphic
- Smooth muscle	- Round cell
- Perivascular	- Biphasic
- Skeletal muscle	- Myxoid
- Vascular	
- Chondro-osseous	
- Nerve sheath	
- Uncertain/undifferentiated	

Basic Principles



Neoplasm vs "Reactive" Change

Benign vs Malignant

Grade:
Low (1) → Resect with clean margin and watch
High (2-3) → Chemo-RT pre or post-resection

Extent of Rx effect
(necrosis)

Margin

French FNCLCC Grading System



Data from Trajani M, Contesso G, Coindre JM, et al. Soft-tissue sarcomas of adults: study of pathological prognostic variables and definition of a histopathological grading system. *Int J Cancer.* 1984;33:37 42.

Tumor Differentiation	
Score 1	Sarcomas that closely resemble normal adult mesenchymal tissues
Score 2	Sarcomas for which histologic typing is certain
Score 3	Embryonal and undifferentiated sarcomas, synovial sarcoma, and sarcomas of uncertain differentiation
Mitotic Count	
Score 1	0–9 mitoses/10 hpf
Score 2	10–19 mitoses/10 hpf
Score 3	≥20 mitoses/10 hpf
Tumor Necrosis	
Score 0	No necrosis
Score 1	<50% tumor necrosis
Score 2	≥50% tumor necrosis
Histologic Grade (tumor differentiation + mitotic count + tumor necrosis)	
Grade 1 (low grade)	Total score: 2 or 3
Grade 2 (intermediate grade)	Total score: 4 or 5
Grade 3 (high grade)	Total score: 6, 7, or 8

FNCLCC, Fédération Nationale des Centres de Lutte Contre le Cancer; hpf, high-power field.



Table 1.2 Tumour differentiation score according to histological type in the FNCLCC grading system

Histological type	Differentiation score
Well-differentiated liposarcoma	1
Well-differentiated leiomyosarcoma	1
Malignant neurofibroma	1
Well-differentiated fibrosarcoma	1
Myxoid liposarcoma	2
Conventional leiomyosarcoma	2
Conventional MPNST	2
Conventional fibrosarcoma	2
Myxofibrosarcoma	2
Myxoid chondrosarcoma	2
Conventional angiosarcoma	2
High-grade myxoid (round cell) liposarcoma	3
Pleomorphic liposarcoma	3
Dedifferentiated liposarcoma	3
Rhabdomyosarcoma	3
Poorly differentiated/pleomorphic leiomyosarcoma	3
Poorly differentiated/epithelioid angiosarcoma	3
Poorly differentiated MPNST	3
Malignant Triton tumour	3
Synovial sarcoma	3
Extraskeletal osteosarcoma	3
Extraskeletal Ewing sarcoma	3
Mesenchymal chondrosarcoma	3
Clear cell sarcoma	3
Epithelioid sarcoma	3
Alveolar soft part sarcoma	3
Malignant rhabdoid tumour	3
Undifferentiated (spindle cell and pleomorphic) sarcoma	3

FNCLCC, Fédération Nationale des Centres de Lutte Contre le Cancer; MPNST, malignant peripheral nerve sheath tumour
Modified from {494}

- Grade predicts probability of metastasis / survival
- Does NOT predict local recurrence (margins*)

Grade is less meaningful in:

- Dediff liposarc
- Round cell liposarc
- Rhabdomyosarcoma
- Ewing sarcoma
- Alveolar soft part sarcoma
- Epithelioid sarcoma
- Clear cell sarcoma

Staging



Trunk and Extremities
___ pTX: Primary tumor cannot be assessed
___ pT0: No evidence of primary tumor
___ pT1: Tumor 5 cm or less in greatest dimension
___ pT2: Tumor more than 5 cm and less than or equal to 10 cm in greatest dimension
___ pT3: Tumor more than 10 cm and less than or equal to 15 cm in greatest dimension
___ pT4: Tumor more than 15 cm in greatest dimension

AJCC 8: T stage varies by site:
- Head & Neck
- Trunk & Extremities
- Abdomen & Thoracic Visceral Organs
- Retroperitoneum
- Orbit



___ pN0: No regional lymph node metastasis
___ pN1: Regional lymph node metastasis

Sarcomas RARELY metastasize to lymph nodes, except:
- alveolar rhabdomyosarcoma
- epithelioid sarcoma
- clear cell sarcoma



___ pM1: Distant metastasis
Specify site(s), if known: ______________________________

Margins

- Marginal excision = removal of the tumor and its pseudocapsule with minimal (or no) surrounding normal tissue
- Wide excision = removal of the tumor with adjacent normal (healthy) tissue surrounding the tumor
- Radical resection = removal of the tumor and the entire anatomic compartment in which it is located (e.g., the surrounding muscle groups)

- Margins should be taken as perpendicular sections
- The presence of an intact fascial plane at the margin should also be reported

General Concepts

- Morphology / Terminology
- Immunostains and what they mean

- Putting Morphology, Immuno, Genetics, & Clinical together



What to do when faced with an unknown spindle cell lesion?

Nuclear Morphology

Wavy / Pointed
Nerve sheath

Cigar / Blunt
Smooth Muscle

Vesicular / Nucleoli
Fibroblast / Myofibroblast



What to do when faced with an unknown spindle cell lesion?

Cytoplasm

Abundant
“epithelioid”

Pink, Eccentric
“plasmacytoid” / “rhabdoid”
(rhabdoid=more pink cytoplasm)

Wispy
“stellate”
fibroblast / myofibroblast



What to do when faced with an unknown spindle cell lesion?

Stroma

Fine, Blue-ish
myxoid

Dense, Pink
collagenous

Misc.
osteoid / chondroid



What to do when faced with an unknown spindle cell lesion?

Architecture

Haphazard
“tissue culture”

Whorls/Short Fascicles
“storiform”

Long Fascicles
“fasciculated” / “herringbone”



What to do when faced with an unknown spindle cell lesion?

Vessels

Thin-walled
Dilated/Branching
“Staghorn / HPC-like”

Anastomosing
Capillary network
“chicken wire”

Thick-walled
Muscular
-



What to do when faced with an unknown spindle cell lesion?

Other Features

Necrosis
Malignancy/Prior Bx

Mitosis
In context!
Typical/Atypical

Infiltrative VS
Pushing Border
-

Immunostains



Keratins

CAM5.2
AE1/AE3
HMW
CK5/6
Specific CKs
(EMA)

Keratins can stain:
Carcinoma
Myoepithelioma
Epithelioid sarcoma
Synovial sarcoma
~Schwannoma
~Ewing sarcoma

Neural Crest

S-100 (Nuc/Cyto)
SOX-10 (Nuc)

Melanocytic

Melan-A (Mart-1)
HMB45
MiTF (Nuc)

S100 can stain:
Melanoma
Adipocytes
Fatty tumors
Chondrocytes
Cartilaginous tumors
Nerve sheath tumors

Myogenic

Desmin
MSA
SMA
Calponin
Caldesmon
SMMHC
Myogenin (Nuc)
(MyoD1) (Nuc)

Desmin can stain:
Smooth muscle
Myofibroblast
Myoepithelium
Skeletal muscle

Vascular

CD31
CD34
D2-40
FLI-1 (Nuc)
ERG (Nuc)

Other

CD45 (LCA)
Vimentin
...the list goes on

Keratins

- CAM5.2 – low molecular weight
  - Non squamous epithelium
- HMW, CK5/6 – high molecular weight
  - Squamous, urothelial, myoepithelial
- AE1/AE3 – broad spectrum (low and high)
  - Most epithelium
- EMA – least specific
  - Epithelial and non-epithelial tissues

Neural Crest

S100
- Melanoma
- Nerve sheath (Schwann cells)
- Granular cell tumor
- Myoepithelium
- Fat/Fatty tumors
- Cartilage/Cartilaginous tumors

SOX-10
- Melanoma
- Nerve sheath (Schwann cells)
- Granular cell tumor
- Myoepithelium
- Salivary/Pancreas acini

Myogenic

	Desmin	MSA Muscle specific actin	SMA Smooth muscle actin	Calponin	Caldesmon	Smooth muscle myosin HC	Myogenin MyoD1 (nuclear)
Smooth muscle	+	+	+	+	+	+	-
Myo-epithelial	+	+	+	+	+	+	-
Myo-fibroblast	+/-	+	+	+/-	-	-	-
Skeletal muscle	+	+	-	-	-	-	+

What is a fibroblast/myofibroblast?
A spindle cell that you cannot prove is anything else by morphology and IHC.
If it has some myogenic markers, you can call it myofibroblast.

Vascular

- CD31 – more sensitive and specific
  - Endothelium
  - Macrophages, plasma cells, Histiocytic sarcoma...
- CD34
  - Endothelium
  - Solitary fibrous tumor, DFSP, Spindle cell lipoma...
- D2-40
  - Lymphatic* endothelium
  - Kaposi sarcoma, angiosarc, EHE, mesothelioma...
- FLI-1 – nuclear, benign and malignant
  - EWS, lymphocytes...
- ERG – nuclear, probably most sensitive/specific
  - Prostate cancer, EWS, Acute myeloid leukemia...

Be sure to check the cells that are staining with your marker!

Translocations

- Wow.
- There are a lot.
  - Some are specific to a tumor.
  - Some are not.
  - Some have associated IHC staining.
  - Some do not.
- We will discuss in context of DDx.



Sarcoma Type	Cytogenetic Alterations	Molecular Alterations
Alveolar rhabdomyosarcoma	t(2;13)(q35;q14)	*PAX3-FOXO1A* fusion
	t(1;13)(p36;q14), double minutes	*PAX7-FOXO1A* fusion
	t(2;2)(q35;p23)	*PAX3-NCOA1* fusion
	t(X;2)(q35;q13)	*PAX3-AFX* fusion
Alveolar soft part sarcoma	t(X;17)(p11;q25)	*TFE3-ASPL* fusion*
Aneurysmal bone cyst	t(16;17)(q22;p13)	*CDH11-USP6* fusion
Angiomatoid fibrous histiocytoma	t(12;16)(q13;p11)	*FUS-ATF1* fusion
	t(12;22)(q13;q12)	*EWSR1-ATF1* fusion*
	t(2;22)(q33;q12)	*EWSR1-CREB1* fusion
Clear cell sarcoma	t(12;22)(q13;q12)	*EWSR1-ATF1* fusion*
	t(2;22)(q33;q12)	*EWSR1-CREB1* fusion
Desmoid fibromatosis	Trisomies 8 and 20 and loss of 5q21	*CTNNB1* or *APC* mutation*
Desmoplastic small round cell tumor	t(11;22)(p13;q12)	*EWSR1-WT1* fusion
Dermatofibrosarcoma protuberans	Ring form of chromosomes 17 and 22	*COL1A1-PDGFB* fusion
	t(17;22)(q21;q13)	*COL1A1-PDGFB* fusion
Endometrial stromal sarcoma	t(7;17)(p15;q21)	*JAZF1-SUZ12* fusion
	t(6;7)(p21;7p15)	*JAZF1-PHF1* fusion
	t(6;10)(p21;p11)	*EPC1-PHF1* fusion
Epithelioid hemangioendothelioma	t(1;3)(p36;q25)	*WWTR1-CAMTA1* fusion
Ewing sarcoma/primitive neuroectodermal tumor (PNET)	t(11;22)(q24;q12)	*EWSR1-FLI1* fusion
	t(21;22)(q12;q12)	*EWSR1-ERG* fusion
	t(2;22)(q33;q12)	*EWSR1-FEV* fusion
	t(7;22)(p22;q12)	*EWSR1-ETV1* fusion
	t(17;22)(q12;q12)	*EWSR1-E1AF* fusion
	inv(22)(q12;q12)	*EWSR1-ZSG* fusion
	t(16;21)(p11;q22)	*FUS-ERG* fusion*
	t(19;der)ins.inv(21;22)	*EWSR1-ERG* fusion
	t(17;22)(q12;q12)	*EWSR1-ETV4* fusion
	t(6;22)(p21;q12)	*EWSR1-POU5F1* fusion
	t(1;22)(q36.1;q12)	*EWSR1-PATZ1* fusion
	t(2;22)(q31;q12)	*EWSR1-SP3* fusion
	t(20;22)(q13;q12)	*EWSR1-NFATc2* fusion
	t(2;16)(q35;p11)	*FUS-FEV* fusion
	t(15;19)(q14;p13.1)	*BRD4-NUT* fusion*
	t(4;19)(q35;q13)	*CIC-DUX4* fusion
Extraskeletal myxoid chondrosarcoma	t(9;22)(q22;q12)	*EWSR1-NR4A3* fusion
	t(9;17)(q22;q11)	*TAF2N-NR4A3* fusion
	t(9;15)(q22;q21)	*TCF12-NR4A3* fusion
	t(3;9)(q11;q22)	*TFG-NR4A3* fusion
Fibrosarcoma, infantile/cellular mesoblastic nephroma	t(12;15)(p13;q26)	*ETV6-NTRK3* fusion*
	Trisomies 8, 11, 17, and 20	
Gastrointestinal stromal tumor	Monosomies 14 and 22; deletion on 1 p	*KIT* or *PDGFRA* mutation; rarely *BRAF*, *SDHA*, *SDHB*, *SDHC*, or *SDHD* mutations
Inflammatory myofibroblastic tumor	t(1;2)(q22;p23)	*TPM3-ALK* fusion*
	t(2;19)(p23;p13)	*TPM4-ALK* fusion
	t(2;17)(p23;q23)	*CLTC-ALK* fusion*
	t(2;2)(p23;q13)	*RANBP2-ALK* fusion
	t(2;2)(p23;q35)	*ATIC-ALK* fusion*
	t(2;11)(p23;p15)	*CARS-ALK* fusion
	t(2;4)(p23;q21)	*SEC31L1-ALK* fusion
	t(2;12)(p23;p12)	*PPFIBP1-ALK* fusion
Low-grade fibromyxoid sarcoma	t(7;16)(q33;p11)	*FUS-CREB3L2* fusion
	t(11;16)(p11;p11)	*FUS-CREB3L1* fusion
Mesenchymal chondrosarcoma	t(8;8)(q13;q21)	*HEY1-NCOA2* fusion
Myoepithelial tumors of soft tissue	t(6;22)(p21;q12) t(19;22)(q13;q12) t(1;22)(q23;q12)	*EWSR1-POU5F1* fusion *EWSR1-ZNF444* fusion *EWSR1-PBX1* fusion
Myxoid/round cell liposarcoma	t(12;16)(q13;p11)	*FUS-DDIT3* fusion
	t(12;22)(q13;q12)	*EWSR1-DDIT3* fusion
Myxoinflammatory fibroblastic sarcoma/hemosiderotic fibrolipomatous tumor	t(1;10)(p22;q24) 3p11–12 (ring chromosomes)	*TGFBR3-MGEA5* fusion Amplification of *VGLL3*, *CHMP2B*
Nodular fasciitis	t(17;22)(p13;q13)	*MYH9-USP6* fusion
Sclerosing epithelioid fibrosarcoma (some cases)	t(7;16)(p22;q24)	*FUS-CREB3L2* fusion
Synovial sarcoma		
Biphasic	t(X;18)(p11;q11)	Predominantly *SS18-SSX1* fusion
Monophasic	t(X;18)(p11;q11)	*SS18-SSX1*, *SSX2*, or *SSX4* fusion
Tenosynovial giant cell tumor	t(1;2)(p13;q35)	*COL6A3-CSF1* fusion
Well-differentiated liposarcoma/dedifferentiated liposarcoma	Ring form of chromosome 12	Amplification of *MDM2*, *CDK4*, and others



Let's Begin



Adipocytic Tumors



Concepts in Fat

Normal fat cell

Brown fat

Fat necrosis, histiocytes

Lipoblast, signet ring

Lipoblast, multivacuolated



Not Lipoblasts

Fat necrosis
- Pitfall: histiocytes can stain with MDM2/CDK4

Lochkern fat cell
- Vacuolated nucleus b/c seen en face
- Benign

German:
Loch=hole
Kern=seed



Benign adipocytic tumors

Lipoma	Hibernoma	Angiolipoma	Lipoblastoma	Spindle cell / pleomorphic lipoma
Adults Superficial	Young adults	Young adults, Painful 2/3 Forearm 2/3 Multiple	Neonates Children (up to 3 yrs) Trunk, Extremities	Middle aged men Poster neck, shoulder, back
Conventional Intramuscular Variants: Myo, Fibro, Myxoid, Chondroid (t11;16)	Brown fat in varying degrees	Capillary-like vessels Fibrin thrombi	Lobulated / Septated Myxoid matrix Lipoblasts to mature fat	Histologic range: Ropey collagen Blunt spindle cells, palisading Hyperchromatic / Floret cells
Can have extensive fat necrosis (histiocytes can be MDM2/CDK4+)	Radiographically can look heterogenous, like ALT	May be familial (AD inheritance)	Can mature in older childhood Lipoblastomatosis if multiple	CD34 + Rb lost (mammary type myofibroblastoma) MDM2, CDK4 negative
12q* (HMGA2) or 6p transloc. 13q deletions	11q* or 10q translocations	None ? May be a hamartoma	8q (PLAG) translocation	16q or 13q loss



Spindle Cell – Pleomorphic Lipoma

- Is a histologic spectrum of genetically identical tumors
  - Ropey collagen
  - CD34+
  - Lack of MDM2/CDK4 expression
  - Loss of Rb (13q)
- Pleomorphic lipoma has floret giant cells
  - DDx Atypical lipomatous tumor
- Usually subQ > dermal



Mammary Type Myofibroblastoma

- Looks like myofibroblastoma of breast
- Spectrum with spindle cell lipoma & cellular angiofibroma
- Inguinal / paratesticular / vulvovaginal areas "milk line"
- Older men / postmenopausal women
- Histology:
  - Well circumscribed
  - Bland spindle cells in fascicles
  - Hyalinized collagen
  - Mast cells
  - Fat (rare fat-free variant)
  - Desmin, CD34+
- 13q, 16q partial monosomy = loss of Rb (like SCL)

Cellular angiofibroma:
Perineal, M=F
CD34, SMA, ER, PR



Malignant adipocytic tumors: 3 groups

Atypical Lipomatous Tumor
Well-Differentiated Liposarc
↓
De-Differentiated Liposarc

Myxoid Liposarc
↓
Round Cell Liposarc

Pleomorphic Liposarc



Atypical lipomatous tumor / Well-differentiated liposarcoma

Tumor	Clinical	Histology	Molecular	Prognosis

ALT
WD-LPS
DD-LPS
	• Adults
	• Extremity (Deep) (ALT) vs
	• Retroperitoneum / Spermatic cord (LPS)
		• Well-Diff: Hyperchromatic, enlarged stromal cells in fat/septae
		  • Lipoblasts NOT USUAL
		  • Variants: Adipocytic (Lipoma-like), Sclerosing, Inflammatory
		• De-Diff: high grade spindle sarcoma, usually nonlipogenic, may have heterologous differentiation (leiomyosarc, rhabdomyosarc, neural-like with whorls/osteoid, MPNST-like)
		  • Other Variants: Myxoid LPS-like, Pleomorphic LPS-like
		  • Some say need 5 mits / 10 hpf to make De-Diff in hypercellular area
			• 12q amplification (giant marker/ring chromosomes)
			  • FISH: MDM2 amplified
			  • IHC: MDM2, CDK4, p16+ (at least 2/3)
			• De-Diff: coamplification of 1p/6q (JUN, ASK1)
				• WD extremities: may recur with incomplete excision, rarely dedifferentiate
				• WD retroper: often recur and 20% dedifferentiate
				• DD: <2 cm indolent
				• DD: recurrence in 40%, metastasis in 20%
				• DD: less aggressive than other high grade sarcomas



Myxoid – Round Cell Liposarcoma

Tumor	Clinical	Histology	Molecular	Prognosis
Myxoid LPS Round Cell LPS	• Young adults • Extremity • NOT retroperitoneum (*metastasis*)	• Myxoid: myxoid stroma / mucin pools, curvilinear vessels, bland spindle cells and lipoblasts, more cellular at periphery, rare mitoses • Round cell: back to back / overlapping cells with high N:C ratio, nucleoli, mitoses	• t(12;16) DDIT3(CHOP)-FUS 95% • t(12;22) DDIT3-EWSR1 5%	• Myxoid: <10% mets to other soft tissue sites/bone • Round cell: if >5% worse prognosis



Pleomorphic Liposarcoma

Tumor	Clinical	Histology	Molecular	Prognosis
Pleomorphic LPS	Elderly Extremity, Deep	High grade spindle sarcoma with Pleomorphic Lipoblasts and Myxofibrosarcoma like areas	Complex karyotype	Aggressive: 30-50% recur/metastasize



ALT – WDLPS – DDLPS	Myxoid LPS – Round Cell LPS	Pleomorphic LPS
• Adults • Deep Extremity (ALT) • Retroperitoneum / Spermatic cord (LPS)	• Young adults • Extremity • Other sites – retroperitoneum, etc = metastasis !!!!	• Elderly • Deep Extremity

• Well-Diff: Hyperchromatic, angulated nuclei
  • Lipoblasts NOT USUAL
  • Variants: Adipocytic, Sclerosing, Inflammatory
• De-Diff: high grade spindle sarcoma (5/10 mits)
• Other Variants: Myxoid LPS-like, Pleomorphic LPS-like
	• Myxoid: myxoid stroma / mucin pools, curvilinear vessels, bland spindle cells and lipoblasts, more cellular at periphery, rare mitoses
	• Round cell: back to back / overlapping cells with high N:C ratio, nucleoli, mitoses
		• High grade spindle sarcoma with
		• Pleomorphic Lipoblasts and
		• Myxofibrosarcoma like areas

• 12q amplification (giant marker/ring chromosomes)
  • FISH: MDM2 amplified
  • IHC: MDM2, CDK4, p16+ (need at least 2 of 3)
	• t(12;16) DDIT3(CHOP)-FUS 95%
	• t(12;22) DDIT3-EWSR1 5%
		• Complex karyotype

• ALT extremities: recur • WD retroper: recur and 20% dedifferentiate • DD: recurrence in 40%, metastasis in 20% • DD: less aggressive than other high grade sarcomas	• Myxoid: <10% mets to other soft tissue sites/bone • Round cell: if >5% worse prognosis	• Aggressive: 30-50% recur/metastasize

Myogenic Tumors

	Smooth Muscle	Skeletal Muscle	Perivascular Myoid
Benign	• Leiomyoma	• Rhabdomyoma variants	• Glomus and variants • Myopericytoma and variants • Angioleiomyoma
Malignant	• Leiomyosarcoma	• Rhabdomyosarcoma variants	• Malignant glomus
IHC Staining	Desmin, SMA, Calponin, Caldesmon	Desmin, Myogenin/MyoD1	SMA, Caldesmon (Desmin in angioleiomyoma)

Leiomyoma

- Can be superficial (cutaneous) or deep
- Deep leiomyomas
  - Retroperitoneum: Mostly adult female
  - Periphery: M=F
- Histology
  - Intersecting fascicles of eosinophilic spindle cells
  - Blunt-end / cigar-shaped nuclei
  - Atypia / mitoses MINIMAL

  - Retroperitoneal LM & LMS can be ER/PR positive even if not gyne tract (M & F) (Springerplus. 2013; 2: 536.)



Leiomyosarcoma

- Most common in retroperitoneum and/or arising from large vessels
- Location
  - Retroperitoneum: Mostly adult female*
  - Cutaneous: Mostly adult male
- Histology
  - Like leiomyoma, however, cytologic atypia may be mild or marked & mitoses may be few or many
    - Threshold depends on site: skin, retroperitoneum, uterus
  - May aberrantly express keratin, EMA, CD34, S100
  - A diagnosis of LMS should not be made in a high grade sarcoma based on staining for myogenic markers alone: Need morphologic evidence of smooth muscle phenotype!
- Prognosis
  - Location & Size
  - Most common sarcoma to metastasize to skin
- EBV-associated smooth muscle tumors in immunocompromised patients:
  - Parenchymal organs
  - More young adults / children
  - +/- lymphoid infiltrate
  - Better behavior



LMS criteria by site

Soft Tissue / Bone / Retroperitoneum	Skin	Uterus – much more strict
Cytologic atypia can be mild / focal, but is often marked	Cytologic atypia needed	If moderate – severe cytologic atypia:
Mitoses may be few	2+ mitoses / 10 HPF	• If coagulative necrosis present -> sarcoma
Necrosis frequent		• If coagulative necrosis absent -> require diffuse atypia and 10+ mitoses
		If no – mild cytologic atypia:
		• Require coagulative necrosis and mitoses 10+

Rhabdomyoma

Cardiac (tuberous sclerosis)	Adult (extracardiac)	Fetal (Gorlin-PTCH1)	Genital
Heart	Head and neck	Head and neck (postauricular)	RARE
Perinatal	Adult M>F	Child (age 2) M>F	Adult F>M
Glycogenated cytoplasm +/- Spider cells	Large cells with granular eosinophilic cytoplasm +/- Striations / inclusions	Myotubes in myxoid stroma +/- Rhabdomyoblasts	Rhabdomyoblasts in loose stroma



Embryonal Rhabdomyosarcoma

Embryonal
• Most common subtype • Children <5 • Head and neck / GU • "Botryoid" protrudes from epithelial-lined organs
• Spindle-stellate cells in alternating dark and light areas (myxoid stroma) • +/- Rhabdomyoblasts (tadpole, strap with striations) • +/- Round cells • Cambium layer in botryoid (subepithelial condensation)
• Nuclear MyoD1 > myogenin • (diffuse strong myogenin in primitive sarcoma = consider alveolar RMS!)
• Better outcomes < age 10 • Better outcomes than alveolar







http://www.auanet.org/images/education/pathology/paratesticuler-tumors/rhabdomyosarcoma-figureD_Big.jpg

Alveolar Rhabdomyosarcoma

Alveolar

- Adolescents/young adults
- Extremities

- Primitive, round cells
- May have prominent nucleoli
- Little cytoplasm to eccentric pink cytoplasm
- Fibrovascular septae separating cells into “alveoli”
- May be discohesive or solid

- Strong diffuse myogenin
- PAX3 (or PAX7) - FOXO1 fusion (chr 2 (or 1) - 13)

- More aggressive than embryonal (except if fusion negative)





Rhabo (SRBCT) Differential Diagnosis

Rhabdomyosarcoma	Desmoplastic small rd cell (DSRCT)	Ewing-PNET
Children	USUALLY intra-abdominal Teenage Males	Bone > Soft tissue Children > Adults
Desmin + Myogenin / MyoD1 + Also: Carcinoma Melanoma Lymphoma	Multiphenotypic differentiation: • Epithelial: Keratin/EMA + • Myogenic: Desmin + dot like • Neural: NSE + • WT-1: carboxy term nuc +, amino term perinuc + • CD99 cytoplasmic +/-	CD99 membranous + FLI1 nuclear + OR ERG nuclear +
PAX3-FOXO1 t(2;13) if alveolar * Poor prognosis if alveolar	EWSR1-WT1 t(11;22) * Poor prognosis	EWSR1-FLI1 t(11;22)(q24;q12) EWSR1-ERG t(11;22)(q12;q12)



Nests of small cells in desmoplastic stroma +/-necrosis, cysts, rosettes

Ewing Like Sarcomas



BCOR-CCNB3 tumors	CIC-DUX4 tumors
• most occur in bone	• 2/3 of EWSR1-negative tumors • most occur in soft tissue
• heterogeneous small round and spindle cells • stroma may be myxoid	• sheets of small round cells +/- spindle cells • stroma may be myxoid • less monotonous than EWS • may have more cytoplasm
• usually patchy CD99 positive • stain for BCOR and CCNB3	• usually patchy CD99 positive • stain for WT1, ETV6, and CIC

Small Round Blue Cell Tumor Differential Diagnosis Summary

Tumor	CD99	FLI-1	EMA	CD45	Desmin	Myogenin	WT-1	TLE-1
Ewing/PNET	+	+	+/-	-	-	-	-	-
Lymphoma	+	+	-	+	-	-	-	-
Rhabdo	+	-	-	-	+	+	-	-
DSRCT	+	-	+	-	+	-	+ C-term nuclear + N-term cyto	-
Synovial Sarc	+/-	-	+	-	-	-	-	+
Osteosarc	-	-	-	-	-	-	-	-

Pleomorphic Rhabdomyosarcoma



Pleomorphic

- Adult M>F
- Deep soft tissue of extremities

- Spindled to epithelioid cells
- Large cells with abundant pink cytoplasm and multinucleation
- Rare striations

- Myogenin, MyoD1+
- May show nuclear MDM2 without amplification

- Very aggressive, worse with increasing age



Spindle Cell Rhabdomyosarcoma

Spindle cell / sclerosing

- M>>F
- Children: Paratesticular
- Adults: Deep soft tissue of head and neck

- Spindled cells with elongated nuclei, pale eosinophilic cytoplasm
- Rare rhabdomyoblasts +/- striations
- +/- Stromal hyalinization / pseudovascular = sclerosing variant

- Desmin +/- SMA
- Myogenin in rhabdomyoblasts +/- spindle cells
- Sclerosing: MyoD1>myogenin

- Good prognosis in childhood
- Worse prognosis in adulthood







Perivascular Myoid Tumors

Glomus	Myofibroma	Myopericytoma	Angioleiomyoma
• Young adults • Distal extremities, Superficial • (Subungual F>M) • Clinically red-blue & painful	• Child or Male Adult • Superficial, Extremities, Head/Neck, Trunk • Infantile Myofibromatosis: AD	• Adult • Superficial, Distal extremities	• Adults, SubQ • Lower extremity F>M • Upper extremity, Head/Neck M>F • Painful
• Uniform round cells with central round nucleus • Amphophilic/eosinophilic cytoplasm • Distinct cell borders (each cell surrounded by basal lamina) • SMA, caldesmon +	• Biphasic: immature plump spindle cells & mature eosinophilic spindle cells in whorls • Associated with HPC-like vessels, myxoid stroma • SMA + (caldesmon rare)	• Spectrum with Myofibroma / Angioleiomyoma • Perivascular concentric growth of spindle cells with eosinophilic cytoplasm • SMA, caldesmon + (desmin rare)	• Well-differentiated smooth muscle cells surrounding vascular channels: slit-like, venous with perivascular whorling, cavernous • SMA, desmin + in most
• Malignant variant: nuclear atypia + mitoses • GLMN (glomulin) gene in familial cases • Associated with NF1	• Usually cured by excision • Worse outcome with visceral involvement in myofibromatosis	• Usually cured by excision • Rarely malignant	• Usually cured by excision



Glomus Variants

- Solid glomus
- Glomangioma:
  - + cavernous vessels
- Glomangiomyoma:
  - + spindled smooth muscle like cells
- Glomangiopericytoma:
  - + HPC-like vessels (sinus HPC)
- Symplastic:
  - + random nuclear atypia



Vascular Tumors

Benign	Intermediate	Malignant
Epithelioid Hemangioma	Epithelioid Hemangioendothelioma	Epithelioid Angiosarcoma
Juvenile Hemangioma	Pseudomyogenic Hemangioendothelioma	Angiosarcoma Variants
		Kaposi Sarcoma



Infantile Hemangioma

- Female children
- Arise in postnatal few weeks, rapidly enlarge over months, peak at 6-12 months, regress over years
- Flat -> elevated red lesion on skin
- Histology:
  - Multinodular, feeder vessel
  - Proliferative phase with plump endothelium
  - Maturation phase with flattened endothelium, prominent basement membrane
  - GLUT1 positive (negative in other hemangiomas)



Epithelioid Hemangioma

(Angiolymphoid hyperplasia with eosinophilia)

- Many ages, mostly adult female
- Head / Distal extremities / Bone
- Neoplasm vs Reactive?
- Can be locally aggressive
- Histology:
  - Nodular proliferation
  - Capillary sized vessels lined by plump endothelial cells
  - Can be solid or spindled
  - Can have large grooved nuclei
  - Eosinophils / Lymphocytes
- CD31 > CD34, ERG
- 1/3 locally recur
- Unlike EHE: vasoformative, no/rare intracytoplasmic lumina



Epithelioid Hemangio-endothelioma (EHE)

- Adult (30s-40s) females
- Extremities, ANY site (lung, liver, bone)
- Multifocal lesions are likely metastases from a single primary
- Histology:
  - May be angiocentric
  - Chords of epithelioid endothelial cells in myxoid stroma
  - Eosinophilic cytoplasm with VACUOLES (intracytoplasmic lumina +/- RBCs)
  - Not architecturally vasoformative
  - CD34, CD31, FLI1, ERG, keratin in up to half
  - WWTR1-CAMTA1 t(1;3)
- Up to 30% metastasize, 15% die
  - Tumors >3 cm and >3 mitoses/50 HPFs have a 5-year disease-specific survival of 59% (versus 100%) (Modern Pathology. 2014. 27, S30–S38).

Differential Diagnosis

Epi. Hemangioma	Epi. Hemangioendothelioma	Epi. Angiosarcoma
Small vessels and solid nodules +/- Eos / inflammatory cells	Chords and single cells	Sheets of epithelioid cells Cytologically atypical, mitotically active
Vasoformative	NOT vasoformative Intracytoplasmic lumina	Vasoformative
CD31+, CD34+, FLI1+, ERG+, Keratin+/-		



Pseudomyogenic Hemangioendothelioma

- Young adult males, lower limbs, multifocal
- Intramuscular > bony
- Histology:
  - Sheets/fascicles of loose, plump spindle cells with bright eosinophilic cytoplasm
  - Mild nuclear atypia
  - +/- myxoid stroma, neutrophils
  - Keratin, ERG
  - CD31 50%, SMA 30%
  - CD34, Desmin neg
- Genetics:
  - t(7;19) SERPINE1-FOSB



Angiosarcoma

1. Primary cutaneous
   - Sun-exposed scalp, older patients
   - +/- heavy lymphoid infiltrate
   - Some have UV signature
2. Primary soft tissue
   - Deep lower extremities
   - Vasoformative to solid, spindled to epithelioid
   - Epithelioid: Large nuclei, prominent nucleo
3. Primary breast
   - Deep breast, women 20s-30s
   - Well-differentiated vasoformative, flattened bland endothelial cells; but diffusely infiltrates
4. Radiation-related & Lymphedema-related (LE: Stewart–Treves syndrome)
   - Cutaneous (breast or edematous regions)
   - Vasoformative to solid, infiltrative
   - MYC amplification (8q24) in >90%
   - FLT4 co-amplification in 25%
5. Prognosis: Older age, retroperitoneal location, larger size, higher Ki67


Vasoformative
Solid
Post radiation atypical vascular lesion
Post radiation angiosarcoma
d
e
Primary breast angiosarcoma
h
i

Modern Pathology. 2014; 27, S30-S36

Kaposi Sarcoma



The American Journal of Dermatopathology
Volume 5 Number 1
February 1983

LETTERS TO THE EDITOR

Kaposi: Let's Pronounce It Correctly

To the Editor:

KAPOSI (1837-1902).

. . ."(1) Kaposi is one proper name which is almost universally mispronounced in America. In the annoted list of eponyms which Dr. Leider provides, the correct pronunciation is given as Kă-pō-sē, shē. However, in Hungarian the "a" is pronounced as a short "o," "very like English 'o' in 'hot,' but the mouth is opened more widely."(2) In addition, Dr. Karl Holubar has pointed out: "As regards pronounciation, the river Kapos, the town Kaposvár and the name Kaposi all carry the accent on the first syllable and not as frequently heard in Anglo-Saxon countries on the second."(3) Hence, the correct pronunciation is Kŏ-pō-shē.

D. Friday King, M.D.
Laura A. C. King, M.D.
Long Beach, California

Kaposi Sarcoma

- Forms:
  - Classic Indolent: Elderly Mediterranean men
  - Endemic African: Adults/children WITHOUT HIV
  - Iatrogenic: Immunosuppressed, usually solid organ transplant
  - AIDS-associated: Immunosuppressed, HIV-1, often men
    - * most aggressive
- KS-associated herpesvirus (HHV8) in all forms
- Mostly skin but also viscera
- Histology:
  - Vascular spaces dissecting collagen
  - Spindle cell proliferation
  - Extravasated RBCs/hemosiderin
  - +/- inflammatory infiltrate
  - +/- hyaline globules (arrow)
  - + for vascular markers and HHV8



Nerve Sheath Tumors

- Components of a Nerve:



Perineurium
- EMA, GLUT1

Schwann cells (myelin)
- S100

Axons
- Neurofilament

Supporting cells
- Vessels
- Fibroblasts

- Can develop a tumor of any of the components:
  - Perineurioma
    - Perineurial cells
  - Schwannoma
    - Schwann cells
  - Neurofibroma
    - ALL (perineurial, Schwann, axons, fibroblasts, mast cells)
  - Hybrid
  - Malignant Peripheral Nerve Sheath Tumor (MPNST)



Nerve Sheath Tumors

Neurofibroma

Usually monotonously cellular
Wavy nuclei in collagenous-myxoid background (shredded carrot collagen)

Pseudo Meissnerian corpuscle

Tactoid body

Schwannoma

Usually encapsulated (perineurium)
Variably cellular
Antoni A (hypercellular) & Antoni B (hypocellular)



Schwannoma

- Types:
  - Conventional
  - Ancient: degenerative smudgy nuclear atypia / extensive hyalinization
  - Cellular: dense, no Verocay bodies
  - Plexiform: multinodular (loose a/w NF2)
  - Microcystic/reticular: preferentially in GI
  - Multiple a/w:
    - NF2: < age 30, bilateral on vestibular nerve ("acoustic neuroma")
    - Schwannomatosis (SMARCB1 gene mutation): adulthood, NOT on vestibular nerve
- Usually cured if excised
- NOT a precursor to spindle cell MPNST (rarely to epithelioid MPNST)

Notes:
- S100 usually diffusely strongly positive
- Collagen IV/Laminin surrounds individual cells
- May stain with AE1/AE3 as AE3 cross reacts with GFAP
- Entrapped NF+ axons may be present at periphery = NOT a sign the tumor is NF



(Malignant) Melanotic Schwannian Tumor

- Adult females, paraspinal
- Schwannian differentiation:
  - S100, SOX10+
- Melanocytic differentiation:
  - HMB45, MelanA
- Half of tumors have psammoma bodies
- Half of these patients have Carney complex
- May METASTASIZE

Malignant Melanotic Schwannian Tumor
A Clinicopathologic, Immunohistochemical, and Gene Expression Profiling Study of 40 Cases, With a Proposal for the Reclassification of "Melanotic Schwannoma"
Jorge Torres-Mora, MD, Sarah Dry, MD,† Xinmin Li, PhD,† Scott Binder, MD,† Mitual Amin, MD,‡ and Andrew L. Folpe, MD**

No longer "psammomatous melanotic schwannoma"

Am J Surg Pathol 2014;38:94–105.



Neurofibroma

- Most common peripheral nerve sheath tumor
  - Most sporadic, solitary, skin
  - Multiple in NF1 (17q), 10% lifetime risk of progression to MPNST
- 5 macroscopic types:
  - Localized cutaneous = benign
  - Diffuse cutaneous
  - Localized intraneural = MPNST precursor
  - Plexiform intraneural = MPNST precursor
  - Massive diffuse soft tissue plexiform = benign but may be adjacent to intraneural

Atypical neurofibroma:
- Hypercellularity
- Hyperchromatic, smudgy nuclei
- Mitoses
- Atypical NF vs LG MPNST….?
- Precursor to MPNST

http://www.webpathology.com/slides-13/slides/SoftTissue_Neurofibroma67_Atypia_NF1.jpg



Plexiform NF

- Bag of worms
- Pathognomonic of NF1

http://www.sarcomaimages.com/images/db/neurofibroma/neurofibroma-2-gross-sarcoma-images.jpg



Perineurioma

- Benign soft tissue, intraneural, or mucosal
- F>M, wide age range, lower limbs
  - Sclerosing variant on hands
- Histology:
  - Storiform growth pattern
  - Long fascicles
  - Collagenous-myxoid stroma
  - EMA, Claudin1, GLUT1, CD34
  - S100 negative
- NF2 mutation/deletion in some
- "Malignant Perineurioma" (perineurial MPNST) is rare and may behave better than classic MPNST



Hybrid BPNST

- Most commonly sporadic Schwannoma / Perineurioma
  - Storiform – fasciculated growth pattern like perineurioma
  - EMA+ in perineurial cells
  - S100+ in Schwann cells
- Less common NF1-associated or sporadic Neurofibroma / Schwannoma
  - Schwann nodules in a typical NF

Schwannoma / Perineurioma

EMA

S100



MPNST

- Sarcoma with Schwann cell differentiation
  - De-novo
  - From a benign NST
  - 50% in NF1 patient from plexiform NF
  - 10% post radiation
- Fusiform mass involving major nerve
- Histology:
  - Conventional
    - Spindled fascicles with marbling (dark-light)
    - HPC-like blood vessels
    - Heterologous: bone, cartilage, skeletal muscle (Triton), glands, melanin
    - Only 50% S100+/SOX10+, usually focal
    - 25% GFAP+
  - Epithelioid
    - Most common type to arise from Schwannoma
    - NOT a/w NF1
    - Most are S100+, strong
    - May be keratin+, INI1 lost 50%
- Poor prognosis
  - trunk, size, grade, Triton, NF1

Conventional

Triton

Epithelioid

Tumors of Fibroblastic / Myofibroblastic / Uncertain Lineage



Let's start with paucicellular spindle cell tumors

We'll talk about:	We've already seen:
Fasciitis-es	BPNSTs
Fibromatoses	Leiomyoma
Fibromas	
Myxoid sarcomas	



Nodular Fasciitis

- Wide age range
  - Young adults*
  - Extrem/Trunk/H&N
- +/- Trauma
- Rapid growth 2-3 months
- Usu < 5 cm
- Histology:
  - Circumscribed OR Infiltrative
  - "Tissue culture" myofibroblasts
    - Wispy cytoplasm
    - Oval nuclei with small nucleoli
  - Myxoid to collagenous stroma
  - Mitotically active
  - SMA+
  - MYH9-USP6 (22q-17p) gene rearrangement in most = self limited NEOPLASM not reactive
- May regress, should NOT recur after excision
  - RARE metastasizing / malignant forms that are histologically identical

In what other tumors is USP6 rearranged?

- Aneurysmal Bone Cyst
- Cellular Fibroma of Tendon Sheath
- Myositis Ossificans



Microcystic spaces, myxoid

Keloid-like collagen
Extravasated RBCs

NODULAR FASCIITIS

Smooth muscle actin
USP6 rearranged 90%

Plump, whorled fibroblasts (younger)

Thin spindle-stellate fibroblasts



Other Fasciitis

- Intravascular
  - Age <30
  - SubQ, small-medium veins
  - May have increased osteoclast like giant cells
- Cranial
  - Age <2, M
  - External skull, may infiltrate skull / meninges
- Proliferative
  - Older adults
  - Upper extremity
  - SubQ (if intramuscular = proliferative myositis)
  - Nod-Fasc like background with large ganglion-like cells
- Ischemic
  - Elderly, M, often immobile, limb girdles
  - Zonal:
    - Central hypocellular fibrin/necrosis
    - Peripheral granulation tissue like with ganglion-like cells

Proliferative



Fibromatoses

Deep (Desmoid)	Superficial	Colli	Juvenile Hyaline	Inclusion Body	Lipo
Kids: extra-abdominal, M=F Adults: abdominal wall, F (pregnant/peripartum) Older: anywhere, any gender Gardner syndrome (APC mutation)	Adults >30 Palmar (Dupuytren contracture) Plantar (in kids too) Penile (Peyronie disease)	"Torticollis" Distal SCM of infants M>F ? Scar-like reaction to intrauterine injury	Rare, infants AR = Consanguinity Skin of face/neck Gums Periarticular Bones	"Infantile digital" Neonates Dorsal 2nd, 3rd, 4th digits	VERY RARE Hand/Foot Newborn-Teen
Infiltrative Long fascicles of bland spindle cells in collagenous stroma with prominent vessels Oval nuclei with small nucleoli	Younger: cellular plump fasciculated spindle cells Older: increased collagen Plantar: multinucleated giant cells	Younger: cellular plump fasciculated spindle cells Older: increased collagen Entrapped muscle	Plump fibroblasts in homogenous non-fibrillar hyaline material	Whorls, fascicles of eosiniphilic spindle cells Perpendicular to dermis Surround skin adnexa Eosinophilic paranuclear inclusion, indents nucleus	50% adipose 50% fasciculated fibroblasts
SMA+ Beta-catenin nuclear+ 75%	SMA+/- Beta-catenin 50% nuclear+	SMA+ Beta-catenin –	SMA – PAS+, Diastase resistant material	SMA+ Inclusions stain red on Trichrome	SMA+, CD34+ Beta-catenin –
Gardner: APC mutation Sporadic: B-cat mutation	Nonspecific	None	ANTXR2 mutation (4q)	None	Nonspecific
Local recurrence common, related to	Local recurrence related to margins	Physical therapy to normalize posture	Local recurrence common	Local recurrence related to margins	Local recurrence common



Desmoid Fibromatosis

Abdominal wall
Intra-abdominal
A/W Gardner Syndrome
Long, pale fascicles
Vesicular nuclei, nucleoli
SMA +
Beta-catenin nuclear + 75%



Fibromas

Tendon Sheath	Nuchal-type	Gardner	Elasto-	Calcifying aponeurotic
Adults Fingers	Adult M Posterior neck & Extranuchal Up to 50% have DIABETES	Children, Trunk, Head/Neck 80% occur in syndrome (FAP, familial desmoid)	Elderly F Between scapula and chest wall	Children/Adolescents Palms
Circumscribed Bland spindle cells in collagenous background, more cellular at periphery Slit like vessels	Ill-defined Intersecting, lobulated collagen bundles Entrap fat, skin adnexa, skeletal muscle Delicate elastic network	Ill-defined Hypocellular coarse collagen sheets Entraps fat, nerves, muscle	Ill-defined Fibroblasts and collagen Abnormal elastic fibers: thick, fragmented, beads on string	Infiltrative cellular spindle cell component Nodules of epithelioid cells with calcification/chondroid stroma Osteoclast-like giant cells
SMA+ focal USP6 in cellular variant	CD34+, SMA --	CD34+, SMA – Beta-catenin nuclear +	Fibroblasts: CD34+, SMA -- Elastin: elastin stain	SMA+ EMA --, S100 --
May recur if incompletely excised	May recur	SCREEN FOR FAP	Cured by excision	May recur years later



Hypocellular Myxoid Sarcomas

Low Grade Fibromyxoid Sarcoma (LGFMS, Evans tumor, Hyalinizing spindle cell tumor with giant rosettes)	Myxofibrosarcoma, low grade
Is by definition LOW GRADE Has a high grade counterpart – sclerosing epithelioid fibrosarcoma (SEF) – which is morphologically distinct	Myxofibrosarcoma can be OF ANY GRADE depending on histologic features Very common in elderly males
Young adults Extremities / Trunk	Elderly Extremities
Well circumscribed Discrete myxoid / collagenous zones Variably cellular Minimal nuclear atypia 30% have collagen rosettes (peripheral epithelioid cells)	Highly infiltrative Multiple nodules of atypical spindle cells within myxoid stroma Curvilinear blood vessels with perivascular atypical tumor cells
EMA+, MUC4+ t(7;16) FUS-CREB3L2 * t(11;16) FUS-CREB3L1	May be SMA+ (myofibroblastic) No specific genetics, complex karyotype
Low recurrence rates Rare late metastases May be seen in association with SEF	Low grade tumors recur and increase in grade High grade tumors metastasize



Sclerosing Epithelioid Fibrosarcoma

- On a spectrum with LGFMS?
- Older adults, lower extremity, bone
- Histology:
  - Epithelioid cells in strands: "single file"
  - Separated by collagenous stroma (resembles osteoid)
- EMA+ 50%, MUC4+ 70%
- Genetics:
  - Hybrid LGFMS-SEF:
    - t(7;16) FUS-CREB3L2
    - t(11;16) FUS-CREB3L1
  - Pure SEF:
    - EWSR1-CREB3L1/2



Myxoid-Spindled Differential Diagnosis Summary

Tumor	EMA	S100	Muc4	Other Diagnostic Tools
Myxoma	-	-	-	+/- CD34
Myxoid neurofibroma	-	+	-	
Myxoid perineurioma	+	-	-	
Myxoid liposarcoma	-	-	-	Lipoblasts CHOP-FUS
Fibromatosis	-	-	-	SMA Beta-catenin
Myxofibrosarcoma LG (MFS, LG)	-	-	-	Older age, infiltrative, atypia
LG Fibromyxoid sarcoma (LGFMS)	+	-	+	Younger age, circumscribed, lack of atypia

Cellular Sarcomas



We'll talk about:	We've already seen:
Synovial Sarcoma	MPNST
SFT	Rhabdomyosarcoma
IMT	Leiomyosarcoma
Fibrosarcoma	Dediff liposarcoma
UPS	SEF

Synovial Sarcoma

- Any age, mostly young adults
- Deep extremities
  - Juxta-articular prompted "synovial" misnomer
- Monophasic
  - Hypercellular spindle cells in vague fascicles, storiform
  - Uniform, bland nuclei
  - HPC-like vessels
  - May have collagenous/myxoid stroma, calcification/ossification
  - t(X;18) SYT-SSX2
- Biphasic
  - Spindle and epithelioid cells
  - t(X;18) SYT-SSX1
- Poorly differentiated
  - Round/spindle cells with severe nuclear atypia, increased mitoses, prominent nucleoli, necrosis
- IHC:
  - EMA > keratin
  - TLE1 nuclear +
  - CD99+, S100+/-
- Prognosis = stage, size, grade
  - May metastasize to lung, bone
  - Worse in adults



Solitary Fibrous Tumor

- Occurs in pleura, soft tissue, head/neck, meninges
  - "HPC" is obsolete except in head/neck
- Adults, M=F
  - Lipomatous SFT = Males
- Histology:
  - Usu well-circumscribed
  - "Patternless pattern": Variably cellular, bland oval spindle cells in collagen
  - HPC-like vessels
  - CD34+, STAT6+ (inv12 NAB2-STAT6)
  - Variable EMA, SMA, S100, keratin
- ALL should be considered sarcomas b/c behavior not always predicted by histology…
  - Deep, large (>10 cm), + margins, high grade histology
  - "Malignant SFTs" are hypercellular, necrotic, increased mitoses (>4/10)
  - "Dediff SFTs" show abrupt transition to high grade sarcoma
- IGF2 secretion = Hypoglycemia



Inflammatory Myofibroblastic Tumor

- Plasma cell granuloma, inflammatory pseudotumor
- Children/young adults > adults, F>M
  - Throughout body
- Histology: Spindled myofibroblasts in 3 patterns:
  1. Hypocellular myxoid
  2. Cellular fasciculated + ganglion-like cells
  3. Scar-like collagenous
  - All with variable infiltrate of lymphocytes, plasma cells, eosinophils
  - Rare obvious sarcomatous transformation
- SMA, MSA, desmin +/-
- ALK + cytoplasmic in up to 60% = ALK rearrangement
  - Nuclear membrane = RANBP2 fusion partner
  - Granular cytoplasmic = CLTC fusion partner
- Fusion pos = Children
- Fusion neg = Adults = higher likelihood of metastasis



Fibrosarcoma

Infantile	Adult
First year of life Distal extremities, trunk, head/neck Rapidly enlarging	Middle age-older adults Extremities, trunk, head/neck Now very rare * except if transformed from DFSP
Infiltrative Herringbone pattern: Highly cellular intersecting fascicles of spindle to round cells	
ETV6-NTRK3 t(12;15)	Nonspecific
Good prognosis, rare metastases	May recur, metastasize



What other tumors have ETV6-NTRK3 rearrangement?

- Cellular congenital mesoblastic nephroma
- Secretory carcinoma (mammary, salivary)
- Leukemia (AML)

Undifferentiated Sarcoma

- Formerly, undifferentiated pleomorphic sarcoma, malignant fibrous histiocytoma
  - Most high grade sarcomas can be shown to demonstrate a specific lineage of differentiation
  - The rest are undifferentiated
- Adults, somatic soft tissue
- Histology:
  - Pleomorphic
  - Spindle cell
  - Round cell
  - Epithelioid
  - Giant cell rich
- Diagnosis of exclusion
- >80% 5-year met.-free survival



Cellular-Spindled Differential Diagnosis Summary

Tumor	Keratins/EMA	S100	Myogenic	Other Diagnostic Tools
Carcinoma	+	-	-	p63/p40 for SCC
Melanoma	-	+	-	SOX10, HMB45, MelanA (~MiTF)
Leiomyosarcoma	-	-	+	
MPNST	-	~50% spindled ~100% epithelioid	Depends (Triton)	
Synovial sarcoma	+	-	-	TLE-1 (SYT-SSX)
DDLPS	-	-	Depends (heterologous)	MDM2, CDK4
DFSP, fibrosarcomatous	-	-	-	CD34, COL1A1-PDGFB
SFT, malignant	-	-	-	CD34, STAT6 (NAB2-STAT6)
UPS	-	-	Depends (myogenic diff)	

Epithelioid Tumors

We'll talk about:	We've already seen:
Epithelioid Sarcoma	Epi Angiosarc
ASPS	Epi MNPST
CCS	
MERT	
PEComa	
Myoepithelioma	
Chordoma	
EMC	

Epithelioid Sarcoma

* Metastasizes to lymph nodes!

Distal (Classic) Type	Proximal Type
More common Young adults (20s/30s), distal upper > lower extremity Nonhealing skin ulcers	Less common Older adults (up to 65), deep soft tissue of trunk/pelvis
Dermal/SubQ nodules of epithelioid-spindled cells with central necrosis / amorphous material Polygonal cells with eosinophilic cytoplasm small nucleoli, few mitoses	Nodules, sheets of large pleomorphic cells with large oval nuclei, prominent nucleoli +/- rhabdoid appearing cells, myxoid stroma
Keratin +, EMA + (differential diagnosis: carcinoma, other keratin-expressing sarcomas (epithelioid angiosarc, epithelioid MPNST, myoepithelioma, chordoma) CD34 + 50%, ERG + in some INI1 (SMARCB1) lost	
Up to 50% metastasize (lung, nodes); Proximal worse prognosis than distal (large size, early mets)	



Differential Diagnosis

Calcifying Aponeurotic Fibroma	Granuloma Annulare	Rheumatoid Nodule	Epithelioid Sarcoma (Distal)
Fibrohistiocytic cells at periphery: CD68/CD163+, SMA+			Keratin/EMA+ INI1 lost ~100%
Calcific / Chondroid core	Necrotic collagen core	Fibrin / Necrotic collagen core	Hyalinized core
Hands/feet Kids	Dorsal hands/legs Superficial: dermis	Periarticular Deep: subcutis	Distal digit Deep derm / subcutis



http://www.pathologyoutlines.com/images/softtissue/02_35L.jpg

a/w HCV, autoimmune thyroiditis, secondary hyperparathyroidism, MDS, metastatic carcinoma

Malig. Extrarenal Rhabdoid Tumor

Malignant extrarenal rhabdoid tumor (MERT), renal rhabdoid tumor, and CNS atypical teratoid/rhabdoid tumor (ATRT) are related high-grade tumors that share histologic features and identical molecular and cytogenetic abnormalities.
- Extrarenal (non-CNS) 45%
- CNS 35%
- Renal 20% (Source: SEER)

- Infants, Children
- Deep axial region: neck, paraspinal, abdomen, pelvis
- LIVER most common visceral location
- Rapid growing
- Histology:
  - Sheets of rhabdoid cells: large round to bean-shaped nuclei, prominent nucleoli, abundant eccentric eosinophilic cytoplasm
  - PAS+, diastase resistant globules
  - Keratin/EMA+
  - May be CD99/S100+
  - INI1 lost (SMARCB1 loss)
- 35% are familial : Germline SMARCB1 mutation
- Poor prognosis in all locations



INI1 (SMARCB1) lost tumors

- Usually tumors with rhabdoid cell component:
  - Epithelioid sarcoma
  - Malignant myoepithelioma 50%
  - Epithelioid MPNST 50%
  - Renal medullary carcinoma
  - ATRT
  - MERT

Chromatin Remodeling?



SWI/SNF (SWItch/Sucrose Non-Fermentable) is a nucleosome-remodeling complex (destabilizes nucleosomes).
Alter gene expression.
Function as tumor suppressor.
LOST in malignancies.

INI1 – SMARCB1
- Epithelioid sarcoma
- ATRT / MERT
- Malignant myoepithelioma
- Epithelioid MPNST
- Renal medullary carcinoma

BRG1 – SMARCA4
- SMARCA4 deficient thoracic sarcoma
- Ovarian hypercalcemic small cell carcinoma
- SMARCA4 deficient undifferentiated uterine sarcoma

Alveolar Soft Part Sarcoma



- Rare, Mostly young adults
  - Before age 30: F>M, Head & neck
  - After age 30: M>F, Deep thigh
- Histology;
  - Nests separated by delicate vessels
    - Loss of cohesion = alveolar appearance
  - Polygonal cells with well-defined borders, abundant eosinophilic granular cytoplasm, large nuclei, prominent nucleoli
  - Rhomboid intracytoplasmic crystals
- Nuclear TFE3+ = TFE3-ASPSCR1 (ASPL) fusion
  - der(17)t(X;17)(p11q25)
- Some may have desmin/S100
  - No keratin, No other neuronal/melanocytic markers
- Recurrence is rare; Mets are common (lung, bone, brain) with increasing pt age and tumor size

Clear Cell Sarcoma



* Metastasizes to lymph nodes 50%!

- "Malignant melanoma of soft parts"
- Young adults, M=F, distal lower extremities, small bowel
- Slow growing
- Histology:
  - Usu < 5 cm, pushing border
  - Nests, short fascicles
  - Epithelioid to spindled cells
  - Pale pink (rarely clear) cytoplasm
  - Prominent nucleoli
  - Wreath-like multinucleated giant cells
- IHC:
  - HMB45 stronger than S100
  - MelanA, MiTF +
  - Keratin/EMA neg
- t(12;22) EWSR1-ATF1
- Poor prognosis

PEComa

Family of tumors:
- Angiomyolipoma
- Clear cell "sugar" tumor of lung
- Lymphangioleiomyomatosis (LAM)

- Perivascular epithelioid cell tumor
  - Melanocytic / Myogenic diff
- F>>M, wide age range
- Retroperitoneum, pelvis, GI
- Histology:
  - Nests surrounded by delicate capillary vessels
  - Polygonal epithelioid cells with abundant granular eosinophilic-to-clear cytoplasm and round nuclei, small nucleoli
  - 15% are sclerosing
  - SMA+ 80% / rare desmin
  - HMB45, MelanA, MiTF / rare S100!
- So called "Malignant PEComa" = mitoses, necrosis, pleomorphism = may metastasize to liver, nodes, lung, bone





Epithelioid Tumors Differential Diagnosis Summary

Tumor	Keratin HMWK	EMA	INI-1	S100	p63 p40	CD34	CD31 ERG
Epithelioid Sarcoma	+	+	Lost	-	-	+ 50%	- +/- ERG
Malignant Rhabdoid Tumor	+	+	Lost	+/-		-	
Myoep-ithelioma	+/- Must have keratin or EMA		Lost 50%	+/- Must have S100, GFAP, calponin/SMA, or p63/p40			-
Epi MPNST	+/-		Lost 50%	+			
Epi Angiosarc	+ 50%	-	Ret				+
Melanoma	-		Ret	+			
SCC	+ +		Ret		+	-	



Melanocytic/S100-SOX10+ tumors

Tumor	S100, SOX10	HMB45, MelanA *	SMA, calponin, p63, GFAP	Keratin, EMA	INI-1	Other
Melanoma	+	+	-	-	Retained	In situ, c-kit, BRAF, NRAS
Epithelioid MPNST	+	-	-	+/- focal	Lost 50%	a/w nerve, NF patient
Myo-epithelioma	+	-	Variably +	+ for one or both	Lost 50% of malignant	EWSR 50%
Clear Cell Sarcoma	+	+	-	-	Retained	EWSR-ATF1
PEComa	Rare S100	+	+ SMA	-	Retained	-

* Caution with MiTF b/c it can be + in a number of mesenchymal neoplasms.



Epithelioid Tumors w/ Myxoid Stroma

Myoepithelioma of soft tissue

Wide age range
Malignant variant (myoepithelial carcinoma) more in children

Limbs, limb girdles

Reticular, trabecular, nested, sheet-like growth
In myxoid to hyalinized stroma
Epithelioid to spindled cells
Small uniform nuclei
Presence of nucleoli, nuclear pleomorphism, increased mitoses, necrosis may indicate malignancy

Need to show both:
Epithelial marker: Keratin or EMA
AND Myoepithelial marker: Calponin, S100, GFAP > SMA
INI1 lost in some malignant cases
EWSR rearranged in 50%

Benign cases may recur
Malignant cases recur and metastasize

http://www.dermpedia.org/conference/dr-hornicks-conference?page=4

Myoepithelial carcinoma



Epithelioid Tumors w/ Myxoid Stroma

Chordoma

Of Notochordal origin
Adults, M>F

2 specific areas of BONE: base of skull OR sacrococcygeal
With soft tissue extension
"Parachordomas" of soft tissue have mostly been determined to be myoepitheliomas

Lobules composed of strands and chords in myxoid stroma
Clear to eosinophilic often "bubbly" cytoplasm (physaliphorous cells)
Small bland nuclei
Chondroid variant (no obvious behavioral difference)
Dedifferentiated variant (adjacent high grade sarcoma)

Keratin+, EMA+, S100+
BRACHYURY+ (highly specific)

Treated with surgery, 7 year median survival
40% sacrococcygeal / vertebral metastasize to lung, etc

Dediff

* Metastasizes to lymph nodes!

Physaliphorous: Hellenistic Greek φυσαλλίς bladder (see PHYSALIS n.) + German -phore to bear



Epithelioid Tumors w/ Myxoid Stroma

Extraskeletal Myxoid Chondrosarcoma

Adults, M>F

THIGH, proximal extremities, limb girdle

Well-demarcated, multilobulated
Interconnected cells forming cords / network
Myxoid / chondromyxoid stroma
Actual cartilage usually ABSENT
Eosinophilic cytoplasm, uniform bland oval nuclei

S100+ 20%, CD117+ 30%
INI1 lost if rhabdoid features
t(9;22) EWSR1-NR4A3 *
t(9;17) TAF15-NR4A3

Aggressive: 50% recur, 50% metastasize usually to lung
Age, size, proximal location do worse

http://www.sarcomaimages.com/images/db/extraskeletal-myxoid-chondrosarcoma/470-extraskeletal-myxoid-chondrosarcoma-26-sarcoma-images.jpg

Yellow lobulated
Hemorrhage common

http://www.nature.com/modpathol/journal/v21/n11/images/journal_cover_large.jpg



Tumor	S100	Keratin	EMA	Brachyury	D240	GFAP
Myoepithelioma	+ 90%	+ 90%	+/- 60%	-	-	+/- 60%
Chordoma	+ 95%	+ 100%	+ 95%	+ 100%	-	-
Chondrosarcoma, extraskeletal myxoid	+ 20-80%	- (40%+)	- (25%+)	-	-	-
Chondrosarcoma, conventional	+ 100%	-	-	-	+ 100%	-
Chordoid meningioma	- (40%+)	- (20%+)	+ 90%	-	+ 80%	-
Chordoid glioma	+ 100%	+ 100%	+ 100%	-	+ 75%	+ 100%

Chordoid glioma
Skeletal myxoid chondrosarcoma
Enchondroma
Low grade chondrosarcoma
Chordoma
Extraskeletal myxoid chondrosaarcoma
D2-40
S100

Am J Surg Pathol. 2009 May;33(5):669-81.
doi: 10.1097/PAS.0b013e318194c566. PMID 19194275



Differential Diagnosis

Epithelioid MPNST	Malig Myoep	Malig Rhabd Tumor	Epithelioid Sarcoma
S100, SOX10 +	S100, SOX10 +/-	S100 -/+	S100, SOX10 -NEG
Keratin+ 50%	Keratin/EMA+	Keratin/EMA+	Keratin/EMA+
INI1 lost 50%	INI1 lost 50%	INI1 lost ~100%	INI1 lost ~100%
A/W Nerve or Schwannoma	EWSR rearr 50% +/- GFAP,p63,calponin	Axial/Paraspinal Kids	Proximal limb girdle Adult

Dermal Based Tumors

We'll talk about:
- DF
- DFSP
- GCF
- JXG
- AFX
- Dermal sarcoma
- PFHT



Dermal Based Tumors

Dermatofibroma (Fibrous histiocytoma)

Wide age range, F>M

Pale brown raised lesion on legs

Epidermal hyperplasia + basal hyperpigmentation
Nodular dermal infiltrate (spares papillary dermis)
Spindle cells in short, haphazard fascicles / storiform
Peripheral infiltration / Entrapment of dermal collagen
May extend into superficial subcutis (growth along preexisting septae)

SMA+/-
CD68+/- histiocytes
FXIIIa+ dendritic cells (NOT TUMOR cells)
CD34 may be + peripherally (DDx DFSP)

Variants, to name a few:
Cellular: may metastasize to nodes, lung
Epithelioid: DDx nevus, 50% epithelioid cells
Lipidized: ankle
Atypical: with monster cells
Deep: DDx DFSP, within subcutis, circumscribed, more cellular, storiform, 40% CD34+, recur 20%, rare metastases

Most common mesenchymal tumor of skin

http://www.webpathology.com/slides-13/slides/SoftTissue_Fibrohistiocytic_Dermatofibroma4.jpg

http://www.pathologyoutlines.com/images/softtissue/03_13B.jpg



Dermal Based Tumors

Dermatofibrosarcoma Protuberans (DFSP)

Young-Mid Adults

Trunk / Proximal extremities, possibly reddish

Diffuse involvement of dermis / subcutis
Honeycomb entrapment of fat, skin appendages
Whorled / storiform pattern
Uniform spindle cells with wavy nuclei, Few mitoses

CD34+ diffuse
t(17;22) COL1A1-PDGFB

Variants:
Myxoid: very pale
Pigmented: Bednar tumor
Fibrosarcomatous transformation: hypercellular, fasciculated (herringbone), mitoses, loss of CD34
Giant cell fibroblastoma: children

Local recurrences related to margins
Metastasis with fibrosarcomatous transformation



Dermal Based Tumors

Giant Cell Fibroblastoma – DFSP variant

CHILDREN, median 6 yrs

Trunk, groin, axilla

Deep dermal / subcutaneous
Honeycomb infiltration of fat
Hypocellular with myxoid/collagenous stroma
Wavy bland spindle cells and tumor giant cells
Pseudovascular spaces lined by tumor cells

CD34+
t(17;22) COL1A1-PDGFB identical to classic DFSP

50% recur, none metastasize

http://www.pedorthpath.com/giant_cell_fibroblastoma_hist_4.jpg

http://www.pathologyoutlines.com/images/softtissue/03_34B.jpg

http://www.pathologyoutlines.com/images/softtissue/03_36.jpg



Dermal Based Tumors

Juvenile Xanthogranuloma

Infants/children

Head & neck, Trunk

Nodular, polypoid growth, Flattened epidermis
In Dermis +/- superficial subcutis
Mononuclear histiocytes
Touton giant cells with peripheral foamy cytoplasm

CD68, CD163+
SMA+ in some spindle cells

May spontaneously regress or cured with excision



Dermal Based Tumors

Atypical Fibroxanthoma	Dermal Sarcoma
Elderly, M>F, Caucasians	DO NOT DIAGNOSE AFX on superficial biopsy, if you cannot see the subcutis! "atypical intradermal spindle cell neoplasm" ANY POSSIBLE AFX THAT HAS: • Infiltration into subcutis • Tumor necrosis • Lymphovascular invasion • Perineural invasion SHOULD BE REGARDED AS DERMAL SARCOMA
Sun damaged skin: Head & neck / scalp	
Nodular dermal growth with pushing border No infiltration of subcutis (may PUSH but not invade) Basically looks like an undifferentiated pleomorphic sarcoma: spindled to epithelioid cells in fascicles, pleomorphism, mitoses	
Immunohistochemical diagnosis of exclusion: May be SMA+ Exclude spindle cell carcinoma, melanoma, differentiated sarcomas	
BENIGN behavior if use strict criteria, rarely local recurrence	Potential to recur and metastasize

Typical giant cells:



Dermal Based Tumors

Plexiform Fibrohistiocytic Tumor
Children, young adults
Upper extremities (Forearm), Slowly growing
Multinodular, poorly-circumscribed Deep dermal / subcutaneous proliferation of: 1. Spindle cells in fascicles/whorls 2. Mononuclear epithelioid histiocytes 3. Osteoclast-like giant cells Rare mitoses, no necrosis
CD68+ histiocytes/giant cells SMA+ spindled fibroblasts/myofibroblasts
May recur, rarely metastasize to lymph node > lung



Giant Cell Rich Tumors

Tenosynovial Giant Cell Tumor, Localized Type	Tenosynovial Giant Cell Tumor, Diffuse Type	Giant Cell Tumor of Soft Tissue
"Giant cell tumor of tendon sheath"	"Pigmented villonodular synovitis (PVNS)"	"Osteoclastoma of soft tissue"
Adults, F>M, FINGERS (mostly near joint) Well-circumscribed mass	Younger adults, F>M, KNEE (mostly in joint) Ill-defined intra/peri-articular lesions	Adults 40s Superficial extremities
1. Mononuclear cells (synovial-like) (CD68, CD163+) 2. Foamy histiocytes 3. Osteoclast-like giant cells (less common in diffuse type) (CD68, CD45+) 4. Hemosiderin		Similar to GCT BONE Multinodular, separated by fibrous septa 1. Mononuclear cells (CD68 & SMA focal) 2. Osteoclast-like giant cells (CD68+)
CSF1 rearrangement // May recur (more in diffuse type) // Possible "Malignant" variant		Occasional recurrence, rare metastasis



Matrix-Forming Tumors in Soft Tissue

Soft Tissue Chondroma	Myositis Ossificans / Fibro-osseous Pseudotumor of Digit	Extraskeletal Osteosarcoma
Mid Adult, M>F	Young Adult, MO: M>F, FP: F>M	Mid-late Adult, M>F
Fingers > hands, toes, feet	Injury / Trauma to all parts of body	Deep soft tissue of thigh
Well-circumscribed nodules of mature hyaline cartilage separated by fibrous tissue Peripheral ossification Mineralized matrix	FP: Bone throughout MO: Cellular area in center of tumor Bony matrix at periphery ↔	Same as subtypes that arise in bone: Osteoblastic > Fibroblastic, Chondroblastic, Telangiectatic, Small cell, Well diff Bony matrix in center of tumor Cellular area at periphery
Occasional recurrence	Cured by excision (many have USP6 fusion)	Aggressive: 25% 5-year survival

Benign Myxoid Tumors

We'll talk about:
Myxoma
Superficial Angiomyxoma
Acral Fibromyxoma
Deep Angiomyxoma
Angiofibroma



Myxoma

- Adult, F>M
- Intramuscular (thigh)
- Juxta-articular (knee)
- Histology:
  - May focally show increased cellularity
  - CD34, SMA +
  - (EMA, MUC4 -) * DDx LGFMS
- GNAS mutations in intramuscular
- Mazabraud syndrome: intramuscular myxoma, skeletal fibrous dysplasia

Superficial Angiomyxoma

- Adult, M=F
- Dermis
- Histology:
  - Stellate-spindled cells in myxoid stroma
  - Prominent thin walled vessels
  - Neutrophils
  - Entrap skin adnexa
- Multiple or external ear = may be Carney complex
- May recur







Superficial Acral Fibromyxoma

- Mostly adults, M > F
- Hands, feet, peri-ungual
- May be painful
- Histology:
  - Dome-shaped mass in dermis
  - Spindle-stellate fibroblasts in myxoid-collagenous stroma
  - Small vessels
  - Mast cells
  - No adnexal entrapment / neutrophils
- Loss of Rb like SCL
- Rare recurrence





Deep (Aggressive) Angiomyxoma

- Adult Females
- Perineum, Pelvis
- Histology:
  - Lobulated, infiltrative
  - Hypocellular stellate-spindle cells in loose matrix
  - VESSELS of varying caliber
  - ER/PR+
  - Variable SMA, desmin, CD34
- HMGA2 rearrangements in some
- Local recurrence with + margins





Angiofibroma of Soft Tissue

- Adult, F>M
- Lower extremity
- Histology:
  - Circumscribed
  - Small bland spindle cells in collagenous-myxoid stroma
  - Numerous branching thin-walled to ectatic vessels
- t(5;8) AHRR-NCOA2 > t(7;8) GTF2I-NCOA2
- Rare recurrence







Benign Myxoid / Vascular

Myxoma	Superficial Angiomyxoma	Superficial Acral Fibromyxoma	Deep Aggressive Angiomyxoma	Soft Tissue Angiofibroma
Adult, F>M	Adult, F=M	Adult, M>F	Adult, F	Adult, F.M
Intramuscular (thigh) Juxta-articular (knee)	Dermal	Hands, Feet, Peri-ungual +/- Painful	Pelvis, Perineum	Deep lower extremity
May focally show increased cellularity CD34, SMA + (EMA, MUC4 -) * DDx LGFMS	Stellate-spindled cells in myxoid stroma Prominent thin walled vessels Neutrophils Entrap skin adnexa	Dome-shaped mass in dermis Spindle-stellate fibroblasts in myxoid-collagenous stroma Small vessels Mast cells No adnexal entrapment / neutrophils	Lobulated, infiltrative Hypocellular stellate-spindle cells in loose matrix VESSELS of varying caliber ER/PR+ Variable SMA, desmin, CD34	Circumscribed Small bland spindle cells in collagenous-myxoid stroma Numerous branching thin-walled to ectatic vessels
GNAS mut: 50% intramuscular Mazabraud syndrome: intramuscular myxoma, fibrous dysplasia	Multiple or external ear = may be Carney complex May recur	Loss of Rb like SCL Rare recurrence	HMGA2 rearrangements in some Local recurrence with + margins	t(5;8) AHRR-NCOA2 > t(7;8) GTF2I-NCOA2 Rare recurrence



Misc

We'll talk about:
FHI
HFLT
MIFS
PMT
AFH



Fibrous Hamartoma of Infancy

- Before age 2, Axilla/Inguinal
- Ill-defined
- Histology:
  1. Fibrous tissue SMA+
  2. Immature mesenchymal tissue SMA-
  3. Adipose tissue S100+
- Cured by local excision
  - May recur if + margins



HFLT – MIFS

Hemosiderotic Fibrolipomatous Tumor (HFLT)	Hybrid HFLT-MIFS	Myxoinflammatory Fibroblastic Sarcoma (MIFS)
Middle age Adults, F>M, slow growing	"HFLT with sarcomatous progression"	Middle age Adults, M=F, slow growing
DISTAL EXTREMITIES (dorsal foot)		DISTAL EXTREMITIES (hands)
1. Fascicles of fibroblastic spindle cells 2. Adipocytes 3. Hemosiderin laden macrophages 4. Osteoclast-like giant cells	Less frequent Reed-Sternberg-like cells More frequent arborizing vasculature similar to myxofibrosarcoma	Multinodular, fibrous and myxoid zones Spindled-epithelioid cells with inflammatory infiltrate Virocyte-like or Reed-Sternberg-like cells
CD34, Calponin + TGFBR3-MGEA5 t(1:10) →	TGFBR3-MGEA5 t(1:10)	Variable CD68, CD34, SMA MGEA5 rearrangement in some (not w/TGFBR3) BRAF rearrangement in others
Recurrence with + margins	May metastasize	Recurrence common, distant metastasis rare



Phosphaturic Mesenchymal Tumor

- VERY RARE
- Middle aged adults
- All soft tissue sites, bone
- Histology:
  - Bland spindle-stellate cells in "smudgy matrix"
  - "Grungy" calcification
  - Microcysts
  - FGF ISH can be performed if needed
- Elevated serum FGF23 = tumor-induced osteomalacia
- May recur, Osteomalacia regresses after excision

Hypophosphatemic Ricketts, Hyperphosphaturia



Angiomatoid Fibrous Histiocytoma

- Teens, young adults
- Limbs, Lymph node-rich areas
- Histology
  1. Fibrous pseudocapsule
  2. Pericapsular lymphoid tissue
  3. Nodules of spindled-histiocytoid cells with syncytial growth
  4. Pseudoangiomatous blood-filled spaces
  - 50% EMA, CD99, desmin
- EWSR1-CREB1 t(2;22) 90%
- FUS-ATF1, EWSR-ATF1
- May recur, Rare metastasis to lymph node

http://www.pathologyoutlines.com/images/softtissue/03_43.jpg

Thank you!

Questions? Comments?
Multiple Choice Questions to Follow

University of Chicago Photographic Archive, [apf7-02389], Special Collections Research Center, University of Chicago Library.



1. A 54 year old male presents with discomfort and a palpable mass on the sole. An excisional biopsy is performed (pictured). What is the most likely diagnosis?

A. Fibroma of tendon sheath.
B. Giant cell tumor of tendon sheath.
C. Superficial fibromatosis.
D. Nodular fasciitis.
E. Clear cell sarcoma.

1. A 54 year old male presents with discomfort and a palpable mass on the sole. An excisional biopsy is performed (pictured). What is the most likely diagnosis?



A. Fibroma of tendon sheath.
B. Giant cell tumor of tendon sheath.
C. Superficial fibromatosis.
D. Nodular fasciitis.
E. Clear cell sarcoma.

Superficial fibromatosis most frequently occurs on the palms, fingers, and soles. It infiltrates tendon, and is composed of sheets and fascicles of plump to stellate spindle cells. The plantar form often contains multinucleated giant cells.

---

2. A 37 year old female presents with knee pain. Imaging shows multiple intra-articular nodules at the periphery of the joint. Arthroscopy with curettage is performed. What is the most likely diagnosis?



A. Rheumatoid arthritis.
B. Acute synovitis.
C. Lipoma arborescens.
D. Tenosynovial giant cell tumor, localized type.
E. Pigmented villonodular synovitis.

---

2. A 37 year old female presents with knee pain. Imaging shows multiple intra-articular nodules at the periphery of the joint. Arthroscopy with curettage is performed. What is the most likely diagnosis?



A. Rheumatoid arthritis.
B. Acute synovitis.
C. Lipoma arborescens.
D. Tenosynovial giant cell tumor, localized type.
E. Pigmented villonodular synovitis.

PVNS (also known as tenosynovial giant cell tumor, diffuse type) occurs as a multinodular lesion in large joints. It classically contains foamy histiocytes, bland mononuclear cells, hemosiderin, and giant cells. The uninodular form is called "localized type" and more frequently occurs in small joints.

---

3. A 68 year old male complains of neck pain. An 11 cm mass is identified in the subcutaneous tissue above the trapezius. What is the most likely immunoprofile of this lesion?



A. MDM2+, CDK4+
B. CD34+, Rb lost
C. SMA+, Beta-catenin+
D. S100+, desmin negative
E. CD68+, CD163+

---

3. A 68 year old male complains of neck pain. An 11 cm mass is identified in the subcutaneous tissue above the trapezius. What is the most likely immunoprofile of this lesion?



A. MDM2+, CDK4+
B. CD34+, Rb lost
C. SMA+, Beta-catenin+
D. S100+, desmin negative
E. CD68+, CD163+

Atypical lipomatous tumor (while more frequent in extremities) can occur in any site. The atypical nuclei, myxoid stroma, variably-sized fat, lack of ropey collagen, and lack of floret cells make ALT more likely than spindle cell / pleomorphic lipoma, even in this location.

---

4. Based on your answer to question 3, what is the most likely genetic anomaly?



A. Amplification of chromosome 12q
B. Loss of chromosome 13q
C. Mutation in CTNNB gene
D. DDIT3-FUS rearrangement
E. 11q rearrangement

4. Based on your answer to question 3, what is the most likely genetic anomaly?



A. Amplification of chromosome 12q
B. Loss of chromosome 13q
C. Mutation in CTNNB gene
D. DDIT3-FUS rearrangement
E. 11q rearrangement

A. Present in Atypical lipomatous tumor / well-diff / dediff liposarc
B. Present in spindle cell / pleomorphic lipoma / mammary type myofibrbllastoma
C. Present in fibromatosis (desmoid)
D. Present in myxoid liposarcoma
E. Present in hibernoma

---

5. You receive a consult of H&E and immunostains but minimal clinical information. The pictured tumor is positive for keratin, desmin (perinuclear dot), and CD99. What is the most likely clinical scenario?



A. A 9 year old female with a tibial lesion.
B. A 76 year old male with lung and vertebral lesions.
C. A 17 year old male with omental lesions.
D. A 55 year old female with perineal lesion.
E. A 34 year old female with ankle lesion.

---

5. You receive a consult of H&E and immunostains but minimal clinical information. The pictured tumor is positive for keratin, desmin (perinuclear dot), and CD99. What is the most likely clinical scenario?



A. A 9 year old female with a tibial lesion.
B. A 76 year old male with lung and vertebral lesions.
C. A 17 year old male with omental lesions.
D. A 55 year old female with perineal lesion.
E. A 34 year old female with ankle lesion.

Desmoplastic small round cell tumor shows multiphenotypic differentiation by immunohistochemistry and EWSR1-WT1 rearrangement. It classically occurs in the abdomen of teenage males.

---

6. Pictured is an excisional biopsy of a raised lesion from the leg of a 43 year old female. What is the most likely diagnosis?



A. Atypical fibroxanthoma
B. Dermal sarcoma
C. Dermatofibroma
D. Dermatofibro-sarcoma protuberans
E. Perineurioma

---

6. Pictured is an excisional biopsy of a raised lesion from the leg of a 43 year old female. What is the most likely diagnosis?



A. Atypical fibroxanthoma
B. Dermal sarcoma
C. Dermatofibroma
D. Dermatofibro-sarcoma protuberans
E. Perineurioma

Dermatofibrosarcoma protuberans shows a storiform growth pattern with honeycomb infiltration of fat. AFX, DF, and perineurioma should have no to minimal fat infiltration. Dermal sarcoma is usually histologically high grade.

---

7. A 26 year old male undergoes excision of a lower leg mass. What is the most likely diagnostic genetic anomaly?



A. Nonspecific genetics
B. t(17;22) COL1A1-PDGFB
C. USP6 rearrangment
D. GNAS mutation
E. t(7;16) FUS-CREB3L2

7. A 26 year old male undergoes excision of a lower leg mass. What is the most likely diagnostic genetic anomaly?



A. Nonspecific genetics
B. t(17;22) COL1A1-PDGFB
C. USP6 rearrangement
D. GNAS mutation
E. t(7;16) FUS-CREB3L2

This is a low grade fibromyxoid sarcoma. GNAS mutation is present in intramuscular myxoma. USP6 in nodular fasciitis. COL1A1 in DFSP. Nonspecific genetics in myxofibrosarcoma.

8. A 13 year old male has a lytic lesion in the femur with soft tissue extension. Biopsy of the soft tissue component is performed. What immunostains are most likely positive in this lesion?



A. CD99, FLI1
B. Desmin, Myogenin
C. CD45, PAX5
D. EMA, TLE1
E. S100, HMB45

8. A 13 year old male has a lytic lesion in the femur with soft tissue extension. Biopsy of the soft tissue component is performed. What immunostains are most likely positive in this lesion?



A. CD99, FLI1
B. Desmin, Myogenin
C. CD45, PAX5
D. EMA, TLE1
E. S100, HMB45

This is a Ewing sarcoma. Some cases (formerly atypical Ewing sarcoma) have large nuclei and prominent nucleoli. Worse behavior in "atypical" Ewing has not been confirmed.

9. A 72 year old female presents with a mass in the lower abdominal wall. What immunopanel will be most helpful in determining the diagnosis?



A. Keratin, ERG, INI1, SOX10
B. Keratin, CD31, ERG, HMB45
C. Vimentin, EMA, GFAP, S100
D. Vimentin, PAX8, ERG, BRG1
E. Keratin, Vimentin, ER, PR

9. A 72 year old female presents with a mass in the lower abdominal wall. What immunopanel will be most helpful in determining the diagnosis?



A. Keratin, ERG, INI1, SOX10
B. Keratin, CD31, ERG, HMB45
C. Vimentin, EMA, GFAP, S100
D. Vimentin, PAX8, ERG, BRG1
E. Keratin, Vimentin, ER, PR

The differential diagnosis is broad, including melanoma, carcinoma, epithelioid sarcoma, epithelioid angiosarcoma, epithelioid MPNST, malignant myoepithelioma. This is an example of epithelioid sarcoma.

10. A 35 year old female presents with a 1.5 cm mass on the left ankle. It is rubbery and mobile. What is the most likely diagnosis?



A. Leiomyoma
B. Angioleiomyoma
C. Fibroma of tendon sheath
D. Burnt out tenosynovial giant cell tumor
E. Angiofibroma of soft tissue



10. A 35 year old female presents with a 1.5 cm mass on the left ankle. It is rubbery and mobile. What is the most likely diagnosis?

A. Leiomyoma
B. Angioleiomyoma
C. Fibroma of tendon sheath
D. Burnt out tenosynovial giant cell tumor
E. Angiofibroma of soft tissue

This is a well-circumscribed spindle cell lesion. It contains vessels with muscular walls similar to the pale eosinophilic smooth muscle bundles in the background. This is an angioleiomyoma.

This page was intentionally left blank.

Liver Pathology

David H. Gonzalo, MD
Associate Professor
University of Florida College of Medicine
Email: Hernand3@ufl.edu
@DHGGIpath
October 2020



General Approach to Liver Biopsy

- Purpose is to answer important clinical questions
- Final interpretation should synthesize all findings:
  - clinical, laboratory, radiographic, and molecular
- Interpret biopsy
  - blinded to clinical information
  - using a systematic approach (pattern recognition)
    - Architecture
    - Portal triad
    - Lobular region
    - Central vein

- Highlights the connective tissue (BLUE) in the portal area



Hepatocytes are separated from sinusoids by the "space of Disse" (highlighted in pink) that contains Ito (stellate) cells that are normally invisible on H&E. Ito cells store Vit A !! Ito cells can turn into myofibroblasts → make collagen → fibrosis ITO CELL HYPERPLASIA IN HYPERVITAMINOSIS A

Liver – Submicroscopic Histology

Sinusoids are low pressure vascular channels lined by endothelial cells that receive blood from hepatic artery and portal vein and deliver it to the central vein and in the process *perfuse the hepatocytes*

Direction of blood flow: Portal vein → Sinusoids → Central vein [*Also, Hepatic artery→Sinusoids→Central vein*]

Direction of bile flow: Hepatocytes → Bile canaliculi → Bile ductule

Sinusoids also contain **Kupffer cells** → Specialized hepatic macrophages!!

Outline

- Patterns of injury
- Infections
- Vascular abnormalities
- Tumors



Patterns of Injury

	Predominant Location	Key Features
Acute hepatitis	Lobular (zone II &III)	Inflammation/necrosis Lobular disarray No fibrosis
Chronic hepatitis	Portal-periportal (zone I)	Portal inflammation Parenchymal injury Piecemeal necrosis
Biliary injury/Cholestasis	Portal-periportal (zone I)	Large duct obstruction: Cholestasis, portal edema, ductular proliferation Chronic: "Cholate stasis", Copper, loss of bile ducts
Steatohepatitis (alcoholic or nonalcoholic)	Central (zone III)	Fat Mallory bodies Pericellular fibrosis



Pattern 1

Acute Hepatitis Pattern:

- Lobular disarray
- Marked lobular inflammation
- Variable necrosis
- No significant fibrosis

O'Grady JG, Schalm SW, Williams R. Acute liver failure: redefining the syndromes. Lancet 1993; 342:273-275



Etiology of Acute Hepatitis

Hepatotropic viruses
Hepatitis A, B, E
Non-hepatotropic viruses
CMV, EBV

Autoimmune hepatitis
Alcoholic hepatitis
Drug (acetaminophen)
Wilson disease







Types of Hepatitis

	A	B	C	D	E
Source	Fecal/oral	Blood/ Parenteral	Blood/ Parenteral	blood/ Parenteral	Fecal/oral
Virus	RNA	DNA	RNA	RNA	RNA
Route of transmission	fecal-oral	percutaneous mucosal	percutaneous mucosal	percutaneous mucosal	fecal-oral
Chronic Infection	No	Yes	Yes	Yes	No
HCC	No	Yes	Yes	Yes	No



Pattern 2
Chronic hepatitis



Chronic Hepatitis

- Clinical definition
  - > 6 months of symptoms and signs and biochemical indicators
- Pathologic
  - Persistent necroinflammation of the liver
- Liver biopsy is used to:
  - Confirm the presence of necroinflammation
  - Evaluate the severity (grading)
  - Assess the scarring (staging)
  - Suggest possible etiology
  - Provide prognosis



Etiology of Chronic Hepatitis

Blue portal areas !!!

- Chronic hepatitis B (+/- HDV) C, D
- Chronic autoimmune hepatitis
- Medication induced
- Wilson's disease
- Hepatic involvement by lymphoma
- Alpha 1 anti-trypsin deficiency

MIMIC CHRONIC HEPATITIS:

- Primary biliary cholangitis (PBC)
- Primary sclerosing cholangitis (PSC)



Histology of Chronic Hepatitis

- Necroinflammation
  - Portal/periportal based inflammation
  - Periportal hepatocyte injury/necrosis
  - Variable interface activity
  - Variable lobular inflammation & hepatocyte injury/necrosis
- +/- Scar/fibrosis
  - Portal



Chronic hepatitis C

Hepatic steatosis: feature of **genotype 3**



Macronodular cirrhosis → Average nodule size > 3 mm (Chronic hepatitis B and Hepatitis C)
Micronodular cirrhosis → Average nodule size < 3 mm (Alcohol; Hemochromatosis)



Hepatitis B

- DNA virus
- Sex, blood born, mother to child
- Chronic infection: Adults 5%, Newborns 90%
- Vaccine available

HBsAg
HBcAg
DNA Polymerase
42nm
HBeAg



Ground glass hepatocytes

Hepatitis B

- Cytoplasmic inclusion pushes nucleus to side due to proliferating endoplasmic reticulum responding to HbsAg
- Positive for HBsAg

Sanded nuclei
Pink nuclear inclusions containing HBcAg

Glycogen pseudoground glass change in hepatocytes.
Wisell J, Boitnott J, Haas et al, Am J Surg Pathol. 2006 Sep;30(9):1085-90



Autoimmune hepatitis

- Predominantly middle aged women
  - Type 1: Most common, ANA and/or ASMA. P-ANCA.
  - Type 2: Most common in children: anti LKM-1 (more severe disease)
  - Type 3: Adults 30-50 years old: soluble liver antigen or liver pancreas antigen.
- History of other autoimmune diseases
- Elevated IgG, elevated AST/ALT 5-10 times upper limit of normal
- Histology: Untreated AIH usually chronic hepatitis with prominent interface activity and lobular injury
- Plasma cells not required for diagnosis
- Only chronic hepatitis that responds to corticosteroid



Autoimmune hepatitis-Histology

- Marked interface inflammation with bridging necrosis and plasma cells

Severe inflammation

Plasma cells

Bridging necrosis



Centrilobular inflammation

Giant cell transformation



α_1-antitrypsin deficiency

- Mutation in the **SERPINA1 gene** on chr 14q. Autosomal codominant.
  - Alpha-1-antitrypsin folds abnormally and retained in the RER.
  - Deficiency leads to emphysema and chronic liver disease.
- Normal gene product is PiM
  - PiS and PiZ are most common deficiency alleles
  - PIZZ is most severe. Mmmm, Mmmmm good, PiZZ poor
- Periportal PAS/D positive globules (also IHC available).
- Globules not present in first 3-4 months of life (serum testing).
- Patterns: neonatal cholestatic, normal liver, cryptogenic cirrhosis.



Hepatocellular A1AT accumulation

PAS-D

α_1-antitrypsin IHC

Clinical and histologic features of adults with alpha-1 antitrypsin deficiency in a non-cirrhotic cohort.
Clark VC, Marek G, Liu C et al. J Hepatol. 2018 Dec;69(6):1357-1364



Hereditary Hemochromatosis

- Autosomal recessive genetic disorder resulting in increased iron absorption from the gut
- HFE gene mutations, 2 most common
  - C282Y and H63D
  - C282Y cysteine to tyrosine substitution at amino acid position 282 (85% of patients are homozygous for this mutation)
  - C282Y has highest iron overload
- Most asymptomatic. Triad: Cirrhosis, diabetes and skin pigmentation
- Iron in periportal hepatocytes moves to zone III
- Genetic testing and quantitative iron in patients with histologic iron overload
- DD: Iron overload due to hemolysis, transfusions.

Hepatocellular iron accumulation

Prussian blue stain

Look at the next 3 slides and give me your diagnosis?





Green copper deposit at periphery of iris



Wilson disease

- Mutated ATP7B gene, a copper transport protein in Golgi with accumulation of copper in liver and other organs
- Diagnosis:
  - ↓Ceruloplasmin—not sensitive or specific
  - 24 hour urinary copper
  - Tissue copper quantification is definite in most cases
  - Targeted next generation sequencing
- Liver failure and neurologic signs
- Liver morphology
  - Steatosis, hepatitis, glycogenated nuclei, Mallory hyaline, copper by Rhodanine (UNRELIABLE)

Hepatic features of Wilson disease.
Boga S, Ala A, Schilsky I. Handb Clin Neurol. 2017;142:91-99.



Pattern 3
Bile duct injury/cholestasis



Bland Cholestasis

Major finding: Lobular cholestasis without significant

- Androgenic steroids
- High-dose estrogen therapy
- Inherited defects in bilirubin metabolism



Sepsis: cholestasis
Bile in proliferating bile ductules



Cholestasis with Other Changes

- Etiology
  - Large bile duct obstruction (LDO)
  - Ischemic cholangitis
  - Primary sclerosing cholangitis (PSC)
  - Primary biliary cirrhosis (PBC)
  - DILI



Large Bile Duct Obstruction (LDO)

- **Cholestasis with hepatocyte changes**
  - **Cholestasis**
  - **Feathery degeneration**
  - **Xanthomatous changes**
  - **Bile infarct**
- **Portal tract**
  - **Edema and bile ductular reaction**
- **Bile duct**
  - **Normal, reactive change, to neutrophilic inflammation**
- **Fibrosis**
  - **Portal fibrosis if cholestasis persists**



LDO: portal edema, bile duct injury

Periportal copper:
chronic cholestasis

LDO: bile ductular reaction

Primary Biliary Cholangitis

(previously known as primary biliary cirrhosis)

- Autoimmune disease with progressive destruction and loss of intrahepatic bile ducts.
- Females, 40 – 60 years old; fatigue, abdominal pain, jaundice, pruritus.
- Alkaline phosphatase elevation.
- AMA positive (M2 subtype most specific)(+ in 95% of cases).↑IgM
- Histology
  - Portal based inflammation, with florid bile duct lesion.
  - Noncaseating granulomas
  - Chronic cholestasis: Increased copper accumulation
  - Late in disease: biliary type cirrhosis.

Primary Biliary Cholangitis and Autoimmune Hepatitis.
Gonzalez RS, Washington *Surg Pathol Clin.* 2018 Jun;11(2):329-349



Florid duct lesion

Granulomas

Primary Sclerosing Cholangitis (PSC)



Beading, strictures at branch points

- Fibroinflammatory disease affecting intra and extrahepatic biliary tree.
- M>F (5 to 1). 25 – 60 year old.
- pANCA (50% of pts). Elevated alk phos.
- Assoc with UC (65% occur in patients with UC)
- Early: biliary obstruction pattern with bile ductular proliferation).
- Intermediate: Fibrous obliteration of bile ducts with periductal fibrosis
- Late changes: Ductopenia and biliary cirrhosis.
- ↑ cholangiocarcinoma
- Small duct PSC: variant involving only small intrahepatic bile ducts with normal cholangiography (5% of all PSC).



PSC: ductular reaction

PSC: bile duct injury and periductal fibrosis



Pattern 4

Fatty liver disease

Steatosis vs steatohepatitis

Macrovesicular steatosis



Main causes

- ✓ Obesity, DM, dyslipidemia
- ✓ Metabolic syndrome
- ✓ PCOS
- ✓ Starvation/Total parenteral nutrition
- ✓ Alcohol
- ✓ Meds:
  - Amiodarone
  - Methotrexate
  - Tamoxifen
  - Corticosteroids
- ✓ Wilson disease
- ✓ Infection (hepatitis C)

Microvesicular steatosis



Pregnancy (AFLD)
Reye's syndrome
Antiretrovirals
Valproic acid
Mushrooms
Tetracycline

Steatohepatitis

Required histologic criteria

- Steatosis (≥ 5%)
- Hepatocellular injury (ballooned hepatocytes):
  - +/- Mallory-Denk bodies
    - Ropey, pink material
    - Intermediate filaments metabolized by cytokeratin from liver cell injury
      - » p62+, ubiquitin+
- Inflammation (lobular)
- +/- Fibrosis (perivenular/pericentral) starting in zone III (Modified Brunt staging system)
- Non-essential features: Megamitochondria, glycogenated nuclei, lipogranulomas

Non-alcoholic steatohepatitis



Alcoholic steatohepatitis



Causes of Mallory Hyaline

- ✓ Wilson disease
- ✓ Liver lesions: FNH and HCC
- ✓ Chronic cholestatic disorders like PBC
- ✓ Steatohepatitis, especially alcoholic
- ✓ Medications: amiodarone



The Pathology of Alcoholic Liver Disease.
Alpert L, Hart J. Clin Liver Dis. 2016 Aug;20(3):473-89.



Masson trichrome

Periductal fibrosis (chicken wire)



Mimicker of steatohepatitis
GLYCOGENIC HEPATOPATHY

- ✓ Hepatomegaly.
- ✓ Marked transaminase elevation associated with poorly controlled DM.
- ✓ Abundant glycogen (diffuse pale staining) in hepatocytes.
- ✓ Absence of inflammation

Outline

- Patterns of injury
- **Infections**
- Vascular abnormalities
- Tumors



FITE

GRANULOMAS IN LIVER…
Think about:

- Sarcoidosis (associated with fibrosis, adjacent or within granulomas)
- Infections (TB, MAI, histoplasmosis)
- Medication (allopurinol)
- PBC
- Foreign substance
- Systemic granulomatous diseases

Hepatic Granulomas: A Review With Emphasis on Infectious Causes.
Lamps LW.. Arch Pathol Lab Med. 2015 Jul;139(7):867-75



Question 1

A 42 year old woman with hepatosplenomegaly, wheezing and coughing is found to have fivefold elevation of alkaline phosphatase and elevated levels of angiotensin converting enzyme (ACE). A liver biopsy demonstrates the following findings. What is your diagnosis:

a. Tuberculosis
b. Hepatitis C
c. Primary biliary cholangitis
d. Sarcoidosis
e. Q fever

The Osler Institute www.osler.org



GMS

thin based bud

Cryptococcus

Mucicarmine



Q fever (Coxiella burnetti)

Fibrin ring granuloma





Echinococcus

- E. granulosus is most frequent species of Echinococcus tapeworm
- Cystic fluid can cause anaphylaxis
- Laminated membrane
  - Zebra-like pattern
  - GMS+
- Scolex
  - Hooklets GMS+, birefringent

HYDATID SAND: contains capsules, free scolices and loose hooklets





Herpes (HSV)

- Liver with necrosis
- Smudged nuclei
- Marginated chromatin
- Multinucleated
- Molding

PUNCHED-OUT AZONAL PATTERN OF NECROSIS THINK:
HSV, ADENOVIRUS

The Osler Institute www.osler.org



Outline

- Patterns of injury
- Infections
- Vascular abnormalities
- Tumors



Ischemia

- Cardiac failure, sepsis, etc
- Nutmeg liver
  - Reddish discoloration representing areas of necrosis and hemorrhage
- Zone 3 coagulative necrosis
  - Can also be seen in acetaminophen toxicity

Portal areas preserved
Zone 3 necrosis



Acetaminophen toxicity

Acellular zonal(3) necrosis: Ischemia and acetaminophen



Budd-Chiari

- Obstruction to hepatic venous outflow.
- From thrombosis of 1 or more hepatic veins at their openings into IVC.
- Etiologies: Hypercoagulable status, infections, malignancies.
- Sinusoidal dilation, congestion, hemorrhage and hepatocyte atrophy.
- DD: Right heart failure and sinusoidal obstruction syndrome.



Sinusoidal obstruction syndrome

- Toxic damage to sinusoids and zone 3 hepatocytes.
- Due to cancer chemotherapy (oxaliplatin) and s/p hematopoietic stem cell transplant due to alkylating myeloablative drugs.
- Dilated sinusoids with hepatocyte atrophy.
- Obliteration of small central veins by subendothelial swelling.
- Sinusoidal fibrosis.

Dilated sinusoids
Obliteration of central vein



Nodular regenerative hyperplasia (NRH)

- Ischemic atrophy & secondary nodular hyperplasia in areas of good blood flow
- Regenerative nodules (1-3 mm) throughout liver, but little to no fibrosis
- Rheumatologic disorders, PV, lymphoma, azathioprine
- Portal hypertension
  - Reticulin network compressed between nodules
  - Variation in size of hepatocytes

Vague liver nodule, no cirrhosis
Reticulin network compressed b/t nodules

Outline

- Patterns of injury
- Infections
- Vascular abnormalities
- Tumors

Hepatic adenoma

- Risk factor: excess estrogen or androgen, glycogen storage disease.
- Young to middle aged women on OC or with fatty liver disease.
- Solitary, well-circumscribed in non cirrhotic liver
- Benign hepatocytes
  - One to two cells thick cords
  - Clear cell change—glycogen
- No portal triads: Naked arteries or veins
- Intact reticulin. Only focal CD34 capillarization
- Surgery to prevent bleeding/rupture
- Malignant transformation is rare, but can occur

Hepatic Adenomas: Classification, Controversies, and Consensus.
Torbenson M. Surg Pathol Clin. 2018 Jun;11(2):351-366





Hepatic adenoma (type 3, inflammatory)

Dilated sinusoids, no portal tracts, hemorrhage
May be multiple masses, no central scar

Bordeaux Classification

Subtype	Hepatocyte nuclear factor 1 α mutated (HNF1α) inactivated	βcatenin-activated adenoma	Inflammatory hepatocellular adenoma	Not otherwise specified
Frequency	30%(90% women)	10% (60% women)	50% (90% women)	10%
Steatosis	Common	Uncommon	Uncommon	Uncommon
Association with HCC	Rare	Common (aprox 50%)		
Histologic features	Steatosis	Cytologic atypia, architectural atypia (pseudoglands)	Inflammation, sinusoidal dilation and congestion	No defining features
Clinical features		Obesity	Obesity	
IHC	Liver fatty acid binding protein (LFABP) negative	Nuclear beta-catenin + (can be focal) Diffuse glutamine-synthetase	JAK/STAT pathway activation via mut in IL6ST and STAT3: C-reactive protein + Serum amyloid A +	No mutations for HNF1α or β catenin

New kids on the block: Sonic hedgehog-activated adenoma and argininosuccinate synthase-positive adenoma. Both high risk of bleeding.

Focal nodular hyperplasia (FNH)

- More common in women (70%) in 3rd-4th decade:
  - No assoc OCPs
- Unencapsulated, well-circumscribed.
- Non-cirrhotic background. No malignant potential.
- Hepatocytes divided into nodules by thick walled arteries in fibrous septae
- Ductular reaction at junction between septa and parenchyma.
- Thick walled vessels.
- Sporadic or post-chemotherapy or radiation in pediatric malignancies.
- Map-like pattern in glutamine synthetase stain.

The Osler Institute www.osler.org



central scar



FNH



H&E	CK 7	Glutamine synthetase-Map-like pattern
Fibrous septae	Proliferating ductules	GS in normal liver



Look at the next 2 slides and give me your diagnosis?

Bile stained nodule in cirrhosis





HEPATOCELLULAR CARCINOMA

- Male predominance, 65 y/o
- Abdominal pain and/or mass.
- Constitutional symptoms.
- Serum AFP elevated in only 1/3 of pts.
- Usually arises in cirrhotic liver (look for occult chronic liver disease).
- Common etiologies: Etoh, Hep B and C, NAFLD.

Malignant liver tumors in cirrhosis

The Osler Institute www.osler.org

HEPATOCELLULAR CARCINOMA

HCC mimics normal liver cells

- 4 major patterns: trabecular, solid, pseudoglandular and macrotrabecular.
- Histologic variants:
  - Steatohepatitic : Marked fat, inflammation and fibrosis
  - Clear cell : Abundant clear cytoplasm (>90% of tumor)(DD: clear cell carcinoma of kidney and ovary)
  - Combined HCC-cholangiocarcinoma: worse prognosis. 2 distinct morphologies on both H&E and IHC.

Morphologic Subtypes of Hepatocellular Carcinoma.
Torbenson MS. Gastroenterol Clin North Am. 2017 Jun;46(2):365-391



Pseudoglandular structures

Single layer of tumor cells, dilated bile canaliculi

Trabecular architecture

Enlarged hepatic trabeculae surrounded by sinusoids without fibrous stroma



Mallory bodies

Fibrinogen

A1AT



HCC vs mimickers IHC panel:

Metastatic neuroendocrine tumors, paragangliomas, adrenal cortical ca

CD10

Canalicular staining

HepPar1

Arginase

Glypican-3



HCC vs benign hepatocytes panel

Normal reticulin

Glypican-3: only 50% well diff HCC

Abnormal reticulin

CD34

Sinusoidal capillarization, endothelial cells line sinusoids



Question 2

A 30 year old man is found to have an 8 cm liver lesion. No underlying chronic liver disease. A biopsy shows malignant cells that are + for CK7 and CD68.

Central scar



Abundant eosinophilic cytoplasm, vesiculated nuclear chromatin with prominent nucleoli and intratumoral fibrosis.



What is your diagnosis:

a) Hepatic hemangioma
b) Fibrolamellar carcinoma
c) Epithelioid hemangioendothelioma
d) Focal nodular hyperplasia
e) Angiosarcoma



Fibrolamellar carcinoma

- 10-35 years old; eosinophilic carcinoma with lamellar fibrosis
- Differs from other HCC
  - No underlying liver disease. AFP normal level
- Pale bodies (fibrinogen and albumin), hyaline globules (alpha -1 AT)
- Positive for CK7 and CD68 (lysosomes)
- Microdeletion in chr 19 with DNAJB1-PRKACA fusion (FISH)
- Similar prognosis to HCC in noncirrhotic livers.
- DD: scirrhous HCC (>50% of tumor has intratumoral fibrosis).

Fibrolamellar carcinoma: A histologically unique tumor with unique molecular findings. Graham RP, Torbenson MS. Semin Diagn Pathol. 2017 Mar;34(2):146-152



Cholangiocarcinoma



Cholangiocarcinoma (CC)

- Malignant adenocarcinoma arising from duct epithelium.
- 60 years of age. Extrahepatic, hilar (Klatskin tumor), intrahepatic.
- Desmoplastic fibrous stroma, unlike HCC; lymphatic invasion common.

  Positive markers= Mucicarmine (not all), CK7+, CK19+ MOC31+.

  Negative markers= HepPar-1, arginase, CK20, Glypican-3
- Risk factors: alcohol, chronic viral hepatitis, fluke infection (Clonorchis sinensis), PSC (hilar), hepatolithiasis, thorotrast, choledocal cysts.
- DD: Metastatic pancreatobiliary carcinoma, metastatic adenoca of other origins, poorly differentiated HCC
- Lymphocyte rich variant: Usually EBV positive.



Cholangiocarcinoma



Hemangioma

- – Most common benign tumor of liver (usually solitary).
- – Young adult women.
- – May be resected due to bleeding risk.
- – Flat endothelial cells and fibrous stroma.
- Histological patterns: cavernous (common), anastomosing and capillary.



Mean age 47 years 60% women



Epithelioid Hemangioendothelioma (EHE)

- Low-grade malignant vascular neoplasm, less aggressive than typical angiosarcoma.
- Majority multinodular; ill defined border.
- Epithelioid tumor cells embedded in abundant myxoid or hyalinized stroma. Often intracytoplasmic vacuoles with RBCs.
- IHC: + for FLI-1, ERG, CD31. Pitfall: Keratins + in 10-30% of cases.
- t(1;3) translocation, resulting in a *WWTR1-CAMTA1* fusion gene.
- Metastases to abdominal LNs, lung, omentum or peritoneum (histology not prognostic).

Hepatic Epithelioid Hemangioendothelioma.
Studer LL, Selby DM. Arch Pathol Lab Med. 2018 Feb;142(2):263-267



Board type question, What is your diagnosis?



INHERITED DEFECTS IN BILIRUBIN METABOLISM

Disease	Gene	Bilirubin elevation	Clinical and histology
Crigler-Najjar type 1	UGT1A1	Unconjugated	Jaundice first week of life Bland cholestasis
Crigler-Najjar type 2	UGT1A1	Unconjugated	Cholestasis teenager Bland cholestasis
Gilbert's syndrome	UGT1A1	Unconjugated	Episodes of jaundice Bland cholestasis
Rotor syndrome	SLCO1B1 SLCO1B32	Conjugated	Episodes of jaundice Bland cholestasis
Dubin-Johnson syndrome	ABCC2	Conjugated	Episodes of jaundice Bland cholestasis Dense brown lipofuscin

Dubin-Johnson syndrome



Acute cellular rejection

- Most common: 3 months after transplantation
- Sometimes due to reduced immunosuppression
- Histology (at least 2 of 3).
  1) Portal inflammation (activated lymphocytes, few eos, PMNs or plasma cells)
  2) Endotheliitis (portal veins or central veins).
  3) Bile duct lymphocytosis with injury

RAI score (max. score: 9)

- DD: Recurrent disease (hep B, C), autoimmune disease, drug effect, etc.

THANK YOU
Questions?

YOU GOT
THIS

1 WEEK BEFORE BOARDS, PRACTICE FOR **HIGH-YIELD ASSOCIATIONS** IN LIVER PATHOLOGY

TUMORAL

Liver fluke (*Clonorchis sinensis*) associated extra-hepatic cholangiocarcinoma: *TP53* and *SMAD4* mut

Small-duct intrahepatic cholangiocarcinoma (mass forming in periphery of liver): risk factors are non-biliary cirrhosis and chronic viral hepatitis): *IDH1/IDH2* mut and loss of BAP1.

Large-duct intrahepatic cholangiocarcinoma risk factors: PSC, hepatolithiasis, liver fluke infections.

Perihilar cholangiocarcinoma (Klatskin tumor): convergence of R and L hepatic ducts causing obstruction.

Combined HCC-cholangiocarcinoma: Both H&E and IHC evidence of both components needed for diagnosis. Hilar lymph node metastatic disease more common than HCC.

Lymphocyte-rich cholangiocarcinoma: + for EBV, more common in Asia.

Fibrolamellar carcinoma: microdeletion in cr.19 leading to *DNAJB1-PRKACA* fusion gene.

Angiosarcoma: exposure to androgen steroids, vinyl chloride exposure, arsenic and Thorotrast contrast.

Type 1 hepatic adenomas: *HNF1 alpha* mutated (loss of LFABP on IHC). Germline mutations in *HNF1 alpha* seen in mature onset diabetes of the young (MODY3).

Type 2 hepatic adenomas (HIGHEST RISK OF MALIGNANT TRANSFORMATION): *Beta-catenin mut* (on IHC: nuclear beta catenin and diffuse glutamine synthetase). Exon 3 mutations in *beta-catenin* (except S45 mutation) highest risk in comparison with Exon 7 or Exon 8 mutations.

Type 3 hepatic adenomas (inflammatory): JAK/STAT pathway activation via mut in *IL6ST* and *STAT3* (on IHC: + for CRP and SAA)

Adenomas with higher risk of bleeding: Inflammatory hepatocellular adenoma, sonic hedgehog-activated adenoma and argininosuccinate-synthase positive adenoma.

Epithelioid hemangioendothelioma (EHE): *WWTR1-CAMTA1* fusion gene. Less common YAP-1-TFE3 fusion.

Focal nodular hyperplasia (FNH): Map-like pattern with Glutamine synthetase stain

Aflatoxin and HCC: Mycotoxin produced by *Aspergillus flavus* (aflatoxin B1; potent carcinogen) can contaminate peanuts.

NON-TUMORAL

Sinusoidal lymphocytosis pattern (sinusoidal beading appearance with little hepatocyte injury: EBV hepatitis (EBV infects B lymphocytes), diphenylhydantoin (Dilantin), CMV, don't miss lymphoma.

Ground glass hepatocytes: Seen in subset of chronic hepatitis B. Virion particles in ER. HepB surface IHC cytoplasmic staining.

Sanded glass nuclei: Hep B with high viral replication. HepB core IHC positivity.

Alpha-1-antitrypsin deficiency: Mut in SERPINA1 gene (chr 14) (lead to accumulation of misfolded alpha-1-antitrypsin protein in hepatocytes). M allele is normal allele. S and Z most common alleles to cause disease (PiZZ is most severe)

Wilson disease: Mut in *ATPB7* gene (chr 13).

Hemochromatosis: Mutations in HFE gene (chr 6). 2 common mutations, C282Y and H63D. C282Y has highest iron overload. Patients at high risk of developing HCC.

Hypervitaminosis A: Stellate cell (AKA Ito cell) hyperplasia. Ito cells can lead to fibrosis in chronic injury states because they convert into myofibroblasts.

Glycogenic hepatopathy: Abundant pale cytoplasm (glycogen accumulation) + elevation in AST/ALT; complication of poorly controlled type 1 diabetes.

Amiodarone toxicity: Striking ballooned cells and Mallory hyaline.

Microvesicular steatosis: Acute fatty liver of pregnancy, Reye's syndrome, tetracycline, antiretrovirals.

Reye's syndrome: Encephalopathy and fatty degeneration of liver in children due to mitochondrial dysfunction after febrile viral-like illness treated with aspirin.

Punched out azonal necrosis: HSV, VZV and Adenovirus.

Zone 3 necrosis without inflammation: Ischemia, acetaminophen (treatment for acetaminophen toxicity: N-acetylcysteine, NAPQI is toxic metabolite), Budd Chiari, cocaine, mushroom poisoning.

Steatohepatitis required features for dx: Steatosis >5% + ballooned hepatocytes + lobular inflammation.

IgG elevated in autoimmune hepatitis; IgM elevated in PBC.

Granulomas in liver: PBC, sarcoidosis, CVID, systemic granulomatous disease, drugs (allopurinol), infections, idiopathic.

Fibrin ring granulomas: *Coxiella burneti* (Q-fever)

Echinococcosis:

- Cystic echinococcosis caused by *E. granulosus*. Individuals raising sheep. Cysts with 3 layers (pericyst, middle-laminated layer and innermost germinal layer. Hydatid sand: disintegrating proscoleces with hooklets, very antigenic).
- Alveolar echinococcosis: Caused by *E. multilocularis*. Contact with fox/coyotes. Necrotizing granulomatous inflammation. No well-formed cyst walls.

Schistosomiasis: larval forms of the parasite – released by freshwater snails – penetrate the skin during contact with infested water. Portal eosinophilic infiltrate with granulomas containing ova with lateral spine (S. mansoni) and dense portal fibrosis.

This page was intentionally left blank.



INFLAMMATION AND REPAIR

Justin C. Kerstetter, MD
Associate Professor of Pathology
Department of Pathology and Human Anatomy
Department of Dermatology
Loma Linda University Health

horror movie

Notes Prepared by Aaron Auerbach, MD

This lecture will cover aspects of the basic science chapters of Robbins

What do I need to know in these chapters of Robbins?



What is your diagnosis?



Interface dermatitis, lichenoid dermatitis,
superficial/deep perivascular and interstitial dermatitis?



Could be
- an interface drug reaction
- chronic/resolving arthropod bite reaction
- infection such as dermatophyte infection
- resolving lichenoid dermatitis (lichen planus, lichen stiatus)
- ?? Discoid lupus erythematosus
  - (if dermal mucin, and findings on direct immunoflourescence)
- early erythrema multiforme
  - (look for dyskeratotic/necrotic keratinocytes)
- ??urticaria
- + many other diagnoses



Early mycosis fungoides

inflammation

Associated with nearly every disease

Seen in nearly every slide

Different in different organs
    Lymphoid aggregates in the liver is abnormal,
    in the terminal ileum may be reactive Peyer patches

Associated with tumors
    Tumor infiltrating
    lymphocytes, melanoma



What is inflammation?

- Inflammation – is a protective response intended to eliminate the initial cause of cell injury and the necrotic cells and tissues arising from the injury

Inflammation

- Acute - minutes to days
  - Characterized by fluid and protein
  - PMN's
  - Exudate SG > 1.020
- Chronic - weeks to years
  - Lymphocytes and macrophages

Five classic local signs of acute inflammation

– Heat	– Calor – vasodilatation
– Redness	– Rubor – vasodilatation
– Swelling	– Tumor – vascular permeability
– Pain	– Dolor – mediator release/PMNs
– Loss of function	– Functio laesa – loss of function

Vascular changes

- Transient vasoconstriction
- Vasodilation
- Exudation of protein rich fluid
- Blood stasis
- Margination
- Emigration/Transmigration


A. NORMAL
No net flow
Net flow out
Net flow in
Arteriole
Capillaries
Venule
B. ACUTE INFLAMMATION
Net flow out
Net flow out
Net flow out
Arteriole
Capillaries
Venule
Hydrostatic pressure
Colloid osmotic pressure

Hydrostatic pressure is pressure outward exerted by a fluid at rest.
Osmotic pressure is pressure applied to prevent the inward flow of water across a semipermeable membrane


Rolling
Adhesion
Transmigration
Selectins
Integrins
Endothelial activation
Leukocyte activation
Stimulus
Chemotaxis, activation

R
A
T

S
I
T

Fig 3-6



Margination

- Normal flow - RBCs and WBCs flow in the center of the vessel
- A *cell poor* plasma is flowing adjacent to endothelium
- As blood flow slows, WBCs collect along the endothelium → Margination



Margination

- Normal flow - RBCs and WBCs flow in the center of the vessel
- A *cell poor* plasma is flowing adjacent to endothelium
- As blood flow slows, WBCs collect along the endothelium → Margination


Sialyl-Lewis X-modified glycoprotein
L-selectin
ENDOTHELIAL ACTIVATION
E-selectin (ELAM-1)
CD34
P-selectin (GMP-140)

Fig 3-9

Rolling

- Selectins transiently bind to receptors
- PMNs bounce or roll along → Rolling

Adhesion

- Mediated by integrins ICAM-1 and VCAM-1


Activated integrins
FIRM ADHESION
Sialyl-Lewis X-modified glycoprotein
PECAM-1 (CD31)
L-selectin
3
CD34
E-selectin (ELAM-1)
ICAM-1
P-selectin (GMP-140)

Transmigration

- Diapedesis (cells crawling)
- Mediated by PECAM-1 & ICAM-1 (Integrins)
- Primary in venules
- Collagenases degrade BM ↑ Permeability


TRANSMIGRATION
PECAM-1 (CD31)
4

Question 5



Question 5

- This electron micrograph illustrates…

A. Locomotion
B. Chemotaxis
C. Diapedesis
D. Pinocytosis
E. Margination

Answer C

- Leukocytes cross the endothelial lining of the capillary and migrate interstitially by the process called diapedesis or extravasation as illustrated in the image below.
- Chemotactic factors drive this process, particularly during an inflammatory response when immune-related cells migrate to and become concentrated at the site of inflammation.

Phagocytosis

1. Recognition & attachment
    Opsonins (IgG and C3) coat target
2. Engulfment
    Pseudopods flow around the particle to be engulfed. Particle is engulfed and fuses with lysosome
3. Killing/degradation
    - O_2 dep: Reactive O_2 species in lysosomes
    - O_2 indep: Bactericidal permeability agents, lysozyme, MBP, lactoferrin


1. RECOGNITION AND ATTACHMENT
Microbes bind to phagocyte receptors
Mannose receptor
Mac-1 integrin
Scavenger receptor
2. ENGULFMENT
Phagocyte membrane zips up around microbe
Microbe ingested in phagosome
Lysosome
Phagosome with ingested microbe
Lysosome with enzymes
Fusion of phagosome with lysosome
INOS
Arginine
NO
ROI
Phagocyte oxidase
O_2
Killing of microbes by ROIs and NO
Killing of microbes by lysosomal enzymes in phagolysosome
Phagolysosome
A
3. KILLING AND DEGRADATION

© Elsevier 2005

Question 3



The process illustrated in this EM is facilitated by…

- A. IgG and C3b
- B. bradykinin
- C. ICAM-1
- D. IL-1
- E. prostcyclin

Ans B. IgG & C3b

- This EM shows phagocytosis in which pseudopodia are seen embracing particles, which are then taken into the cell. The particles are coated by IgG antibodies and C3b, both of which are opsins.
- ICAM-1 is an integrin which is seen during adhesion. Bradykinin and prostacyclin are both vasodilators. Interleukin-1 acts to mediate the inflammatory response, in part by activating chemokines.

Boards like to test clinical situations

Genetic	Defect
LAD 1	•B chain of CD11/CD18 integrins
LAD 2	•Sialylated oligosaccharide
CGD	•Defective chemotaxis
X-linked—most common	•NADPH oxidase (membrane)
AR	•NADPH oxidase (cytoplasm)
MPO deficiency	•Absent H2O2 system, cant make HCL •no problem w infections
Chediak-Higashi syndrome	•Lysosomal defect
Acquired	
Thermal injury, DM, CA, sepsis	•Chemotaxis
Dialysis, DM	•Adhesion
Leukemia, anemia, sepsis, DM, neonates, malnutrition	•Phagocytosis & microbicidal activity

Leukocyte adhesion deficiency 1 (LAD-1)

- ↓ Beta chain of CD11/CD18 integrin on polys
- Neutrophils can roll but do not stick
- Recurrent bacterial infections
- Inflammatory lesions lack neutrophils
- But, ↑ neutrophils in the circulation

Chediak-Higashi Syndrome

- It's so rare you'll never see it in your practice, but it is often tested
- PB: WBC with multiple bluish inclusions per cell
- Defect in phagocytosis, no phagolysosome formation, no lysosome degranulation
- Defects in cells w granules
- melanosomes→oculocutaneous albinism
- Mutation in LYST lysosomal trafficking regulator gene
- EBV+ lymphoproliferative disorders



Large lysosome vesicles

Chronic Granulomatous Disease

- Usually x-linked, less often autosomal recessive
- Defect in NADPH oxidase system
- Young, men with infections and granulomas at autopsy
  - recurrent catalase + bacterial and fungal infections
- Test: Nitoblue tetrazolium reductase test negative
  - Inability to reduce NBT
  - ↓ O2 consumption, ↓ H2O2 and O2- production

Complement system

- Role in immunity
  - Membrane Attack Complex (MAC C5-9)
  - Punches a hole in bacterial membrane

If you have deficiencies in the MAC

you are predisposed to infections that are encapsulated

Neisseria, Strept, Hemoph, Listeria

Roles of complement in inflammation

1. Vascular effects (c3a & c5a)
   - ↑vascular permeability and vasodilation
   - Similar to histamine
2. Activates lipoxygenase pathway of arachidonic acid metabolism (c5a)
3. Leukocyte activation, adhesion and chemotaxis (c5a)
4. Phagocytosis
   - c3b acts as opsonin

Fig 3-16


CLASSIC PATHWAY
Antigen-antibody (IgG or IgM) complex
C1
Activated C1
C4 + C2
Classic pathway C3 convertase
C4b2a
Classic pathway C5 convertase
C4b2a3b
Also generated via plasmin or lysosomal proteases
C3a
C5a
C3
C3b
C5
C6 C7 C8 C9
C5b
C5-9
Membrane attack complex
C3
C3b
Microbial surfaces Polysaccharides
Factor B
Factor D
C3bBb
Alternative pathway C3 convertase
Stabilized by properdin
C3bBb3b
Alternative pathway C5 convertase
ALTERNATIVE PATHWAY

Complement related diseases

- Decay accelerating factor" (DAF) acts to destabilize the convertases
- Paroxysmal nocturnal hemoglobinuria occurs when DAF is not functional
  - -you cannot turn off the complement system
  - -RBCs are destroyed and platelets bind to complement
  - -Hemolytic anemia and thrombosis


Antigen-antibody (IgG or IgM) complex
CLASSIC PATHWAY
C1
Activated C1
C4 + C2
Classic pathway C3 convertase
C4b2a
Classic pathway C5 convertase
C4b2a3b
Also generated via plasmin or lysosomal proteases
C3a
C5a
C3
C3b
C5
C6 C7 C8 C9
C5b
C5-9
Membrane attack complex
C3
C3b
Microbial surfaces Polysaccharides
Factor B
Factor D
C3bBb
Alternative pathway C3 convertase
Stabilized by properdin
C3bBb3b
Alternative pathway C5 convertase
ALTERNATIVE PATHWAY

Arachidonic Acid

- AA is released from the cell membrane by phospholipases
- AA metabolism occurs via two pathways named for the enzymes that initiate the reactions; lipoxygenase and cyclooxygenase

AA metabolites (eicosanoids)

Cyclooxygenases synthesize

Prostaglandins

Thromboxanes

Lipoxygenases synthesize

Leukotrienes

Lipoxins


Cell membrane phospholipids
Phospholipases
Steroids inhibit
HETEs
HPETEs
Other lipoxygenases
ARACHIDONIC ACID
5-Lipoxygenase
Cyclooxygenase
COX-1 and COX-2 inhibitors aspirin, indomethacin inhibit
5-HETE
5-HPETE
Chemotaxis
12-Lipoxygenase
Prostaglandin G₂ (PGG₂)
Leukotriene B₄
Leukotriene A₄ (LTA₄)
Prostaglandin H₂ (PGH₂)
Leukotriene C₄ (LTC₄)
Leukotriene D₄ (LTD₄)
Leukotriene E₄ (LTE₄)
Vasoconstriction Bronchospasm Increased permeability
Prostacyclin PGI₂
Causes vasodilation, inhibits platelet aggregation
Thromboxane A₂ TXA₂
Causes vasoconstriction, promotes platelet aggregation
Lipoxin A₄ (LXA₄)
Lipoxin B₄ (LXB₄)
Vasodilation Inhibit neutrophil chemotaxis Stimulate monocyte adhesion
PGD₂
PGE₂
PGF₂α
Vasodilation Potentiate edema

© Elsevier 2005

IL1/TNF cytokines


Bacterial products, immune complexes, toxins, physical injury, other cytokines
MACROPHAGE (and other cell) ACTIVATION
IL-1 / TNF

ACUTE-PHASE REACTIONS

Fever
↑Sleep
↓Appetite
↑Acute-phase proteins
Hemodynamic effects (shock)
Neutrophilia

ENDOTHELIAL EFFECTS

↑Leukocyte adherence
↑PGI synthesis
↑Procoagulant activity
↓Anticoagulant activity
↑IL-1, IL-8, IL-6, PDGF

FIBROBLAST EFFECTS

↑Proliferation
↑Collagen synthesis
↑Collagenase
↑Protease
↑PGE synthesis

LEUKOCYTE EFFECTS

↑Cytokine secretion (IL-1, IL-6)

© Elsevier 2005

Chronic inflammation

- Difficult to define
- Inflammation of prolonged duration in which active inflammation, tissue injury and the healing proceed simultaneously
- Examples
  - Persistent infections - AFB, fungi, Treponemes
    - Delayed hypersensitivity
    - Granulomatous inflammation
  - Prolonged exposure potentially toxic agents
    - Silica
  - Autoimmunity - RA, lupus
  - atherosclerosis

Chronic inflammation

- Infiltration with mononuclear cells (macrophages, lymphocytes & plasma cells)
- Tissue destruction
- Repair involving angiogenesis and fibrosis
- Macrophage is the main player

Macrophage

Tox. O_2
Proteases
Collagenases
Chemotx factors
Coag factors
AA metabolites
NO PDGF
FGF TGF-β

Key macrophage events

1. Recruitment from circulation
2. Are immobilized at the site of inflammation
3. Differentiation (microglia, kupffer, alveolar macrophage, osteoclasts)

Granulomatous Inflammation

- Aggregations of epithelioid histiocytes
- Bacteria
  - Tuberculosis
- Parasites
  - Schistosomiasis
- Wegners disease
- Prostate
  - BCG, Post Turp
- Fungi
  - Histoplasmosis
- Metal/Dust
  - Berylliosis
  - Silicosis
- Foreign body
  - Splinter
  - Suture
  - Graft material
- Sarcoidosis

Granuloma



Question 8



Yes this is a really dark picture just like the boards

Question 8

- This lesion is typical of disease characterized by which of the following?

A. A deficiency of hydrogen peroxide generation
B. Infections due to catalase-producing microorganisms
C. Occurrence mainly in males
D. All of the above are correct
E. None of the above is correct

Answer D

- Chronic granulomatous disease is due to defects in the oxidative burst in leukocytes. The most common form is X-liked caused by a defect in the membrane component of the NADPH oxidase. This is most commonly seen in boys. The patients have increased susceptability of catalase producing bacteria.

Books that I like for board review

- Rosai (for the pictures)
- Leftkowitz (AP board review) for questions
- The clinical compendium for CP
- Random other atlases

Questions?

Questions?

This page was intentionally left blank.